

PACKING THE HYPERCUBE

DAVID OFFNER

*Department of Mathematics and Computer Science
Westminster College, Pennsylvania, USA*

e-mail: offnerde@westminster.edu

Abstract

Let G be a graph that is a subgraph of some n -dimensional hypercube Q_n . For sufficiently large n , Stout [20] proved that it is possible to pack vertex-disjoint copies of G in Q_n so that any proportion $r < 1$ of the vertices of Q_n are covered by the packing. We prove an analogous theorem for edge-disjoint packings: For sufficiently large n , it is possible to pack edge-disjoint copies of G in Q_n so that any proportion $r < 1$ of the edges of Q_n are covered by the packing.

Keywords: hypercube, packing, decomposition.

2010 Mathematics Subject Classification: 05C88.

REFERENCES

- [1] W. Aiello and F.T. Leighton, *Hamming codes, hypercube embeddings, and fault tolerance*, SIAM J. Comput. **37** (2007) 783–803.
doi:10.1137/S0097539798332464
- [2] B. Barden, R. Libeskind-Hadas, J. Davis, and W. Williams, *On edge-disjoint spanning trees in hypercubes*, Inform. Process. Lett. **70** (1999) 13–16.
doi:10.1016/S0020-0190(99)00033-2
- [3] D. Bryant, S. El-Zanati, C. Vanden Eynden, and D. Hoffman, *Star decompositions of cubes*, Graphs Combin. **17** (2001) 55–59.
doi:10.1007/s003730170054
- [4] G. Cybenko, D.W. Krumme, and K.N. Venkataraman, *Fixed hypercube embedding*, Inform. Process. Lett. **25** (1987) 35–39.
doi:10.1016/0020-0190(87)90090-1
- [5] M.E. Deza, *On Hamming geometry of unitary cubes*, Cybernetics and Control Theory, Soviet Phys. Dokl. (1961) 940–943.

- [6] D.Ž. Djoković, *Distance-preserving subgraphs of hypercubes*, J. Combin. Theory (B) **14** (1973) 263–267.
doi:10.1016/0095-8956(73)90010-5
- [7] A. Felzenbaum, R. Holzman, and D. Kleitman, *Packing lines in a hypercube*, Discrete Math. **117** (1993) 107–112.
doi:10.1016/0095-8956(73)90010-5
- [8] J.F. Fink, *On the decomposition of n -cubes into isomorphic trees*, J. Graph Theory **14** (1990) 405–411.
doi:10.1002/jgt.3190140403
- [9] M.R. Garey and R.L. Graham, *On cubical graphs*, J. Combin. Theory (B) **18** (1975) 84–95.
doi:10.1016/0095-8956(75)90067-2
- [10] V.A. Gorbatov and A.A. Kazanskiy, *Characterization of graphs embedded in n -cubes*, Engineering Cybernetics, Soviet J. Comput. Systems Sci. **20** (1982) 96–102.
- [11] N. Graham and F. Harary, *Covering and packing in graphs—V: Mispacking subcubes in hypercubes*, Comput. Math. Appl. **15** (1988) 267–270.
doi:10.1016/0898-1221(88)90211-8
- [12] F. Harary, J. Hayes, and H. Wu, *A survey of the theory of hypercube graphs*, Comput. Math. Appl. **15** (1988) 277–289.
doi:10.1016/0898-1221(88)90213-1
- [13] P. Horak, J. Širáň, and W. Wallis, *Decomposing cubes*, J. Aust. Math. Soc. (A) **61** (1996) 119–128.
doi:10.1017/S1446788700000112
- [14] F.T. Leighton, *Introduction to Parallel Algorithms and Architectures: Arrays, Trees, Hypercubes* (Morgan Kaufmann Publishers, 1991).
- [15] M. Livingston and Q. Stout, *Embeddings in hypercubes*, Math. Comput. Modelling **11** (1988) 222–227.
doi:10.1016/0895-7177(88)90486-4
- [16] E.M. Palmer, *On the spanning tree packing number of a graph: a survey*, Discrete Math. **230** (2001) 13–21.
doi:10.1016/S0012-365X(00)00066-2
- [17] M. Ramras, *Symmetric edge-decompositions of hypercubes*, Graphs Combin. **7** (1991) 65–87.
doi:10.1007/BF01789464
- [18] M. Ramras, *Symmetric vertex partitions of hypercubes by isometric trees*, J. Combin. Theory (B) **54** (1992) 239–248.
doi:10.1016/0095-8956(92)90055-3
- [19] M. Ramras, *Fundamental subsets of edges of hypercubes*, Ars Combin. **46** (1997) 3–24.

- [20] Q. Stout, *Packings in hypercubes*, presented at 21st Southeastern Int'l. Conf. on Combinatorics, Graph Theory, and Computing, Boca Raton, FL, (1990).
<http://www.eecs.umich.edu/~qstout/abs/hypack.html>.
- [21] J.H. van Lint, Introduction to Coding Theory, 3rd Edition (Springer, 1998).
- [22] S. Wagner and M. Wild, *Decomposing the hypercube Q_n into n isomorphic edge-disjoint trees*, Discrete Math. **312** (2012) 1819–1822.
doi:10.1016/j.disc.2012.01.033
- [23] R. Wilson, *Decompositions of complete graphs into subgraphs isomorphic to a given graph*, Congr. Numer. **15** (1976) 647–659.

Received 4 February 2012

Revised 30 November 2012

Accepted 3 December 2012