

NOTE

A DIFFERENT SHORT PROOF OF BROOKS' THEOREM

LANDON RABERN

Arizona State University
School of Mathematical & Statistical Sciences
e-mail: landon.rabern@gmail.com

Abstract

Lovász gave a short proof of Brooks' theorem by coloring greedily in a good order. We give a different short proof by reducing to the cubic case.

Keywords: coloring, clique number, maximum degree.

2010 Mathematics Subject Classification: 05C15.

REFERENCES

- [1] R.L. Brooks, *On colouring the nodes of a network*, in: Math. Proc. Cambridge Philos. Soc. **37** Cambridge Univ. Press (1941) 194–197.
- [2] R. Diestel, Graph Theory (Fourth Ed., Springer Verlag, 2010).
- [3] A.D. King, *Hitting all maximum cliques with a stable set using lopsided independent transversals*, J. Graph Theory **67** (2011) 300–305.
doi:10.1002/jgt.20532
- [4] A.V. Kostochka, *Degree, density, and chromatic number*, Metody Diskret. Anal. **35** (1980) 45–70 (in Russian).
- [5] L. Lovász, *Three short proofs in graph theory*, J. Combin. Theory (B) **19** (1975) 269–271.
doi:10.1016/0095-8956(75)90089-1
- [6] L. Rabern, *On hitting all maximum cliques with an independent set*, J. Graph Theory **66** (2011) 32–37.
doi:10.1002/jgt.20487
- [7] H. Tverberg, *On Brooks' theorem and some related results*, Math. Scand. **52** (1983) 37–40.

Received 28 May 2012
Accepted 3 January 2013