

NOTE

PACKING TREES INTO n -CHROMATIC GRAPHS

ANDRÁS GYÁRFÁS

Alfréd Rényi Institute of Mathematics

e-mail: gyarf@renyi.hu

Abstract

We show that if a sequence of trees T_1, T_2, \dots, T_{n-1} can be packed into K_n then they can be also packed into any n -chromatic graph.

Keywords: tree packing.

2010 Mathematics Subject Classification: 05C70.

REFERENCES

- [1] D. Gerbner, B. Keszegh and C. Palmer, *Generalizations of the tree packing conjecture*, Discuss. Math. Graph Theory **32** (2012) 569–582.
doi:10.7151/dmgt.1628
- [2] D. Gerbner, B. Keszegh and C. Palmer, *Red-blue alternating paths*, Third Emléktábla Workshop, p.25.
<http://www.renyi.hu/~emlektab/index/booklet.html>
- [3] A. Gyárfás and J. Lehel, *Packing trees of different order into K_n* , Combinatorics, Proc. Fifth Hungarian Coll. Keszthely, 1976, Vol II. North Holland. Colloq. Math. Soc. J. Bolyai **18** 463–469.
- [4] A. Gyárfás, E. Szemerédi and Zs. Tuza, *Induced subtrees in graphs of large chromatic number*, Discrete Math. **30** (1980) 235–244.
doi:10.1016/0012-365X(80)90230-7
- [5] S. Zaks and C.L. Liu, *Decomposition of graphs into trees*, Proceedings of 8-th South-eastern Conference on Combinatorics, Graph Theory and Computing, Louisiana State Univ., Baton Rouge, La. Util. Math., Congr. Numer. **XIX** (1977) 643–654.

Received 13 June 2013
Accepted 26 June 2013