

NOTE

FRACTIONAL ASPECTS OF THE ERDŐS-FABER-LOVÁSZ CONJECTURE

JOHN BOSICA AND CLAUDE TARDIF

Department of Mathematics and Computer Science
Royal Military College of Canada
PO Box 17000 Stn Forces, Kingston, ON Canada, K7K 7B4

e-mail: John.Bosica@rmc.ca
Claude.Tardif@rmc.ca

Abstract

The Erdős-Faber-Lovász conjecture is the statement that every graph that is the union of n cliques of size n intersecting pairwise in at most one vertex has chromatic number n . Kahn and Seymour proved a fractional version of this conjecture, where the chromatic number is replaced by the fractional chromatic number. In this note we investigate similar fractional relaxations of the Erdős-Faber-Lovász conjecture, involving variations of the fractional chromatic number. We exhibit some relaxations that can be proved in the spirit of the Kahn-Seymour result, and others that are equivalent to the original conjecture.

Keywords: Erdős-Faber-Lovász Conjecture, fractional chromatic number.

2010 Mathematics Subject Classification: 05C15, 05C72.

REFERENCES

- [1] N.G. de Bruijn and P. Erdős, *On a combinatorial problem*, *Nederl. Akad. Wetensch. Indag. Math* **10** (1948) 421–423.
- [2] P. Erdős, R.C. Mullin, V.T. Sós and D.R. Stinson, *Finite linear spaces and projective planes*, *Discrete Math.* **47** (1983) 49–62.
doi:10.1016/0012-365X(83)90071-7
- [3] J. Kahn, *Coloring nearly-disjoint hypergraphs with $n + o(n)$ colors*, *J. Combin. Theory (A)* **59** (1992) 31–39.
doi:10.1016/0097-3165(92)90096-D
- [4] J. Kahn and P.D. Seymour, *A fractional version of the Erdős-Faber-Lovász conjecture*, *Combinatorica* **12** (1992) 155–160.
doi:10.1007/BF01204719

- [5] E.R. Scheinerman and D.H. Ullman, Fractional Graph Theory (Wiley-Interscience Series in Discrete Mathematics and Optimization, John Wiley & Sons, New York, 1997).

Received 22 May 2013
Revised 28 November 2013
Accepted 20 February 2014