

A NOTE ON LONGEST PATHS IN CIRCULAR ARC GRAPHS

FELIX JOOS

Institut für Optimierung und Operations Research
Universität Ulm, Ulm, Germany

e-mail: felix.joos@uni-ulm.de

Abstract

As observed by Rautenbach and Sereni [SIAM J. Discrete Math. **28** (2014) 335–341] there is a gap in the proof of the theorem of Balister *et al.* [Combin. Probab. Comput. **13** (2004) 311–317], which states that the intersection of all longest paths in a connected circular arc graph is nonempty. In this paper we close this gap.

Keywords: circular arc graphs, longest paths intersection.

2010 Mathematics Subject Classification: 05C38.

REFERENCES

- [1] P.N. Balister, E. Györi, J. Lehel and R.H. Schelp, *Longest paths in circular arc graphs*, *Combin. Probab. Comput.* **13** (2004) 311–317.
doi:10.1017/S0963548304006145
- [2] T. Gallai, *Problem 4*, in: *Theory of graphs*, Proceedings of the Colloquium held at Tihany, Hungary, September, 1966., P. Erdős and G. Katona Eds., Academic Press, New York-London; Akademiai Kiad, Budapest (1968).
- [3] J.M. Keil, *Finding Hamiltonian circuits in interval graphs*, *Inform. Process. Lett.* **20** (1985) 201–206.
doi:10.1016/0020-0190(85)90050-X
- [4] D. Rautenbach and J.-S. Sereni, *Transversals of longest paths and cycles*, *SIAM J. Discrete Math.* **28** (2014) 335–341.
doi:10.1137/130910658
- [5] A. Shabbira, C.T. Zamfirescu and T.I. Zamfirescu, *Intersecting longest paths and longest cycles: A survey*, *Electron. J. Graph Theory Appl.* **1** (2013) 56–76.
doi:10.5614/ejgta.2013.1.1.6

Received 16 December 2013

Revised 6 August 2014

Accepted 6 August 2014