

ON SPECTRA OF VARIANTS OF THE CORONA OF TWO GRAPHS AND SOME NEW EQUIENERGETIC GRAPHS

CHANDRASHEKAR ADIGA AND B.R. RAKSHITH

*Department of Studies in Mathematics
 University of Mysore, Manasagangothri
 Mysore – 570 006, India*

e-mail: c.adiga@hotmail.com
 ranmsc08@yahoo.co.in

Abstract

Let G and H be two graphs. The join $G \vee H$ is the graph obtained by joining every vertex of G with every vertex of H . The corona $G \circ H$ is the graph obtained by taking one copy of G and $|V(G)|$ copies of H and joining the i -th vertex of G to every vertex in the i -th copy of H . The neighborhood corona $G \star H$ is the graph obtained by taking one copy of G and $|V(G)|$ copies of H and joining the neighbors of the i -th vertex of G to every vertex in the i -th copy of H . The edge corona $G \diamond H$ is the graph obtained by taking one copy of G and $|E(G)|$ copies of H and joining each terminal vertex of i -th edge of G to every vertex in the i -th copy of H . Let G_1, G_2, G_3 and G_4 be regular graphs with disjoint vertex sets. In this paper we compute the spectrum of $(G_1 \vee G_2) \cup (G_1 \star G_3)$, $(G_1 \vee G_2) \cup (G_2 \star G_3) \cup (G_1 \star G_4)$, $(G_1 \vee G_2) \cup (G_1 \circ G_3)$, $(G_1 \vee G_2) \cup (G_2 \circ G_3) \cup (G_1 \circ G_4)$, $(G_1 \vee G_2) \cup (G_1 \diamond G_3)$, $(G_1 \vee G_2) \cup (G_2 \diamond G_3) \cup (G_1 \diamond G_4)$, $(G_1 \vee G_2) \cup (G_2 \circ G_3) \cup (G_1 \star G_3)$, $(G_1 \vee G_2) \cup (G_2 \circ G_3) \cup (G_1 \diamond G_4)$ and $(G_1 \vee G_2) \cup (G_2 \star G_3) \cup (G_1 \diamond G_4)$. As an application, we show that there exist some new pairs of equienergetic graphs on n vertices for all $n \geq 11$.

Keywords: spectrum, corona, neighbourhood corona, edge corona, energy of a graph, equienergetic graphs.

2010 Mathematics Subject Classification: 05C50.

REFERENCES

- [1] C. Adiga, R. Balakrishnan and W. So, *The skew energy of a digraph*, Linear Algebra Appl. **432** (2010) 1825–1835.
 doi:10.1016/j.laa.2009.11.034

- [2] R. Balakrishnan, *The energy of a graph*, Linear Algebra Appl. **387** (2004) 287–295.
doi:10.1016/j.laa.2004.02.038
- [3] R.B. Bapat, *Energy of a graph is never an odd integer*, Bull. Kerala Math. Assoc. **1** (2004) 129–132.
- [4] S. Barik, S. Pati and B.K. Sarma, *The spectrum of the corona of two graphs*, SIAM J. Discrete Math. **21** (2007) 47–56.
doi:10.1137/050624029
- [5] S.B. Bozkurt, A.D. Gungor and I. Gutman, *Note on distance energy of graphs*, MATCH. Commun. Math. Comput. Chem. **64** (2010) 129–134.
- [6] V. Brankov, D. Stevanović and I. Gutman, *Equienergetic chemical trees*, J. Serb. Chem. Soc. **69** (2004) 549–553.
doi:10.2298/JSC0407549B
- [7] S.-Y. Cui and G.-X. Tian, *The spectrum and the signless Laplacian spectrum of coronae*, Linear Algebra Appl. **437** (2012) 1692–1703.
doi:10.1016/j.laa.2012.05.019
- [8] D. Cvetković, M. Doob and H. Sachs, *Spectra of Graphs: Theory and Application* (Academic Press, New York, 1980).
- [9] W.L. Ferrar, *A Text-Book of Determinants, Matrices and Algebraic Forms* (Oxford University Press, 1953).
- [10] R. Frucht and F. Harary, *On the corona of two graphs*, Aequationes Math. **4** (1970) 322–325.
doi:10.1007/BF01844162
- [11] S. Gong, X. Li, G. Xu, I. Gutman and B. Furtula, *Borderenergetic graphs*, MATCH Commun. Math. Comput. Chem. **74** (2015) 321–332.
- [12] I. Gutman, *The energy of a graph*, Ber. Math.-Statist. Sect. Forschungsz. Graz **103** (1978) 1–22.
- [13] I. Gutman, *The energy of a graph: old and new results*, in: Algebraic Combinatorics and Applications, A. Betten, A. Kohnert, R. Laue and A. Wassermann (Ed(s)), (Berlin, Springer, 2000) 196–211.
- [14] I. Gutman, *Topology and stability of conjugated hydrocarbons. The dependence of total π -electron energy on molecular topology*, J. Serb. Chem. Soc. **70** (2005) 441–456.
doi:10.2298/JSC0503441G
- [15] I. Gutman, D. Kiani, M. Mirzakhah and B. Zhou, *On incidence energy of a graph*, Linear Algebra Appl. **431** (2009) 1223–1233.
doi:10.1016/j.laa.2009.04.019
- [16] Y. Hou and W.-C. Shiu, *The spectrum of the edge corona of two graphs*, Electron. J. Linear Algebra **20** (2010) 586–594.
doi:10.13001/1081-3810.1395

- [17] G. Indulal, *The spectrum of neighborhood corona of graphs*, Kragujevac J. Math. **35** (2011) 493–500.
- [18] G. Indulal and A. Vijayakumar, *On a pair of equienergetic graphs*, MATCH. Commun. Math. Comput. Chem. **55** (2006) 83–90.
- [19] X. Li, Y. Shi and I. Gutman, *Graph Energy* (Springer, New York, 2012).
doi:10.1007/978-1-4614-4220-2
- [20] X. Li, M. Wei and S. Gong, *A computer search for the borderenergetic graphs of order 10*, MATCH Commun. Math. Comput. Chem. **74** (2015) 333–342.
- [21] B. Liu, Y. Huang and Z. You, *A survey on the Laplacian-energy-like invariant*, MATCH. Commun. Math. Comput. Chem. **66** (2011) 713–730.
- [22] J. Liu and B. Liu, *On a pair of equienergetic graphs*, MATCH. Commun. Math. Comput. Chem. **59** (2008) 275–278.
- [23] X. Liu and S. Zhou, *Spectra of the neighbourhood corona of two graphs*, Linear Multilinear Algebra **62** (2014) 1205–1219.
doi:10.1080/03081087.2013.816304
- [24] C. McLeman and E. McNicholas, *Spectra of coronae*, Linear Algebra Appl. **435** (2011) 998–1007.
doi:10.1016/j.laa.2011.02.007
- [25] H.S. Ramane, I. Gutman, H.B. Walikar and S.B. Halkarni, *Equienergetic complement graphs*, Kragujevac J. Sci. **27** (2005) 67–74.
- [26] H.S. Ramane and H.B. Walikar, *Construction of equienergetic graphs*, MATCH Commun. Math. Comput. Chem. **57** (2007) 203–210.
- [27] D. Stevanović, *Energy and NEPS of graphs*, Linear Multilinear Algebra **53** (2005) 67–74.
doi:10.1080/03081080410001714705
- [28] D. Stevanović and I. Stanković, *Remarks on hyperenergetic circulant graphs*, Linear Algebra Appl. **400** (2005) 345–348.
doi:10.1016/j.laa.2005.01.001
- [29] L. Xu and Y. Hou, *Equienergetic bipartite graphs*, MATCH Commun. Math. Comput. Chem. **57** (2007) 363–370.

Received 29 April 2015

Revised 25 May 2015

Accepted 25 May 2015