

## FORBIDDEN SUBGRAPHS FOR HAMILTONICITY OF 1-TOUGH GRAPHS

BINLONG LI

*Department of Applied Mathematics, School of Science  
Northwestern Polytechnical University  
Xi'an, Shaanxi 710072, P.R. China, and  
European Centre of Excellence NTIS  
306 14 Pilsen, Czech Republic*  
**e-mail:** libinlong@mail.nwpu.edu.cn

HAJO J. BROERSMA

*Faculty of EEMCS, University of Twente  
7500 AE Enschede, The Netherlands*  
**e-mail:** h.j.broersma@utwente.nl

AND

SHENGGUI ZHANG

*Department of Applied Mathematics, School of Science  
Northwestern Polytechnical University  
Xi'an, Shaanxi 710072, P.R. China*  
**e-mail:** sgzhang@nwpu.edu.cn

### Abstract

A graph  $G$  is said to be 1-tough if for every vertex cut  $S$  of  $G$ , the number of components of  $G - S$  does not exceed  $|S|$ . Being 1-tough is an obvious necessary condition for a graph to be hamiltonian, but it is not sufficient in general. We study the problem of characterizing all graphs  $H$  such that every 1-tough  $H$ -free graph is hamiltonian. We almost obtain a complete solution to this problem, leaving  $H = K_1 \cup P_4$  as the only open case.

**Keywords:** forbidden subgraph, 1-tough graph,  $H$ -free graph, hamiltonian graph.

**2010 Mathematics Subject Classification:** 05C45.

## REFERENCES

- [1] D. Bauer, H.J. Broersma, J. van den Heuvel and H.J. Veldman, *Long cycles in graphs with prescribed toughness and minimum degree*, Discrete Math. **141** (1995) 1–10.  
doi:10.1016/0012-365X(93)E0204-H
- [2] D. Bauer, H.J. Broersma and E. Schmeichel, *Toughness in graphs—a survey*, Graphs Combin. **22** (2006) 1–35.  
doi:10.1007/s00373-006-0649-0
- [3] D. Bauer, H.J. Broersma and H.J. Veldman, *Not every 2-tough graph is hamiltonian*, Discrete Appl. Math. **99** (2000) 317–321.  
doi:10.1016/S0166-218X(99)00141-9
- [4] P. Bedrossian, *Forbidden Subgraph and Minimum Degree Conditions for Hamiltonicity* (Ph.D. Thesis, Memphis State University, 1991).
- [5] J.A. Bondy and U.S.R. Murty, *Graph Theory* (Graduate Texts in Mathematics, Springer **244**, 2008).
- [6] H.J. Broersma, *On some intriguing problems in hamiltonian graph theory—a survey*, Discrete Math. **251** (2002) 47–69.  
doi:10.1016/S0012-365X(01)00325-9
- [7] H.J. Broersma, V. Patel, and A. Pyatkin, *On toughness and hamiltonicity of  $2K_2$ -free graphs*, J. Graph Theory **75** (2014) 244–255.  
doi:10.1002/jgt.21734
- [8] V. Chvátal, *Tough graphs and hamiltonian circuits*, Discrete Math. **5** (1973) 215–228.  
doi:10.1016/0012-365X(73)90138-6
- [9] R.J. Faudree and R.J. Gould, *Characterizing forbidden pairs for hamiltonian properties*, Discrete Math. **173** (1997) 45–60.  
doi:10.1016/S0012-365X(96)00147-1
- [10] H.A. Jung, *On a class of posets and the corresponding comparability graphs*, J. Combin. Theory, Ser. B **24** (1978) 125–133.  
doi:10.1016/0095-8956(78)90013-8
- [11] M.M. Matthews and D.P. Sumner, *Hamiltonian results in  $K_{1,3}$ -free graphs*, J. Graph Theory **8** (1984) 139–146.  
doi:10.1002/jgt.3190080116
- [12] Z.G. Nikoghosyan, *Disconnected forbidden subgraphs, toughness and Hamilton cycles*, ISRN Combinatorics **2013** (2013) ID 673971.  
doi:10.1155/2013/673971

Received 18 March 2015

Revised 6 January 2016

Accepted 7 January 2016