

SHARP UPPER BOUNDS ON THE SIGNLESS LAPLACIAN SPECTRAL RADIUS OF STRONGLY CONNECTED DIGRAPHS¹

WEIGE XI AND LIGONG WANG²

*Department of Applied Mathematics
 School of Science, Northwestern Polytechnical University
 Xi'an, Shaanxi 710072, P.R. China*

e-mail: xiyanxwg@163.com
 lgwangmath@163.com

Abstract

Let $G = (V(G), E(G))$ be a simple strongly connected digraph and $q(G)$ be the signless Laplacian spectral radius of G . For any vertex $v_i \in V(G)$, let d_i^+ denote the outdegree of v_i , m_i^+ denote the average 2-outdegree of v_i , and N_i^+ denote the set of out-neighbors of v_i . In this paper, we prove that:

(1) $q(G) = d_1^+ + d_2^+$, ($d_1^+ \neq d_2^+$) if and only if G is a star digraph $\overset{\longleftrightarrow}{K}_{1,n-1}$, where d_1^+, d_2^+ are the maximum and the second maximum outdegree, respectively ($\overset{\longleftrightarrow}{K}_{1,n-1}$ is the digraph on n vertices obtained from a star graph $K_{1,n-1}$ by replacing each edge with a pair of oppositely directed arcs).

(2) $q(G) \leq \max \left\{ \frac{1}{2} \left(d_i^+ + \sqrt{d_i^{+2} + 8d_i^+ m_i^+} \right) : v_i \in V(G) \right\}$ with equality if and only if G is a regular digraph.

(3) $q(G) \leq \max \left\{ \frac{1}{2} \left(d_i^+ + \sqrt{d_i^{+2} + \frac{4}{d_i^+} \sum_{v_j \in N_i^+} d_j^+ (d_j^+ + m_j^+)} \right) : v_i \in V(G) \right\}$.

Moreover, the equality holds if and only if G is a regular digraph or a bipartite semiregular digraph.

(4) $q(G) \leq \max \left\{ \frac{1}{2} \left(d_i^+ + 2d_j^+ - 1 + \sqrt{(d_i^+ - 2d_j^+ + 1)^2 + 4d_i^+} \right) : (v_j, v_i) \in E(G) \right\}$. If the equality holds, then G is a regular digraph or $G \in \Omega$, where Ω is a class of digraphs defined in this paper.

Keywords: digraph, signless Laplacian spectral radius.

2010 Mathematics Subject Classification: 05C50, 15A18.

¹The research was supported by the National Natural Science Foundation of China (Grant No. 11171273).

²Corresponding author.

REFERENCES

- [1] Ş.B. Bozkurt and D. Bozkurt, *On the signless Laplacian spectral radius of digraphs*, Ars Combin. **108** (2013) 193–200.
- [2] B.K. Bulter and P.H. Siegel, *Sharp bounds on the spectral radius of nonnegative matrices and digraphs*, Linear Algebra Appl. **439** (2013) 1468–1478.
doi:10.1016/j.laa.2013.04.029
- [3] Y.Q. Chen and L.G. Wang, *Sharp bounds for the largest eigenvalue of the signless Laplacian of a graph*, Linear Algebra Appl. **433** (2010) 908–913.
doi:10.1016/j.laa.2010.04.026
- [4] S.Y. Cui, C.X. Tian and J.J. Guo, *A sharp upper bound on the signless Laplacian spectral radius of graphs*, Linear Algebra Appl. **439** (2013) 2442–2447.
doi:10.1016/j.laa.2013.06.015
- [5] K.Ch. Das, *A characterization on graphs which achieve the upper bound for the largest Laplacian eigenvalue of graphs*, Linear Algebra Appl. **376** (2004) 173–186.
doi:10.1016/j.laa.2003.06.009
- [6] K.Ch. Das, *The Laplacian spectrum of a graph*, Comput. Math. Appl. **48** (2004) 715–724.
doi:10.1016/j.camwa.2004.05.005
- [7] K.Ch. Das, *Maximizing the sum of the squares of the degrees of a graph*, Discrete Math. **285** (2004) 57–66.
doi:10.1016/j.disc.2004.04.007
- [8] E. Gudiño and J. Rada, *A lower bound for the spectral radius of a digraph*, Linear Algebra Appl. **433** (2010) 233–240.
doi:10.1016/j.laa.2010.02.012
- [9] A.D. Güngör, K.C. Das and A.S. Çevik, *Sharp upper bounds on the spectral radius of the signless Laplacian matrix of a graph*, Appl. Math. Comput. **219** (2013) 5025–5032.
doi:10.1016/j.amc.2012.11.039
- [10] W.X. Hong and L.H. You, *Spectral radius and signless Laplacian spectral radius of strongly connected digraphs*, Linear Algebra Appl. **457** (2014) 93–113.
doi:10.1016/j.laa.2014.05.007
- [11] R.A. Horn and C.R. Johnson, *Matrix Analysis* (Cambridge University Press, New York, 1985).
doi:10.1017/CBO9780511810817
- [12] W.W. Lang and L.G. Wang, *Sharp bounds for the signless Laplacian spectral radius of digraphs*, Appl. Math. Comput. **238** (2014) 43–49.
doi:10.1016/j.amc.2014.04.001
- [13] M.H. Liu, B.L. Liu and B. Cheng, *Ordering (signless) Laplacian spectral radii with maximum degrees of graphs*, Discrete Math. **338** (2015) 159–163.
doi:10.1016/j.disc.2014.09.007

- [14] C.S. Oliveira, L.S. de Lima, N.M.M. de Abreu and P. Hansen, *Bounds on the index of the signless Laplacian of a graph*, Discrete Appl. Math. **158** (2010) 355–360.
doi:10.1016/j.dam.2009.06.023
- [15] Y.L. Pan, *Sharp upper bounds for the Laplacian graph eigenvalues*, Linear Algebra Appl. **355** (2002) 287–295.
doi:10.1016/S0024-3795(02)00353-1
- [16] G.H. Xu, K.F. Fang and J. Shen, *Bounds on the spectral radii of digraphs in terms of walks*, Appl. Math. Comput. **219** (2012) 3721–3728.
doi:10.1016/j.amc.2012.09.073
- [17] G.L. Yu, Y.R. Wu and J.L. Shu, *Sharp bounds on the signless Laplacian spectral radii of graphs*, Linear Algebra Appl. **434** (2011) 683–687.
doi:10.1016/j.laa.2010.09.029

Received 26 March 2015
Revised 13 January 2016
Accepted 13 January 2016