

ONE MORE TURÁN NUMBER AND RAMSEY NUMBER FOR THE LOOSE 3-UNIFORM PATH OF LENGTH THREE

JOANNA POLCYN

*Adam Mickiewicz University
Poznań, Poland*

e-mail: joaska@amu.edu.pl

Abstract

Let P denote a 3-uniform hypergraph consisting of 7 vertices a, b, c, d, e, f, g and 3 edges $\{a, b, c\}$, $\{c, d, e\}$, and $\{e, f, g\}$. It is known that the r -color Ramsey number for P is $R(P; r) = r + 6$ for $r \leq 9$. The proof of this result relies on a careful analysis of the Turán numbers for P . In this paper, we refine this analysis further and compute the fifth order Turán number for P , for all n . Using this number for $n = 16$, we confirm the formula $R(P; 10) = 16$.

Keywords: Ramsey numbers, Turán numbers.

2010 Mathematics Subject Classification: 05D10.

REFERENCES

- [1] R. Csákány and J. Kahn, *A homological approach to two problems on finite sets*, J. Algebraic Combin. **9** (1999) 141–149.
doi:10.1023/A:1018630111976
- [2] P. Erdős, C. Ko and R. Rado, *Intersection theorems for systems of finite sets*, Q. J. Math. **12** (1961) 313–320.
doi:10.1093/qmath/12.1.313
- [3] P. Frankl and Z. Füredi, *Exact solution of some Turán-type problems*, J. Combin. Theory Ser. A **45** (1987) 226–262.
doi:10.1016/0097-3165(87)90016-1
- [4] P. Frankl and Z. Füredi, *Non-trivial intersecting families*, J. Combin. Theory Ser. A **41** (1986) 150–153.
doi:10.1016/0097-3165(86)90121-4
- [5] Z. Füredi, T. Jiang and R. Seiver, *Exact solution of the hypergraph Turán problem for k -uniform linear paths*, Combinatorica **34** (2014) 299–322.
doi:10.1007/s00493-014-2838-4

- [6] A. Gyárfás and G. Raeisi, *The Ramsey number of loose triangles and quadrangles in hypergraphs*, Electron. J. Combin. **19** (2012) # R30.
- [7] J. Han and Y. Kohayakawa, *The maximum size of a non-trivial intersecting uniform family that is not a subfamily of the Hilton-Milner family*, Proc. Amer. Math. Soc. **145** (2017) 73–87.
doi:101090/proc/13221
- [8] A.J.W. Hilton, and E.C. Milner, *Some intersection theorems for systems of finite sets*, Q. J. Math. **18** (1967) 369–384.
doi:10.1093/qmath/18.1.369
- [9] E. Jackowska, *The 3-color Ramsey number for a 3-uniform loose path of length 3*, Australas. J. Combin. **63** (2015) 314–320.
- [10] E. Jackowska, J. Polcyn and A. Ruciński, *Turán numbers for linear 3-uniform paths of length 3*, Electron. J. Combin. **23(2)** (2016) #P2.30.
- [11] E. Jackowska, J. Polcyn and A. Ruciński, *Multicolor Ramsey numbers and restricted Turán numbers for the loose 3-uniform path of length three*, submitted.
- [12] A. Kostochka, D. Mubayi and J. Verstraëte, *Turán problems and shadows I: paths and cycles*, J. Combin. Theory Ser. A **129** (2015) 57–79
doi:10.1016/j.jcta.2014.09.005
- [13] G.R. Omid and M. Shahsiah, *Ramsey numbers of 3-uniform loose paths and loose cycles*, J. Combin. Theory Ser. A **121** (2014) 64–73.
doi:10.1016/j.jcta.2013.09.003
- [14] J. Polcyn and A. Ruciński, *A hierarchy of maximal intersecting triple systems*, Opuscula Math. **37(4)** (2017), to appear.
arXiv:1608.06114
- [15] J. Polcyn and A. Ruciński, *Refined Turán and Ramsey numbers for the loose 3-uniform path of length three*, Discrete Math. **340** (2017) 107–118.
doi:10.1016/j.disc.2016.08.006

Received 28-12-2015

Revised 9-05-2016

Accepted 9-05-2016