

THE TOTAL ACQUISITION NUMBER OF THE RANDOMLY WEIGHTED PATH

ANANT GODBOLE

Department of Mathematics and Statistics
East Tennessee State University

e-mail: godbolea@mail.etsu.edu

ELIZABETH KELLEY

Department of Mathematics
University of Minnesota

e-mail: ekelley@g.hmc.edu

EMILY KURTZ

Department of Mathematics
Wellesley College

e-mail: ekurtz@wellesley.edu

PAWEŁ PRAŁAT

Department of Mathematics
Ryerson University

e-mail: pralat@ryerson.ca

AND

YIGUANG ZHANG

Department of Applied Mathematics and Statistics
The Johns Hopkins University

e-mail: yzhan132@jhu.edu

Abstract

There exists a significant body of work on determining the acquisition number $a_t(G)$ of various graphs when the vertices of those graphs are each initially assigned a unit weight. We determine properties of the acquisition

number of the path, star, complete, complete bipartite, cycle, and wheel graphs for variations on this initial weighting scheme, with the majority of our work focusing on the expected acquisition number of randomly weighted graphs. In particular, we bound the expected acquisition number $E(a_t(P_n))$ of the n -path when n distinguishable “units” of integral weight, or chips, are randomly distributed across its vertices between $0.242n$ and $0.375n$. With computer support, we improve it by showing that $E(a_t(P_n))$ lies between $0.29523n$ and $0.29576n$. We then use subadditivity to show that the limiting ratio $\lim E(a_t(P_n))/n$ exists, and simulations reveal more exactly what the limiting value equals. The Hoeffding-Azuma inequality is used to prove that the acquisition number is tightly concentrated around its expected value. Additionally, in a different context, we offer a non-optimal acquisition protocol algorithm for the randomly weighted path and exactly compute the expected size of the resultant residual set.

Keywords: total acquisition number, Poissonization, dePoissonization, Maxwell-Boltzman and Bose-Einstein allocation.

2010 Mathematics Subject Classification: 05C75.

REFERENCES

- [1] D. Bal, P. Bennett, A. Dudek and P. Prałat, *The total acquisition number of random graphs* (2016), arXiv:1402.2854.
- [2] E. Czabarka, M. Marsili and L.A. Székely, *Threshold functions for distinct parts: revisiting Erdős-Lehner*, in: Information Theory, Combinatorics, and Search Theory (in Memory of Rudolph Ahlswede), H. Aydinian, F. Cicalese and C. Deppe (Eds.), Springer-Verlag, Lecture Notes in Comput. Sci. **7777** (2013) 463–471. doi:10.1007/978-3-642-36899-8_22
- [3] D.E. Lampert and P.J. Slater, *The acquisition number of a graph*, Congr. Numer. **109** (1995) 203–210.
- [4] T. LeSaulnier, N. Prince, P. Wenger, D. West and P. Worah, *Total acquisition in graphs*, SIAM J. Discrete Math. **27** (2013) 1800–1819. doi:10.1137/110856186
- [5] P. Wenger, *Fractional acquisition in graphs*, Discrete Appl. Math. **178** (2014) 142–148. doi:10.1016/j.dam.2014.06.010
- [6] D. West, N. Prince and P. Wenger, *Unit acquisition in graphs*, preprint.
- [7] P. Prałat, C++ program and results can be found at <http://www.math.ryerson.ca/~pralat/>.

Received 14 August 2015

Revised 15 July 2016

Accepted 15 July 2016