

SHARP UPPER BOUNDS ON THE CLAR NUMBER OF FULLERENE GRAPHS¹

YANG GAO

*School of Traffic and Transportation
Lanzhou Jiaotong University
Lanzhou, Gansu 730070, P.R. China*

e-mail: gaoy@mail.lzjtu.cn

AND

HEPING ZHANG²

*School of Mathematics and Statistics
Lanzhou University
Lanzhou, Gansu 730000, P.R. China*

e-mail: zhanghp@lzu.edu.cn

Abstract

The Clar number of a fullerene graph with n vertices is bounded above by $\lfloor n/6 \rfloor - 2$ and this bound has been improved to $\lfloor n/6 \rfloor - 3$ when n is congruent to 2 modulo 6. We can construct at least one fullerene graph attaining the upper bounds for every even number of vertices $n \geq 20$ except $n = 22$ and $n = 30$.

Keywords: fullerene, Clar number, Clar set, leapfrog transformation.

2010 Mathematics Subject Classification: 05C10, 05C62, 05C70, 92E10.

REFERENCES

- [1] P.W. Fowler and D.E. Manolopoulos, *An Atlas of Fullerenes* (Oxford University Press, Oxford, 1995).
- [2] Y. Gao and H. Zhang, *The Clar number of fullerenes on surfaces*, MATCH Commun. Math. Comput. Chem. **72** (2014) 411–426.

¹This work is supported by NSFC (grant no. 11371180), and the Fundamental Research Funds for the Central Universities (grant nos. lzujbky-2017-ct01, lzujbky-2016-ct12).

²Corresponding author.

- [3] Y. Gao and H. Zhang, *Clar structure and Fries set of fullerenes and (4, 6)-fullerenes on surfaces*, J. Appl. Math. (2014) Article ID 196792, 11 pages.
- [4] Y. Gao, Q. Li and H. Zhang, *Fullerenes with the maximum Clar number*, Discrete Appl. Math. **202** (2016) 58–69.
doi:10.1016/j.dam.2015.08.007
- [5] C. Godsil and G. Royle, Algebraic Graph Theory (Springer-Verlag, New York, 2001).
doi:10.1007/978-1-4613-0163-9
- [6] B. Grünbaum and T. Motzkin, *The number of hexagons and the simplicity of geodesics on certain polyhedra*, Canad. J. Math. **15** (1963) 744–751.
doi:10.4153/CJM-1963-071-3
- [7] E. Hartung, *Fullerenes with complete Clar structure*, Discrete Appl. Math. **161** (2013) 2952–2957.
doi:10.1016/j.dam.2013.06.009
- [8] T. Pisanski and M. Randić, *Bridges between geometry and graph theory*, in: Geometry at Work: Papers in Applied Geometry Vol. 53, C.A. Gorini (Ed(s)), (Washington, DC, Mathematical Association of America, 2000) 174–194.
- [9] D. Ye and H. Zhang, *Extremal fullerene graphs with the maximum Clar number*, Discrete Appl. Math. **157** (2009) 3152–3173.
doi:10.1016/j.dam.2009.06.007
- [10] H. Zhang and F. Zhang, *The Clar covering polynomial of hexagonal systems I*, Discrete Appl. Math. **69** (1996) 147–167.
doi:10.1016/0166-218X(95)00081-2
- [11] H. Zhang and D. Ye, *An upper bound for Clar number of fullerene graphs*, J. Math. Chem. **42** (2007) 123–133.
doi:10.1007/s10910-006-9061-5
- [12] H. Zhang, D. Ye and Y. Liu, *A combination of Clar number and Kekulé count as an indicator of relative stability of fullerene isomers of C_{60}* , J. Math. Chem. **48** (2010) 733–740.
doi:10.1007/s10910-010-9706-2

Received 12 February 2016

Revised 26 October 2016

Accepted 26 October 2016