

ON INDEPENDENT $[1, 2]$ -SETS IN TREES ¹

SAHAR A. ALEID, JOSÉ CÁCERES

AND

MARÍA LUZ PUERTAS

Department of Mathematics
University of Almeria, Spain

e-mail: sahar.aleid@ual.es
jcaceres@ual.es
mpuertas@ual.es

Abstract

An $[1, k]$ -set S in a graph G is a dominating set such that every vertex not in S has at most k neighbors in it. If the additional requirement that the set must be independent is added, the existence of such sets is not guaranteed in every graph. In this paper we solve some problems previously posed by other authors about independent $[1, 2]$ -sets. We provide a necessary condition for a graph to have an independent $[1, 2]$ -set, in terms of spanning trees, and we prove that this condition is also sufficient for cactus graphs. We follow the concept of excellent tree and characterize the family of trees such that any vertex belongs to some independent $[1, 2]$ -set. Finally, we describe a linear algorithm to decide whether a tree has an independent $[1, 2]$ -set. This algorithm can be easily modified to obtain the cardinality of a smallest independent $[1, 2]$ -set of a tree.

Keywords: domination, independence, spanning trees, excellent trees.

2010 Mathematics Subject Classification: 05C69.

REFERENCES

- [1] T.A. Beyer, A. Proskurowski, S.T. Hedetniemi and S. Mitchell, *Independent domination in trees*, Congr. Numer. **19** (1997) 321–328.
- [2] G. Chartrand, L. Lesniak and P. Zhang, *Graphs and Digraphs*, 5th edition (CRC Press, 2011).

¹Partially supported by Phoenix Project (Erasmus Mundus Programme), MTM2014-60127-P (Ministerio de Economía y Competitividad) and FQM305 (Junta de Andalucía).

- [3] M. Chellali, S.T. Hedetniemi and A. McRae, *[1, 2]-set in graphs*, Discrete Appl. Math. **161** (2013) 2885–2893.
doi:10.1016/j.dam.2013.06.012
- [4] M. Chellali, O. Favaron, T.W. Haynes, S.T. Hedetniemi and A. McRae, *Independent $[1, k]$ -sets in graphs*, Australas. J. Combin. **59** (2014) 144–156.
- [5] E. Cockayne, S. Goodman and S. Hedetniemi, *A linear algorithm for the domination number of a tree*, Inform. Process. Lett. **4** (1975) 41–44.
doi:10.1016/0020-0190(75)90011-3
- [6] G.F. Fricke, T.W. Haynes, S.M. Hedetniemi, S.T. Hedetniemi and R.C. Laskar, *Excellent trees*, Bull. Inst. Combin. Appl. **34** (2002) 27–38.
- [7] J.H. Hattingh and M.A. Henning, *Restrained domination excellent trees*, Ars Combin. **87** (2008) 337–351.
- [8] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, *Fundamentals of Domination in Graphs* (Marcel Dekker Inc, 1998).
- [9] T.W. Haynes and M.A. Henning, *A characterization of i -excellent trees*, Discrete Math. **248** (2002) 69–77.
doi:10.1016/S0012-365X(01)00274-6
- [10] M.A. Henning, *Total domination excellent trees*, Discrete Math. **263** (2003) 93–104.
doi:10.1016/S0012-365X(02)00572-1
- [11] R. Laskar, J. Pfaff, S.M. Hedetniemi and S.T. Hedetniemi, *On the algorithmic complexity of total domination*, SIAM J. Algebraic Discrete Methods **5** (1984) 420–425.
doi:10.1137/0605040
- [12] K.S. Natarajan and L.J. White, *Optimum domination in weighted trees*, Inform. Process. Lett. **7** (1978) 261–265.
doi:10.1016/0020-0190(78)90012-1
- [13] S.L. Mitchell and S.T. Hedetniemi, *Edge domination in trees*, Congr. Numer. **19** (1977) 489–509.
- [14] P.J. Slater, *R -domination in graphs*, J. ACM **23** (1976) 446–450.
doi:10.1145/321958.321964
- [15] M. Yannakakis and F. Gavril, *Edge dominating sets in graphs*, SIAM J. Appl. Math. **38** (1980) 364–372.
doi:10.1137/0138030

Received 5 October 2016
 Revised 23 December 2016
 Accepted 23 January 2017