

ORIENTED INCIDENCE COLOURINGS OF DIGRAPHS

CHRISTOPHER DUFFY

Department of Mathematics and Statistics
University of Saskatchewan
Saskatoon, Canada

e-mail: christopher.duffy@usask.ca

GARY MACGILLIVRAY

Department of Mathematics and Statistics
University of Victoria
Victoria, Canada

e-mail: gmacgill@math.uvic.ca

PASCAL OCHEM

LIRMM, CNRS,
Université de Montpellier,
Montpellier, France

e-mail: pascal.ochem@lirmm.fr

AND

ANDRÉ RASPAUD

LaBRI UMR CNRS 5800
Université Bordeaux 1,
Bordeaux, France

e-mail: raspaud@labri.fr

Abstract

Brualdi and Quinn Massey [6] defined incidence colouring while studying the strong edge chromatic index of bipartite graphs. Here we introduce a similar concept for digraphs and define the oriented incidence chromatic number. Using digraph homomorphisms, we show that the oriented incidence chromatic number of a digraph is closely related to the chromatic number of the underlying simple graph. This motivates our study of the oriented incidence chromatic number of symmetric complete digraphs. We give

upper and lower bounds for the oriented incidence chromatic number of these graphs, as well as digraphs arising from common graph constructions and decompositions. Additionally we construct, for all $k \geq 2$, a target digraph H_k for which oriented incidence k colouring is equivalent to homomorphism to H_k .

Keywords: digraph homomorphism, graph colouring, incidence colouring, computational complexity.

2010 Mathematics Subject Classification: 05C15.

REFERENCES

- [1] K. Appel, W. Haken and J. Koch, *Every planar map is four colorable. Part rm II: Reducibility*, Illinois J. Math. **21** (1977) 491–567.
- [2] J. Bang-Jensen, P. Hell and G. MacGillivray, *The complexity of colouring by semi-complete digraphs*, SIAM J. Discrete Math. **1** (1988) 281–298.
doi:10.1137/0401029
- [3] L. Barto, M. Kozik and T. Niven, *The CSP dichotomy holds for digraphs with no sources and no sinks (a positive answer to a conjecture of Bang-Jensen and Hell)*, SIAM J. Comput. **38** (2009) 1782–1802.
doi:10.1137/070708093
- [4] J. Bensmail, C. Duffy and S. Sen, *Analogues of cliques for (m, n) -colored mixed graphs*, Graphs Combin. **33** (2017) 735 – 750.
doi:10.1007/s00373-017-1807-2
- [5] J.A. Bondy and U.S.R. Murty, *Graph Theory* (Springer-Verlag, London, 2008).
- [6] R.A. Brualdi and J.J. Quinn Massey, *Incidence and strong edge colorings of graphs*, Discrete Math. **122** (1993) 51–58.
doi:10.1016/0012-365X(93)90286-3
- [7] B. Courcelle, *The monadic second order logic of graphs VI: on several representations of graphs by relational structures.*, Discrete Appl. Math. **54** (1994) 117–129.
doi:10.1016/0166-218X(94)90019-1
- [8] W.D. Fellner, *On minimal graphs*, Theoret. Comput. Sci. **17** (1982) 103–110.
doi:10.1016/0304-3975(82)90135-9
- [9] R.L. Graham and N.J.A. Sloane, *Lower bounds for constant weight codes*, IEEE Trans. Inform. Theory **26** (1980) 37–43.
doi:10.1109/TIT.1980.1056141
- [10] P. Hell and J. Nešetřil, *On the complexity of H -coloring*, J. Combin. Theory Ser. B **48** (1990) 92–110.
doi:10.1016/0095-8956(90)90132-J
- [11] M. Hosseini Dolama, E. Sopena and X. Zhu, *Incidence coloring of k -degenerated graphs*, Discrete Math. **283** (2004) 121–128.
doi:10.1016/j.disc.2004.01.015

- [12] W. Klostermeyer and G. MacGillivray, *Homomorphisms and oriented colorings of equivalence classes of oriented graphs*, Discrete Math. **274** (2004) 161–172.
doi:10.1016/S0012-365X(03)00086-4
- [13] A.V. Kostochka, E. Sopena and X. Zhu, *Acyclic and oriented chromatic numbers of graphs*, J. Graph Theory **24** (1997) 331–340.
doi:10.1002/(SICI)1097-0118(199704)24:4<331::AID-JGT5>3.0.CO;2-P
- [14] G. MacGillivray and K. Sherk, *A theory of 2-dipath colourings*, Australas. J. Combin. **60** (2014) 11–26.
- [15] G. MacGillivray and J.S. Swarts, *Obstructions to homomorphisms involving the graft extension*, manuscript.
- [16] T.H. Marshall, *Homomorphism bounds for oriented planar graphs*, J. Graph Theory **55** (2007) 175–190.
doi:10.1002/jgt.20233
- [17] J. Nešetřil and A. Pultr, *On classes of relations and graphs determined by subobjects and factorobjects*, Discrete Math. **22** (1978) 287–300.
doi:10.1016/0012-365X(78)90062-6
- [18] S. Sen, *Maximum order of a planar oclique is 15*, in: Proc. 23rd International Workshop on Combinatorial Algorithms, W.F. Smyth and S. Arumugam (Ed(s)), (Springer, 2012) 130–142.
doi:10.1007/978-3-642-35926-2_16
- [19] E. Sopena, *Oriented graph coloring*, Discrete Math. **229** (2001) 359–369.
doi:10.1016/S0012-365X(00)00216-8
- [20] P.K. Sun, *Incidence coloring of regular graphs and complement graphs*, Taiwanese J. Math. **16** (2012) 2289–2295.
doi:10.11650/twjm/1500406852
- [21] E. Welzl, *Color-families are dense*, Theoret. Comput. Sci. **17** (1982) 29–41.
doi:10.1016/0304-3975(82)90129-3
- [22] J. Wu, *Some results on the incidence coloring number of a graph*, Discrete Math. **309** (2009) 3866–3870.
doi:10.1016/j.disc.2008.10.027
- [23] K. Young, *2-Dipath and Proper 2-Dipath Colouring* (Masters Thesis, University of Victoria, 2009).

Received 12 September 2016
 Revised 12 July 2017
 Accepted 13 July 2017

Appendix

Colour Class	Vertex List
1	(1, (7, 1)), (3, (3, 2)), (2, (4, 2)), (3, (4, 3)), (3, (3, 5)), (5, (4, 5)), (5, (6, 5)), (3, (3, 7)), (2, (7, 2)), (5, (7, 5)), (6, (7, 6)), (1, (6, 1)), (6, (6, 4)), (6, (6, 2))
2	(6, (1, 6)), (7, (1, 7)), (2, (3, 2)), (4, (3, 4)), (4, (4, 1)), (4, (4, 2)), (2, (5, 2)), (4, (4, 5)), (6, (5, 6)), (4, (4, 7)), (7, (3, 7)), (7, (5, 7)), (6, (6, 7)), (6, (6, 3))
3	(1, (1, 2)), (1, (1, 3)), (1, (1, 4)), (1, (1, 5)), (1, (1, 6)), (1, (1, 7)), (3, (2, 3)), (4, (2, 4)), (5, (2, 5)), (6, (2, 6)), (7, (2, 7))
4	(1, (2, 1)), (1, (3, 1)), (1, (4, 1)), (5, (3, 5)), (5, (5, 4)), (5, (5, 1)), (5, (5, 2)), (7, (4, 7)), (7, (7, 2)), (7, (7, 3)), (6, (3, 6)), (5, (5, 6)), (6, (4, 6)), (7, (7, 6)), (7, (7, 1))
5	(2, (2, 1)), (1, (5, 1)), (1, (6, 1)), (1, (3, 1)), (2, (2, 4)), (2, (2, 5)), (3, (3, 4)), (3, (5, 3)), (4, (5, 4)), (3, (3, 6)), (2, (6, 2)), (7, (7, 4)), (7, (7, 5)), (4, (6, 4)), (7, (6, 7))
6	(2, (1, 2)), (3, (1, 3)), (4, (1, 4)), (5, (1, 5)), (2, (2, 3)), (4, (4, 3)), (2, (2, 6)), (2, (2, 7)), (5, (5, 3)), (3, (6, 3)), (3, (7, 3)), (4, (7, 4)), (5, (5, 7)), (4, (4, 6)), (5, (6, 5))

Table 1. An oriented incidence colouring \vec{K}_7 with six colours.

Colour Class	Vertex List
1	(6, (6, 8)), (1, (2, 1)), (3, (3, 1)), (3, (3, 2)), (2, (2, 4)), (3, (3, 4)), (4, (5, 4)), (6, (2, 6)), (6, (6, 1)), (6, (6, 4)), (6, (6, 5)), (4, (7, 4)), (8, (5, 8)), (3, (5, 3)), (1, (7, 1)), (6, (7, 6)), (3, (7, 3)), (1, (5, 1))
2	(8, (8, 5)), (8, (8, 6)), (2, (2, 1)), (2, (2, 3)), (8, (3, 8)), (1, (3, 1)), (4, (4, 1)), (4, (3, 4)), (4, (4, 5)), (4, (4, 7)), (6, (3, 6)), (2, (2, 6)), (4, (4, 6)), (6, (5, 6)), (2, (5, 2)), (7, (5, 7)), (2, (2, 7))
3	(1, (8, 1)), (2, (8, 2)), (3, (8, 3)), (4, (8, 4)), (5, (8, 5)), (6, (8, 6)), (7, (8, 7)), (1, (1, 2)), (1, (1, 3)), (1, (1, 4)), (1, (1, 5)), (1, (1, 6)), (7, (7, 2)), (7, (7, 4)), (7, (7, 5)), (7, (7, 6)), (7, (7, 3))
4	(1, (1, 8)), (1, (1, 7)), (2, (2, 8)), (3, (3, 8)), (2, (2, 4)), (3, (4, 3)), (8, (4, 8)), (1, (4, 1)), (3, (3, 6)), (1, (6, 1)), (2, (6, 2)), (5, (7, 5)), (2, (2, 5)), (3, (3, 5)), (5, (4, 5)), (5, (6, 5)), (2, (7, 2)), (3, (3, 7)), (8, (6, 8)), (8, (7, 8))
5	(8, (1, 8)), (4, (4, 8)), (5, (5, 8)), (7, (7, 8)), (8, (2, 8)), (3, (1, 3)), (4, (1, 4)), (3, (2, 3)), (4, (4, 2)), (4, (4, 3)), (5, (2, 5)), (5, (5, 1)), (3, (6, 3)), (4, (6, 4)), (5, (5, 3)), (7, (2, 7)), (7, (7, 1)), (5, (5, 6)), (7, (6, 7))
6	(8, (8, 1)), (8, (8, 2)), (8, (8, 3)), (8, (8, 4)), (8, (8, 7)), (2, (1, 2)), (2, (3, 2)), (5, (1, 5)), (6, (1, 6)), (7, (1, 7)), (2, (4, 2)), (5, (5, 2)), (6, (6, 2)), (6, (6, 3)), (6, (4, 6)), (5, (3, 5)), (7, (4, 7)), (5, (5, 4)), (5, (5, 7)), (6, (6, 7)), (7, (3, 7))

Table 2. An oriented incidence colouring \vec{K}_8 with six colours.




Figure 1. Oriented incidence colourings of $\vec{K}_4, \vec{K}_5, \vec{K}_6$ with the minimum number of colours. The colouring of \vec{K}_4 is obtained by deleting any vertex in the colouring of \vec{K}_5 .