

1-RESTRICTED OPTIMAL RUBBLING ON GRAPHS

ROBERT A. BEELER

*Department of Mathematics and Statistics
East Tennessee State University
Johnson City, TN 37614-1700 USA*

e-mail: beelerr@etsu.edu

TERESA W. HAYNES

*Department of Mathematics and Statistics
East Tennessee State University
Johnson City, TN 37614-1700 USA*

*Department of Mathematics
University of Johannesburg
Auckland Park, South Africa*

e-mail: haynes@etsu.edu

AND

KYLE MURPHY

*Department of Mathematics
Iowa State University
Ames, IA 50011 USA*

e-mail: kylem2@iastate.edu

Abstract

Let G be a graph with vertex set V and a distribution of pebbles on the vertices of V . A pebbling move consists of removing two pebbles from a vertex and placing one pebble on a neighboring vertex, and a rubbing move consists of removing a pebble from each of two neighbors of a vertex v and placing a pebble on v . We seek an initial placement of a minimum total number of pebbles on the vertices in V , so that no vertex receives more than one pebble and for any given vertex $v \in V$, it is possible, by a sequence of pebbling and rubbing moves, to move at least one pebble to v . This minimum number of pebbles is the 1-restricted optimal rubbing number. We determine the 1-restricted optimal rubbing numbers for Cartesian products. We also present bounds on the 1-restricted optimal rubbing number.

Keywords: graph pebbling, graph rubbing, optimal rubbing, t -restricted optimal pebbling.

2010 Mathematics Subject Classification: 05C78.

REFERENCES

- [1] R.A. Beeler, T.W. Haynes and K. Murphy, *An introduction to t -restricted optimal rubbing*, Congr. Numer., to appear.
- [2] C. Belford and N. Sieben, *Rubbling and optimal rubbing of graphs*, Discrete Math. **309** (2009) 3436–3446.
doi:10.1016/j.disc.2008.09.035
- [3] T.A. Clarke, R.A. Hochberg and G.H. Hurlbert, *Pebbling in diameter two graphs and products of paths*, J. Graph Theory **25** (1997) 119–128.
doi:10.1002/(SICI)1097-0118(199706)25:2<119::AID-JGT3>3.0.CO;2-P
- [4] R. Feng and J.Y. Kim, *Graham’s pebbling conjecture on product of complete bipartite graphs*, Sci. China Ser. A–Math. **44** (2001) 817–822.
doi:10.1007/BF02880130
- [5] H.-L. Fu and C.-L. Shiue, *The optimal pebbling number of the complete m -ary tree*, Discrete Math. **222** (2000) 89–100.
doi:10.1016/S0012-365X(00)00008-X
- [6] D.S. Herscovici, *Graham’s pebbling conjecture on products of cycles*, J. Graph Theory **42** (2003) 141–154.
doi:10.1002/jgt.10080
- [7] G.Y. Katona and L.F. Papp, *The optimal rubbing number of ladders, prisms and Möbius-ladders*, Discrete Appl. Math. **209** (2016) 227–246.
doi:10.1016/j.dam.2015.10.026
- [8] G.Y. Katona and N. Sieben, *Bounds on the rubbing and optimal rubbing numbers of graphs*, Graphs Combin. **29** (2013) 535–551.
doi:10.1007/s00373-012-1146-2
- [9] S.S. Wang, *Pebbling and Graham’s conjecture*, Discrete Math. **226** (2001) 431–438.
doi:10.1016/S0012-365X(00)00177-1

Received 1 May 2017
Revised 23 October 2017
Accepted 24 October 2017