

ON THE METRIC DIMENSION OF DIRECTED AND UNDIRECTED CIRCULANT GRAPHS

TOMÁŠ VETRÍK

Department of Mathematics and Applied Mathematics
University of the Free State
Bloemfontein, South Africa

e-mail: vetrik@ufs.ac.za

Abstract

The undirected circulant graph $C_n(\pm 1, \pm 2, \dots, \pm t)$ consists of vertices v_0, v_1, \dots, v_{n-1} and undirected edges $v_i v_{i+j}$, where $0 \leq i \leq n-1$, $1 \leq j \leq t$ ($2 \leq t \leq \frac{n}{2}$), and the directed circulant graph $C_n(1, t)$ consists of vertices v_0, v_1, \dots, v_{n-1} and directed edges $v_i v_{i+1}, v_i v_{i+t}$, where $0 \leq i \leq n-1$ ($2 \leq t \leq n-1$), the indices are taken modulo n . Results on the metric dimension of undirected circulant graphs $C_n(\pm 1, \pm t)$ are available only for special values of t . We give a complete solution of this problem for directed graphs $C_n(1, t)$ for every $t \geq 2$ if $n \geq 2t^2$. Grigorious *et al.* [*On the metric dimension of circulant and Harary graphs*, Appl. Math. Comput. 248 (2014) 47–54] presented a conjecture saying that $\dim(C_n(\pm 1, \pm 2, \dots, \pm t)) = t + p - 1$ for $n = 2tk + t + p$, where $3 \leq p \leq t + 1$. We disprove it by showing that $\dim(C_n(\pm 1, \pm 2, \dots, \pm t)) \leq t + \frac{p+1}{2}$ for $n = 2tk + t + p$, where $t \geq 4$ is even, p is odd, $1 \leq p \leq t + 1$ and $k \geq 1$.

Keywords: metric dimension, resolving set, circulant graph, distance.

2010 Mathematics Subject Classification: 05C35, 05C12.

REFERENCES

- [1] A. Borchert and S. Gosselin, *The metric dimension of circulant graphs and Cayley hypergraphs*, Util. Math. **106** (2018) 125–147.
- [2] G. Chartrand, L. Eroh, M.A. Johnson and O.R. Oellermann, *Resolvability in graphs and the metric dimension of a graph*, Discrete Appl. Math. **105** (2000) 99–113.
doi:10.1016/S0166-218X(00)00198-0
- [3] K. Chau and S. Gosselin, *The metric dimension of circulant graphs and their Cartesian products*, Opuscula Math. **37** (2017) 509–534.
doi:10.7494/OpMath.2017.37.4.509

- [4] C. Grigorious, P. Manuel, M. Miller, B. Rajan and S. Stephen, *On the metric dimension of circulant and Harary graphs*, Appl. Math. Comput. **248** (2014) 47–54.
doi:10.1016/j.amc.2014.09.045
- [5] M. Imran, *On the metric dimension of barycentric subdivision of Cayley graphs*, Acta Math. Appl. Sin. Eng. Ser. **32** (2016) 1067–1072.
doi:10.1007/s10255-016-0627-0
- [6] M. Imran, A.Q. Baig, S.A. Ul. Hag Bokhary and I. Javaid, *On the metric dimension of circulant graphs*, Appl. Math. Lett. **25** (2012) 320–325.
doi:10.1016/j.aml.2011.09.008
- [7] I. Javaid, M.N. Azhar and M. Salman, *Metric dimension and determining number of Cayley graphs*, World Appl. Sci. J. **18** (2012) 1800–1812.
- [8] I. Javaid, M.T. Rahim and K. Ali, *Families of regular graphs with constant metric dimension*, Util. Math. **75** (2008) 21–33.
- [9] S. Khuller, B. Raghavachari and A. Rosenfeld, *Landmarks in graphs*, Discrete Appl. Math. **70** (1996) 217–229.
doi:10.1016/0166-218X(95)00106-2
- [10] R.A. Melter and I. Tomescu, *Metric bases in digital geometry*, Comput. Vision Graphics Image Process. **25** (1984) 113–121.
doi:10.1016/0734-189X(84)90051-3
- [11] M. Salman, I. Javaid and M.A. Chaudhry, *Resolvability in circulant graphs*, Acta Math. Sin. Eng. Ser. **28** (2012) 1851–1864.
doi:10.1007/s10114-012-0417-4
- [12] S.W. Saputro, R. Simanjuntak, S. Uttunggadewa, H. Assiyatun, E.T. Baskoro, A.N.M. Salman and M. Baća, *The metric dimension of the lexicographic product of graphs*, Discrete Math. **313** (2013) 1045–1051.
doi:10.1016/j.disc.2013.01.021
- [13] T. Vetrík, *The metric dimension of circulant graphs*, Canad. Math. Bull. **60** (2017) 206–216.
doi:10.4153/CMB-2016-048-1

Received 25 May 2017
 Revised 23 January 2018
 Accepted 23 January 2018