

## INCIDENCE COLORING—COLD CASES

FRANTIŠEK KARDOŠ<sup>1</sup>, MÁRIA MACEKOVÁ<sup>2</sup>

MARTINA MOCKOVČIAKOVÁ<sup>3</sup>, ÉRIC SOPENA<sup>1</sup>

AND

ROMAN SOTÁK<sup>2</sup>

<sup>1</sup>*Univ. Bordeaux, Bordeaux INP, CNRS, LaBRI,  
UMR5800, F-33400 Talence, France*

<sup>2</sup>*Institute of Mathematics  
P.J. Šafárik University in Košice  
Jesenná 5, 04001 Košice, Slovakia*

<sup>3</sup>*European Centre of Excellence NTIS  
University of West Bohemia  
Pilsen, Czech Republic*

**e-mail:** frantisek.kardos@labri.fr  
maria.macekova@upjs.sk  
mmockov@ntis.zcu.cz  
eric.sopena@labri.fr  
roman.sotak@upjs.sk

### Abstract

An *incidence* in a graph  $G$  is a pair  $(v, e)$  where  $v$  is a vertex of  $G$  and  $e$  is an edge of  $G$  incident to  $v$ . Two incidences  $(v, e)$  and  $(u, f)$  are adjacent if at least one of the following holds: (i)  $v = u$ , (ii)  $e = f$ , or (iii) edge  $vu$  is from the set  $\{e, f\}$ . An *incidence coloring* of  $G$  is a coloring of its incidences assigning distinct colors to adjacent incidences. The minimum number of colors needed for incidence coloring of a graph is called the *incidence chromatic number*.

It was proved that at most  $\Delta(G) + 5$  colors are enough for an incidence coloring of any planar graph  $G$  except for  $\Delta(G) = 6$ , in which case at most 12 colors are needed. It is also known that every planar graph  $G$  with girth at least 6 and  $\Delta(G) \geq 5$  has incidence chromatic number at most  $\Delta(G) + 2$ .

In this paper we present some results on graphs regarding their maximum degree and maximum average degree. We improve the bound for planar graphs with  $\Delta(G) = 6$ . We show that the incidence chromatic number is at

most  $\Delta(G) + 2$  for any graph  $G$  with  $\text{mad}(G) < 3$  and  $\Delta(G) = 4$ , and for any graph with  $\text{mad}(G) < \frac{10}{3}$  and  $\Delta(G) \geq 8$ .

**Keywords:** incidence coloring, incidence chromatic number, planar graph, maximum average degree.

**2010 Mathematics Subject Classification:** 05C15.

# REFERENCES

- [1] M. Bonamy, B. L  v  que and A. Pinlou, *2-distance coloring of sparse graphs*, J. Graph Theory **77** (2014) 190–218.  
doi:10.1002/jgt.21782
- [2] R.A. Brualdi and J.J. Quinn Massey, *Incidence and strong edge colorings of graphs*, Discrete Math. **122** (1993) 51–58.  
doi:10.1016/0012-365X(93)90286-3
- [3] B. Guiduli, *On incidence coloring and star arboricity of graphs*, Discrete Math. **163** (1997) 275–278.  
doi:10.1016/0012-365X(95)00342-T
- [4] S.L. Hakimi, J. Mitchem and E. Schmeichel, *Star arboricity of graphs*, Discrete Math. **149** (1996) 93–98.  
doi:10.1016/0012-365X(94)00313-8
- [5] M. Hosseini Dolama,   . Sopena and X. Zhu, *Incidence coloring of  $k$ -degenerated graphs*, Discrete Math. **283** (2004) 121–128.  
doi:10.1016/j.disc.2004.01.015
- [6] M. Hosseini Dolama and   . Sopena, *On the maximum average degree and the incidence chromatic number of a graph*, Discrete Math. Theor. Comput. Sci. **7** (2005) 203–216.
- [7] D.P. Sanders and Y. Zhao, *Planar graphs of maximum degree seven are class I*, J. Combin. Theory Ser. B **83** (2001) 201–212.  
doi:10.1006/jctb.2001.2047
- [8] D. Yang, *Fractional incidence coloring and star arboricity of graphs*, Ars Combin. **105** (2012) 213–224.

Received 9 October 2017

Revised 25 April 2018

Accepted 26 April 2018