

## ROMAN $\{2\}$ -BONDAGE NUMBER OF A GRAPH

AHMAD MORADI, DOOST ALI MOJDEH

AND

OMID SHARIFI

*Department of Mathematical Sciences*  
*University of Mazandaran, Babolsar, Iran*

**e-mail:** a.moradi@umz.ac.ir  
damojdeh@umz.ac.ir  
omid.sharifi1365@gmail.com

### Abstract

For a given graph  $G = (V, E)$ , a Roman  $\{2\}$ -dominating function  $f : V(G) \rightarrow \{0, 1, 2\}$  has the property that for every vertex  $u$  with  $f(u) = 0$ , either  $u$  is adjacent to a vertex assigned 2 under  $f$ , or is adjacent to at least two vertices assigned 1 under  $f$ . The Roman  $\{2\}$ -domination number of  $G$ ,  $\gamma_{\{R2\}}(G)$ , is the minimum of  $\sum_{u \in V(G)} f(u)$  over all such functions. In this paper, we initiate the study of the problem of finding Roman  $\{2\}$ -bondage number of  $G$ . The Roman  $\{2\}$ -bondage number of  $G$ ,  $b_{\{R2\}}$ , is defined as the cardinality of a smallest edge set  $E' \subseteq E$  for which  $\gamma_{\{R2\}}(G - E') > \gamma_{\{R2\}}(G)$ . We first demonstrate complexity status of the problem by proving that the problem is NP-Hard. Then, we derive useful parametric as well as fixed upper bounds on the Roman  $\{2\}$ -bondage number of  $G$ . Specifically, it is known that the Roman bondage number of every planar graph does not exceed 15 (see [S. Akbari, M. Khatirinejad and S. Qajar, *A note on the Roman bondage number of planar graphs*, *Graphs Combin.* 29 (2013) 327–331]). We show that same bound will be preserved while computing the Roman  $\{2\}$ -bondage number of such graphs. The paper is then concluded by computing exact value of the parameter for some classes of graphs.

**Keywords:** domination, Roman  $\{2\}$ -domination, Roman  $\{2\}$ -bondage number.

**2010 Mathematics Subject Classification:** 05C69.

- [1] S. Akbari, M. Khatirinejad and S. Qajar, *A note on the Roman bondage number of planar graphs*, *Graphs Combin.* **29** (2013) 327–331.  
doi:10.1007/s00373-011-1129-8
- [2] A. Bahremandpour, F.-T. Hu, S.M. Sheikholeslami and J.-M. Xu, *On the Roman bondage number of a graph*, *Discrete Math. Algorithms Appl.* **5** (2013) #1350001.  
doi:10.1142/S1793830913500018
- [3] E.W. Chambers, B. Kinnersley, N. Prince and D.B. West, *Extremal problems for Roman domination*, *SIAM J. Discrete Math.* **23** (2009) 1575–1586.  
doi:10.1137/070699688
- [4] M. Chellali, T.W. Haynes, S.T. Hedetniemi and A.A. McRae, *Roman  $\{2\}$ -domination*, *Discrete Appl. Math.* **204** (2016) 22–28.  
doi:10.1016/j.dam.2015.11.013
- [5] E.J. Cockayne, P.M. Dreyer Jr., S.M. Hedetniemi and S.T. Hedetniemi, *Roman domination in graphs*, *Discrete Math.* **278** (2004) 11–22.  
doi:10.1016/j.disc.2003.06.004
- [6] J.F. Fink, M.S. Jacobson, L.F. Kinch and J. Roberts, *The bondage number of a graph*, *Discrete Math.* **86** (1990) 47–57.  
doi:10.1016/0012-365X(90)90348-L
- [7] M.R. Garey and D.S. Johnson, *Computers and Intractability: A Guide to the Theory of NP-Completeness* (W.H. Freeman, 1979).
- [8] J. Kleinberg and E. Tardos, *Algorithm Design* (Pearson Education, India, 2006).
- [9] P. Roushini Leely Pushpam and T.N.M. Malini Mai, *Weak roman domination in graphs*, *Discuss. Math. Graph Theory* **31** (2011) 115–128.  
doi:10.7151/dmgt.1532
- [10] N. Jafari Rad and L. Volkmann, *Roman bondage in graphs*, *Discuss. Math. Graph Theory* **31** (2011) 763–773.  
doi:10.7151/dmgt.1578
- [11] M. Krzywkowski, *2-bondage in graphs*, *Int. J. Comput. Math.* **90** (2013) 1358–1365.  
doi:10.1080/00207160.2012.752817
- [12] T. Turaci, *On the average lower bondage number of a graph*, *RAIRO Oper. Res.* **50** (2016) 1003–1012.  
doi:10.1051/ro/2015062

Received 21 July 2017  
Revised 19 February 2018  
Accepted 12 March 2018