

ADDITIVE LIST COLORING OF PLANAR GRAPHS WITH GIVEN GIRTH

AXEL BRANDT¹

Department of Mathematics & Statistics
Northern Kentucky University, Highland Heights, KY, USA

e-mail: brandta2@nku.edu

SOGOL JAHANBEKAM²

Department of Mathematics San Jose State University
San Jose, CA, USA

e-mail: sogol.jahanbekam@sjsu.edu

AND

JENNIFER WHITE

Department of Mathematics
Saint Vincent College, Latrobe, PA, USA

e-mail: jennifer.white@stvincent.edu

Abstract

An *additive coloring* of a graph G is a labeling of the vertices of G from $\{1, 2, \dots, k\}$ such that two adjacent vertices have distinct sums of labels on their neighbors. The least integer k for which a graph G has an additive coloring is called the *additive coloring number* of G , denoted $\chi_{\Sigma}(G)$. Additive coloring is also studied under the names lucky labeling and open distinguishing. In this paper, we improve the current bounds on the additive coloring number for particular classes of graphs by proving results for a list version of additive coloring. We apply the discharging method and the Combinatorial Nullstellensatz to show that every planar graph G with girth at least 5 has $\chi_{\Sigma}(G) \leq 19$, and for girth at least 6, 7, and 26, $\chi_{\Sigma}(G)$ is at most 9, 8, and 3, respectively. In 2009, Czerwiński, Grytczuk, and Żelazny conjectured that $\chi_{\Sigma}(G) \leq \chi(G)$, where $\chi(G)$ is the chromatic number of G . Our result for

¹Partially funded by NSF GK-12 Transforming Experiences Grant DGE-0742434

²Research supported in part by NSF grant CMMI-1727743.

the class of non-bipartite planar graphs of girth at least 26 is best possible and affirms the conjecture for this class of graphs.

Keywords: lucky labeling, additive coloring, reducible configuration, discharging method, Combinatorial Nullstellensatz.

2010 Mathematics Subject Classification: 05C78, 05C15, 05C22, 05C10.

REFERENCES

- [1] A. Ahadi, A. Dehghan, M. Kazemi and E. Mollaahmadi, *Computation of lucky number of planar graphs is NP-hard*, Inform. Process. Lett. **112** (2012) 109–112.
doi:10.1016/j.ipl.2011.11.002
- [2] A. Ahadi and A. Dehghan, *The inapproximability for the $(0,1)$ -additive number*, Discrete Math. Theor. Comput. Sci. **17** (2016) 217–226.
- [3] S. Akbari, M. Ghanbari, R. Manaviyat and S. Zare, *On the lucky choice number of graphs*, Graphs Combin. **29** (2013) 157–163.
doi:10.1007/s00373-011-1112-4
- [4] N. Alon, *Combinatorial Nullstellensatz*, Combin. Probab. Comput. **8** (1999) 7–29.
doi:10.1017/S0963548398003411
- [5] M. Axenovich, J. Harant, J. Przybyło, R. Soták, M. Voigt and J. Weidelich, *A note on adjacent vertex distinguishing colorings of graphs*, Discrete Appl. Math. **205** (2016) 1–7.
doi:10.1016/j.dam.2015.12.005
- [6] T. Bartnicki, B. Bosek, S. Czerwiński, J. Grytczuk, G. Matecki and W. Żelazny, *Additive coloring of planar graphs*, Graphs Combin. **30** (2014) 1087–1098.
doi:10.1007/s00373-013-1331-y
- [7] A. Brandt, M. Ferrara, M. Kumbhat, S. Loeb, D. Stolee and M. Yancey, *I, F -partitions of sparse graphs*, European J. Combin. **57** (2016) 1–12.
doi:10.1016/j.ejc.2016.03.003
- [8] Y. Bu, D. Cranston, M. Montassier, A. Raspaud and W. Wang, *Star coloring of sparse graphs*, J. Graph Theory **62** (2009) 201–219.
doi:10.1002/jgt.20392
- [9] G. Chartrand, F. Okamoto and P. Zhang, *The sigma chromatic number of a graph*, Graphs Combin. **26** (2010) 755–773.
doi:10.1007/s00373-010-0952-7
- [10] D.W. Cranston and D.B. West, *An introduction to the discharging method via graph coloring*, Discrete Math. **340** (2017) 766–793.
doi:10.1016/j.disc.2016.11.022
- [11] S. Czerwiński, J. Grytczuk and W. Żelazny, *Lucky labelings of graphs*, Inform. Process. Lett. **109** (2009) 1078–1081.
doi:10.1016/j.ipl.2009.05.011

- [12] A. Dehghan, M. Sadeghi and A. Ahadi, *The complexity of the sigma chromatic number of cubic graphs*, (2014).
arXiv:math.CO/1403.6288v1
- [13] A. Dehghan, M. Sadeghi and A. Ahadi, *Algorithmic complexity of proper labeling problems*, Theoret. Comput. Sci. **495** (2013) 25–36.
doi:10.1016/j.tcs.2013.05.027
- [14] H. Grötzsch, *Zur Theorie der diskreten Gebilde. VII. Ein Dreifarbensatz für dreikreisfreie Netze auf der Kugel*, Wiss. Z. Martin-Luther-Univ. Halle-Wittenberg. Math.-Nat. Reihe **8** (1958/1959) 109–120.
- [15] M. Kalkowski, M. Karoński and F. Pfender, *Vertex-coloring edge-weightings: towards the 1-2-3-conjecture*, J. Combin. Theory Ser. B **100** (2010) 347–349.
doi:10.1016/j.jctb.2009.06.002
- [16] M. Karoński, T. Łuczak and A. Thomason, *Edge weights and vertex colours*, J. Combin. Theory Ser. B **91** (2004) 151–157.
doi:10.1016/j.jctb.2003.12.001
- [17] B. Seamone, *The 1-2-3 conjecture and related problems: a survey* (2012).
arXiv:math.CO/1211.5122
- [18] B. Seamone, *Derived Colourings of Graphs* (PhD Thesis, Carleton University, 2012).
- [19] R. Steinberg and C.A. Tovey, *Planar Ramsey numbers*, J. Combin. Theory Ser. B **59** (1993) 288–296.
doi:10.1006/jctb.1993.1070
- [20] C. Thomassen, *A short list color proof of Grötzsch’s theorem*, J. Combin. Theory Ser. B **88** (2003) 189–192.
doi:10.1016/S0095-8956(03)00029-7
- [21] D.B. West, *Introduction to Graph Theory* (Prentice Hall, Inc., Upper Saddle River, NJ, 1996).

Received 29 May 2017

Revised 18 May 2018

Accepted 18 May 2018