

THE DOUBLE ROMAN DOMATIC NUMBER OF A DIGRAPH

LUTZ VOLKMANN

*Lehrstuhl II für Mathematik
RWTH Aachen University
52056 Aachen, Germany*

e-mail: volkm@math2.rwth-aachen.de

Abstract

A *double Roman dominating function* on a digraph D with vertex set $V(D)$ is defined in [G. Hao, X. Chen and L. Volkmann, *Double Roman domination in digraphs*, Bull. Malays. Math. Sci. Soc. (2017).] as a function $f : V(D) \rightarrow \{0, 1, 2, 3\}$ having the property that if $f(v) = 0$, then the vertex v must have at least two in-neighbors assigned 2 under f or one in-neighbor w with $f(w) = 3$, and if $f(v) = 1$, then the vertex v must have at least one in-neighbor u with $f(u) \geq 2$. A set $\{f_1, f_2, \dots, f_d\}$ of distinct double Roman dominating functions on D with the property that $\sum_{i=1}^d f_i(v) \leq 3$ for each $v \in V(D)$ is called a *double Roman dominating family* (of functions) on D . The maximum number of functions in a double Roman dominating family on D is the *double Roman domatic number* of D , denoted by $d_{dR}(D)$. We initiate the study of the double Roman domatic number, and we present different sharp bounds on $d_{dR}(D)$. In addition, we determine the double Roman domatic number of some classes of digraphs.

Keywords: digraph, double Roman domination, double Roman domatic number.

2010 Mathematics Subject Classification: 05C20, 05C40.

REFERENCES

- [1] H. Abdollahzadeh Ahangar, J. Amjadi, M. Atapour, M. Chellali and S.M. Sheikholeslami, *Double Roman trees*, Ars Combin., to appear.
- [2] H. Abdollahzadeh Ahangar, J. Amjadi, M. Chellali, S. Nazari-Moghaddam and S.M. Sheikholeslami, *Trees with double Roman domination number twice the domination number plus two*, Iran. J. Sci. Technol. Trans. A Sci. (2018), in press.
doi:10.1007/s40995-018-0535-7

- [3] H. Abdollahzadeh Ahangar, M. Chellali and S.M. Sheikholeslami, *On the double Roman domination in graphs*, Discrete Appl. Math. **232** (2017) 1–7.
doi:10.1016/j.dam.2017.06.014
- [4] R.A. Beeler, T.W. Haynes and S.T. Hedetniemi, *Double Roman domination*, Discrete Appl. Math. **211** (2016) 23–29.
doi:10.1016/j.dam.2016.03.017
- [5] G. Hao, X. Chen and L. Volkmann, *Double Roman domination in digraphs*, Bull. Malays. Math. Sci. Soc. (2017), in press.
doi:10.1007/s40840-017-0582-9
- [6] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, *Fundamentals of Domination in Graphs* (Marcel Dekker, Inc., New York, 1998).
- [7] N. Jafari Rad and H. Rahbani, *Some progress on double Roman domination in graphs*, Discuss. Math. Graph Theory, in press.
doi:10.7151/dmgt.2069
- [8] E.A. Nordhaus and J.W. Gaddum, *On complementary graphs*, Amer. Math. Monthly **63** (1956) 175–177.
doi:10.2307/2306658
- [9] L. Volkmann, *The double Roman domatic number of a graph*, J. Combin. Math. Combin. Comput. **104** (2018) 205–215.
- [10] B. Zelinka, *Domatic number and degrees of vertices of a graph*, Math. Slovaca **33** (1983) 145–147.
- [11] X. Zhang, Z. Li, H. Jiang and Z. Shao, *Double Roman domination in trees*, Inform. Process. Lett. **134** (2018) 31–34.
doi:10.1016/j.ipl.2018.01.004

Received 8 March 2018

Revised 15 June 2018

Accepted 15 June 2018