

LINEAR LIST COLORING OF SOME SPARSE GRAPHS

MING CHEN

College of Mathematics Physics and Information Engineering
Jiaxing University, Zhejiang 314001, China

e-mail: chen2001ming@163.com

YUSHENG LI

AND

LI ZHANG¹

School of Mathematical Sciences
Tongji University Shanghai 200092, China

e-mail: li_yusheng@tongji.edu.cn
lizhang@tongji.edu.cn

Abstract

A linear k -coloring of a graph is a proper k -coloring of the graph such that any subgraph induced by the vertices of any pair of color classes is a union of vertex-disjoint paths. A graph G is linearly L -colorable if there is a linear coloring c of G for a given list assignment $L = \{L(v) : v \in V(G)\}$ such that $c(v) \in L(v)$ for all $v \in V(G)$, and G is linearly k -choosable if G is linearly L -colorable for any list assignment with $|L(v)| \geq k$. The smallest integer k such that G is linearly k -choosable is called the linear list chromatic number, denoted by $lc_l(G)$. It is clear that $lc_l(G) \geq \left\lceil \frac{\Delta(G)}{2} \right\rceil + 1$ for any graph G with maximum degree $\Delta(G)$. The maximum average degree of a graph G , denoted by $mad(G)$, is the maximum of the average degrees of all subgraphs of G . In this note, we shall prove the following. Let G be a graph, (1) if $mad(G) < \frac{8}{3}$ and $\Delta(G) \geq 7$, then $lc_l(G) = \left\lceil \frac{\Delta(G)}{2} \right\rceil + 1$; (2) if $mad(G) < \frac{18}{7}$ and $\Delta(G) \geq 5$, then $lc_l(G) = \left\lceil \frac{\Delta(G)}{2} \right\rceil + 1$; (3) if $mad(G) < \frac{20}{7}$ and $\Delta(G) \geq 5$, then $lc_l(G) \leq \left\lceil \frac{\Delta(G)}{2} \right\rceil + 2$.

Keywords: linear coloring, maximum average degree, planar graphs, discharging.

2010 Mathematics Subject Classification: 05C15.

¹Corresponding author.

REFERENCES

- [1] D.W. Cranston and G. Yu, *Linear choosability of sparse graphs*, Discrete Math. **311** (2011) 1910–1917.
doi:10.1016/j.disc.2011.05.017
- [2] W. Dong, B. Xu and X. Zhang, *Improved bounds on linear coloring of plane graphs*, Sci. China Math. **53** (2010) 1895–1902.
doi:10.1007/s11425-010-3073-0
- [3] W. Dong and W. Lin, *On linear coloring of planar graphs with small girth*, Discrete Appl. Math. **173** (2014) 35–44.
doi:10.1016/j.dam.2014.03.019
- [4] L. Esperet, M. Montassier and A. Raspaud, *Linear choosability of graphs*, Discrete Math. **308** (2008) 3938–3950.
doi:10.1016/j.disc.2007.07.112
- [5] C. Li, W. Wang and A. Raspaud, *Upper bounds on the linear chromatic number of a graph*, Discrete Math. **311** (2011) 232–238.
doi:10.1016/j.disc.2010.10.023
- [6] C.H. Liu and G. Yu, *Linear colorings of subcubic graphs*, European J. Combin. **34** (2013) 1040–1050.
doi:10.1016/j.ejc.2013.02.008
- [7] Y. Wang and Q. Wu, *Linear coloring of sparse graphs*, Discrete Appl. Math. **160** (2012) 664–672.
doi:10.1016/j.dam.2011.10.028
- [8] R. Yuster, *Linear coloring of graphs*, Discrete Math. **185** (1998) 293–297.
doi:10.1016/S0012-365X(97)00209-4

Received 12 May 2017
 Revised 31 January 2018
 Accepted 31 July 2018