Discussiones Mathematicae Graph Theory 41 (2021) 905–921 https://doi.org/10.7151/dmgt.2219

Full PDF DMGT Page

THE LAGRANGIAN DENSITY OF {123, 234, 456} AND THE TURÁN NUMBER OF ITS EXTENSION

PINGGE CHEN

College of Science Hunan University of Technology e-mail: chenpingge@hnu.edu.cn

JINHUA LIANG

College of Mathematics and Econometrics Hunan University

e-mail: jh_liang@hnu.edu.cn

AND

YUEJIAN PENG¹

Institute of Mathematics Hunan University

e-mail: ypeng1@hnu.edu.cn

Abstract

Given a positive integer n and an r-uniform hypergraph F, the Turán number ex(n, F) is the maximum number of edges in an F-free r-uniform hypergraph on n vertices. The Turán density of F is defined as $\pi(F) = \lim_{n\to\infty} \frac{ex(n,F)}{\binom{n}{r}}$. The Lagrangian density of F is $\pi_{\lambda}(F) = \sup\{r!\lambda(G) : G$ is F-free}, where $\lambda(G)$ is the Lagrangian of G. Sidorenko observed that $\pi(F) \leq \pi_{\lambda}(F)$, and Pikhurko observed that $\pi(F) = \pi_{\lambda}(F)$ if every pair of vertices in F is contained in an edge of F. Recently, Lagrangian densities of hypergraphs and Turán numbers of their extensions have been studied actively. For example, in the paper [A hypergraph Turán theorem via Lagrangians of intersecting families, J. Combin. Theory Ser. A 120 (2013) 2020–2038], Hefetz and Keevash studied the Lagrangian density of the 3uniform graph spanned by {123, 456} and the Turán number of its extension. In this paper, we show that the Lagrangian density of the 3-uniform graph

¹Corresponding author.

spanned by $\{123, 234, 456\}$ achieves only on K_5^3 . Applying it, we get the Turán number of its extension, and show that the unique extremal hypergraph is the balanced complete 5-partite 3-uniform hypergraph on n vertices.

Keywords: Turán number, hypergraph Lagrangian, Lagrangian density. 2010 Mathematics Subject Classification: 05C65.

References

- A. Brandt, D. Irwin and T. Jiang, Stability and Turán numbers of a class of hypergraphs via Lagrangians, Combin. Probab. Comput. 26 (2017) 367–405. https://doi.org/10.1017/S0963548316000444
- [2] P. Chen, J. Liang and Y. Peng, Lagrangian density of {123, 234, 567} and Turán numbers of its extension, submitted.
- [3] P. Frankl and Z. Füredi, Extremal problems whose solutions are the blowups of the small witt-designs, J. Combin. Theory Ser. A 52 (1989) 129–147. https://doi.org/10.1016/0097-3165(89)90067-8
- P. Frankl and V. Rödl, Hypergraphs do not jump, Combinatorica 4 (1984) 149–159. https://doi.org/10.1007/BF02579215
- [5] D. Hefetz and P. Keevash, A hypergraph Turán theorem via lagrangians of intersecting families, J. Combin. Theory Ser. A 120 (2013) 2020–2038. https://doi.org/10.1016/j.jcta.2013.07.011
- [6] S. Hu, Y. Peng and B. Wu, Lagrangian densities of the disjoint union of 3-uniform linear paths and matchings and Turán numbers of their extensions, submitted.
- [7] T. Jiang, Y. Peng and B. Wu, Lagrangian densities of some sparse hypergraphs and Turán numbers of their extensions, European J. Combin. 73 (2018) 20–36. https://doi.org/10.1016/j.ejc.2018.05.001
- [8] G. Katona, T. Nemetz and M. Simonovits, On a problem of Turán in the theory of graphs, Mat. Lapok (N.S.) 15 (1964) 228–238.
- P. Keevash, Hypergrah Turán problems, in: Surveys in Combinatorics 2011, London Math. Soc. Lecture Note Ser. 392 (Cambridge Univ. Press, Cambridge, 2011) 83–140. https://doi.org/10.1017/CB09781139004114
- [10] T.S. Motzkin and E.G. Straus, Maxima for graphs and a new proof of a theorem of Turán, Canad. J. Math. 17 (1965) 533–540. https://doi.org/10.4153/CJM-1965-053-6
- [11] D. Mubayi, A hypergraph extension of Turán's theorem, J. Combin. Theory Ser. B 96 (2006) 122–134. https://doi.org/10.1016/j.jctb.2005.06.013

- D. Mubayi and O. Pikhurko, A new generalization of Mantel's theorem to k-graphs, J. Combin. Theory Ser. B 97 (2007) 669–678. https://doi.org/10.1016/j.jctb.2006.11.003
- S. Norin and L. Yepremyan, Turán number of generalized triangles, J. Combin. Theory Ser. A 146 (2017) 312–343. https://doi.org/10.1016/j.jcta.2016.09.003
- [14] S. Norin and L. Yepremyan, *Turán numbers of extensions*, J. Combin. Theory Ser. A 155 (2018) 476–492. https://doi.org/10.1016/j.jcta.2017.08.004
- [15] O. Pikhurko, An exact Turán result for the generalized triangle, Combinatorica 28 (2008) 187–208. https://doi.org/10.1007/s00493-008-2187-2
- [16] O. Pikhurko, Exact computation of the hypergraph Turán function for expanded complete 2-graphs, J. Combin. Theory Ser. B 103 (2013) 220–225. https://doi.org/10.1016/j.jctb.2012.09.005
- [17] A.F. Sidorenko, The maximal number of edges in a homogeneous hypergraph containing no prohibited subgraphs, Mat. Zametki 41 (1987) 433–455. https://doi.org/10.1027.BF01158259
- [18] A.F. Sidorenko, Asymptotic solution for a new class of forbidden r-graphs, Combinatorica 9 (1989) 207–215. https://doi.org/10.1007/BF02124681
- [19] B. Wu and Y. Peng, Lagrangian densities of 3-uniform linear paths and Turán numbers of their extensions, submitted.
- [20] B. Wu, Y. Peng and P. Chen, On a conjecture of Hefetz and Keevash on Lagrangians of intersecting hypergraphs and Turán numbers. arXiv:1701.06126v3
- [21] Z. Yan and Y. Peng, λ-perfect hypergraphs and Lagrangian densities of hypergraph cycles, Discrete Math. (2019). https://doi.org/10.1016/j.disc.2019.03.024
- [22] A.A. Zykov, On some properties of linear complexes, Mat. Sb. 24 (1949) 163–188.

Received 4 August 2017 Revised 18 March 2019 Accepted 18 March 2019