Discussiones Mathematicae
Graph Theory 41 (2021) 1199–1213
https://doi.org/10.7151/dmgt.2269

THE GENERAL POSITION PROBLEM ON KNESER GRAPHS AND ON SOME GRAPH OPERATIONS

Modjtaba Ghorbani, Hamid Reza Maimani

Mostafa Momeni, Farhad Rahimi Mahid

Department of Mathematics, Faculty of Science
Shahid Rajaee Teacher Training University
Tehran, 16785 - 136, I.R. Iran

e-mail: ghorbani30@gmail.com
maimani@ipm.ir
momeni.mosi@yahoo.com
farhad.rahimi@sru.ac.ir

Sandi Klavžar

Faculty of Mathematics and Physics, University of Ljubljana, Slovenia
Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia
Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia

e-mail: sandi.klavzar@fmf.uni-lj.si

and

Gregor Rus

Faculty of Organizational Sciences, University of Maribor, Slovenia
Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia

e-mail: gregor.rus4@um.si

Abstract

A vertex subset S of a graph G is a general position set of G if no vertex of S lies on a geodesic between two other vertices of S. The cardinality of a largest general position set of G is the general position number (gp-number) $\mathrm{gp}(G)$ of G. The gp-number is determined for some families of Kneser graphs, in particular for $K(n,2)$, $n \geq 4$, and $K(n,3)$, $n \geq 9$. A sharp lower bound on the gp-number is proved for Cartesian products of graphs. The gp-number is also determined for joins of graphs, coronas over graphs, and line graphs of complete graphs.

Keywords: general position set, Kneser graphs, Cartesian product of graphs, corona over graphs, line graphs.

2010 Mathematics Subject Classification: 05C12, 05C69, 05C76.

References

[1] B.S. Anand, S.V. Ullas Chandran, M. Changat, S. Klavžar and E.J. Thomas, *Characterization of general position sets and its applications to cographs and bipartite graphs*, Appl. Math. Comput. **359** (2019) 84–89.
https://doi.org/10.1016/j.amc.2019.04.064

[2] G. Boruzanli Ekinci and J.B. Gauci, *The super-connectivity of Kneser graphs*, Discuss. Math. Graph Theory **39** (2019) 5–11.
https://doi.org/10.7151/dmgt.2051

[3] B. Brešar and M. Valencia-Pabon, *Independence number of products of Kneser graphs*, Discrete Math. **342** (2019) 1017–1027.
https://doi.org/10.1016/j.disc.2018.12.017

[4] K.N. Chadha and A.A. Kulkarni, *On independent cliques and linear complementarity problems* (2018).
arXiv:1811.09798

[5] H.E. Dudeney, Amusements in Mathematics (Nelson, Edinburgh, 1917).

[6] P. Erdős, C. Ko and R. Rado, *Intersection theorems for systems of finite sets*, Q. J. Math. **12** (1961) 313–320.
https://doi.org/10.1093/qmath/12.1.313

[7] V. Froese, I. Kanj, A. Nichterlein and R. Niedermeier, *Finding points in general position*, Internat. J. Comput. Geom. Appl. **27** (2017) 277–296.
https://doi.org/10.1142/S021819591750008X

[8] W. Imrich, S. Klavžar and D.F. Rall, Topics in Graph Theory: Graphs and Their Cartesian Product (A K Peters, New York 2008).
https://doi.org/10.1201/b10613

[9] M.M. Kanté, R.M. Sampaio, V.F. dos Santos and J.L. Szwarcfiter, *On the geodetic rank of a graph*, J. Comb. **8** (2017) 323–340.
https://doi.org/10.4310/JOC.2017.v8.n2.a5

[10] S. Klavžar, B. Patkós, G. Rus and I.G. Yero, *On general position sets in Cartesian grids* (2019).
arXiv:1907.04535v3

[11] S. Klavžar and I.G. Yero, *The general position problem and strong resolving graphs*, Open Math. **17** (2019) 1126–1135.
https://doi.org/10.1515/math-2019-0088

[12] C.Y. Ku and K.B. Wong, *On no-three-in-line problem on m-dimensional torus*, Graphs Combin. **34** (2018) 355–364.
https://doi.org/10.1007/s00373-018-1878-8

[13] J.H. van Lint and R.M. Wilson, A Course in Combinatorics (Cambridge University Press, Cambridge, 1992).
https://doi.org/10.1017/CBO9780511987045

[14] P. Manuel and S. Klavžar, *A general position problem in graph theory*, Bull. Aust. Math. Soc. **98** (2018) 177–187.
https://doi.org/10.1017/S0004972718000473

[15] P. Manuel and S. Klavžar, *The graph theory general position problem on some interconnection networks*, Fund. Inform. **163** (2018) 339–350.
https://doi.org/10.3233/FI-2018-1748

[16] A. Misiak, Z. Stępień, A. Szymaszkiewicz, L. Szymaszkiewicz and M. Zwierzchowski, *A note on the no-three-in-line problem on a torus*, Discrete Math. **339** (2016) 217–221.
https://doi.org/10.1016/j.disc.2015.08.006

[17] T. Mütze and P. Su, *Bipartite Kneser graphs are Hamiltonian*, Combinatorica **37** (2017) 1207–1219.
https://doi.org/10.1007/s00493-016-3434-6

[18] B. Patkós, *On the general position problem on Kneser graphs* (2019).
arXiv:1903.08056v2

[19] M. Payne and D.R. Wood, *On the general position subset selection problem*, SIAM J. Discrete Math. **27** (2013) 1727–1733.
https://doi.org/10.1137/120897493

[20] A. Por and D.R. Wood, *No-three-in-line-in-3D*, Algorithmica **47** (2007) 481–488.
https://doi.org/10.1007/s00453-006-0158-9

[21] S.V. Ullas Chandran and G. Jaya Parthasarathy, *The geodesic irredundant sets in graphs*, Int. J. Math. Combin. **4** (2016) 135–143.
https://doi.org/10.5281/zenodo.826834

[22] M. Valencia-Pabon and J.-C. Vera, *On the diameter of Kneser graphs*, Discrete Math. **305** (2005) 383–385.
https://doi.org/10.1016/j.disc.2005.10.001

Received 11 March 2019
Revised 13 November 2019
Accepted 20 November 2019