

BALANCEDNESS AND THE LEAST LAPLACIAN EIGENVALUE OF SOME COMPLEX UNIT GAIN GRAPHS

FRANCESCO BELARDO

MAURIZIO BRUNETTI

*Department of Mathematics and Applications
University ‘Federico II’, Naples, Italy*

e-mail: fbelardo@unina.it, mbrunett@unina.it

AND

NATHAN REFF

*Department of Mathematics
The College at Brockport: State University of New York
Brockport, NY 14420, USA*

e-mail: nreff@brockport.edu

Abstract

Let $\mathbb{T}_4 = \{\pm 1, \pm i\}$ be the subgroup of 4-th roots of unity inside $\mathbb{T}$, the multiplicative group of complex units. A complex unit gain graph Φ is a simple graph $\Gamma = (V(\Gamma) = \{v_1, \dots, v_n\}, E(\Gamma))$ equipped with a map $\varphi : \vec{E}(\Gamma) \rightarrow \mathbb{T}$ defined on the set of oriented edges such that $\varphi(v_i v_j) = \varphi(v_j v_i)^{-1}$. The gain graph Φ is said to be balanced if for every cycle $C = v_{i_1} v_{i_2} \cdots v_{i_k} v_{i_1}$ we have $\varphi(v_{i_1} v_{i_2}) \varphi(v_{i_2} v_{i_3}) \cdots \varphi(v_{i_k} v_{i_1}) = 1$.

It is known that Φ is balanced if and only if the least Laplacian eigenvalue $\lambda_n(\Phi)$ is 0. Here we show that, if Φ is unbalanced and $\varphi(\Phi) \subseteq \mathbb{T}_4$, the eigenvalue $\lambda_n(\Phi)$ measures how far is Φ from being balanced. More precisely, let $\nu(\Phi)$ (respectively, $\epsilon(\Phi)$) be the number of vertices (respectively, edges) to cancel in order to get a balanced gain subgraph. We show that

$$\lambda_n(\Phi) \leq \nu(\Phi) \leq \epsilon(\Phi).$$

We also analyze the case when $\lambda_n(\Phi) = \nu(\Phi)$. In fact, we identify the structural conditions on Φ that lead to such equality.

Keywords: gain graph, Laplacian eigenvalues, balanced graph, algebraic frustration.

2010 Mathematics Subject Classification: 05C50, 05C22.

REFERENCES

- [1] R.B. Bapat, D. Kalita and S. Pati, *On weighted directed graphs*, Linear Algebra Appl. **436** (2012) 99–111.
doi:10.1016/j.laa.2011.06.035
- [2] F. Belardo, *Balancedness and the least eigenvalue of Laplacian of signed graphs*, Linear Algebra Appl. **446** (2014) 133–147.
doi:10.1016/j.laa.2014.01.001
- [3] S. Fallat and Y.-Z. Fan, *Bipartiteness and the least eigenvalue of signless Laplacian of graphs*, Linear Algebra Appl. **436** (2012) 3254–3267.
doi:10.1016/j.laa.2011.11.015
- [4] K. Guo and B. Mohar, *Hermitian adjacency matrix of digraphs and mixed graphs*, J. Graph Theory **85** (2017) 217–248.
doi:10.1002/jgt.22057
- [5] R.A. Horn and C. R. Johnson, *Matrix Analysis* (Cambridge Univ. Press, New York, 2012).
doi:10.1017/CBO9781139020411
- [6] D. Kalita, *Properties of first eigenvectors and first eigenvalues of nonsingular weighted directed graphs*, Electron. J. Linear Algebra **30** (2015) 227–242.
doi:10.13001/1081-3810.3029
- [7] N. Reff, *Spectral properties of complex unit gain graphs*, Linear Algebra Appl. **436** (2012) 3165–3176.
doi:10.1016/j.laa.2011.10.021
- [8] N. Reff, *Oriented gain graphs, line graphs and eigenvalues*, Linear Algebra Appl. **506** (2016) 316–328.
doi:10.1016/j.laa.2016.05.040
- [9] Y.-Y. Tan and Y.Z. Fan, *On edge singularity and eigenvectors of mixed graphs*, Acta Math. Sin. (Engl. Ser.) **24** (2008) 139–146.
doi:10.1007/s10114-007-1000-2
- [10] Y. Wang, S.-C. Gong and Y.-Z. Fan, *On the determinant of the Laplacian matrix of a complex unit gain graph*, Discrete Math. **341** (2018) 81–86.
doi:10.1016/j.disc.2017.07.003
- [11] T. Zaslavsky, *Biased graphs. I. Bias, balance, and gains*, J. Combin. Theory Ser. B **47** (1989) 32–52.
doi:10.1016/0095-8956(89)90063-4
- [12] T. Zaslavsky, *A mathematical bibliography of signed and gain graphs and allied areas*, Electron. J. Combin. (1998) #DS8.

Received 11 June 2019

Revised 15 October 2019

Accepted 24 October 2019