

Discussiones Mathematicae
Graph Theory 24 (2004) 545–549

LUCIEN HADDAD AND CLAUDE TARDIF

Department of Mathematics and Computer Science
Royal Military College of Canada
PO Box 17000, Station "Forces"
Kingston, Ontario K7K 7B4 Canada

Abstract

The Erdős-Faber-Lovász conjecture states that if a graph G is the union of n cliques of size n no two of which share more than one vertex, then $\chi(G) = n$. We provide a formulation of this conjecture in terms of maximal partial clones of partial operations on a set.

Keywords: chromatic number, Erdős-Faber-Lovász conjecture, maximal partial clones.

2000 Mathematics Subject Classification: 05C15 (primary: graph colouring) 08A55 (secondary: Partial algebras).

1. Introduction

Suppose that there are n committees each with n members each, such that each pair of committees has at most one member in common. The committees hold their meetings in the committee room which has n chairs. Is it then possible for every person to select a chair which he/she will use in all the meetings of all the committees to which he/she belongs? The Erdős-Faber-Lovász conjecture states that the answer is yes; in graph theoretic terms, the conjecture can be restated as follows:

If a graph G is the union of n cliques of size n no two of which share more than one vertex, then $\chi(G) = n$

(see [1, 8, 9]). The n constituent n -cliques of such an instance of the Erdős-Faber-Lovász conjecture can be viewed as the hyperedges in a hypergraph, hence the conjecture also admits a formulation in terms of strong colouring of hypergraphs:

If an n -uniform hypergraph \mathcal{H} has exactly n hyperedges no two of which share more than one point, then the strong chromatic number of \mathcal{H} is n .

A hypergraph which satisfies the hypotheses of the Erdős-Faber-Lovász conjecture will be called an *instance* of the conjecture. Given such an instance \mathcal{H} , we can define the relational structure $(V_{\mathcal{H}}, R_{\mathcal{H}})$ whose base set $V_{\mathcal{H}}$ is the same as that of \mathcal{H} , where $R_{\mathcal{H}}$ is the n -ary relational structure consisting of all the n -tuples (x_1, x_2, \dots, x_n) such that $\{x_1, x_2, \dots, x_n\}$ is an hyperedge of \mathcal{H} . Thus if \mathcal{H} has n hyperedges, then $R_{\mathcal{H}}$ consists of the $n \cdot n!$ n -tuples obtained by linearly ordering each hyperedge of \mathcal{H} . The reason for considering $(V_{\mathcal{H}}, R_{\mathcal{H}})$ is that it allows to define products, homomorphisms and partial homomorphisms, and reinterpret the Erdős-Faber-Lovász conjecture from the point of view of partial clone theory. We prove the following.

Theorem 1. *The Erdős-Faber-Lovász conjecture is true if and only if for every instance \mathcal{H} of the conjecture, the relational structure $(V_{\mathcal{H}}, R_{\mathcal{H}})$ determines a maximal partial clone.*

In the next section we introduce the necessary definitions and results needed to prove Theorem 1.

2. Partial Operations and Partial Clones

An n -ary relational structure is a couple (V, R) where V is a set and $R \subseteq V^n$. It is called *areflexive* if for every $(x_1, \dots, x_n) \in R$, the elements x_1, \dots, x_n are all distinct, and *totally symmetric* if for every $(x_1, \dots, x_n) \in R$ and every permutation π of $\{1, \dots, n\}$, we have $(x_{\pi(1)}, \dots, x_{\pi(n)}) \in R$. A *homomorphism* between two n -ary relational structures (V, R) and (V', R') is a map ϕ from V to V' such that $(\phi(x_1), \dots, \phi(x_n)) \in R'$ for all $(x_1, \dots, x_n) \in R$. A map ϕ from a subset dom_{ϕ} of V to V' is called a *partial homomorphism* if $(\phi(x_1), \dots, \phi(x_n)) \in R'$ for all $x_1, \dots, x_n \in \text{dom}_{\phi}$ such that $(x_1, \dots, x_n) \in R$.

For example, instances \mathcal{H} , \mathcal{H}' of the Erdős-Faber-Lovász conjecture give rise to the areflexive, totally symmetric n -ary relational structures

$(V_{\mathcal{H}}, R_{\mathcal{H}})$, $(V_{\mathcal{H}'}, R_{\mathcal{H}'})$, and also to totally symmetric binary relational structures, namely the corresponding graphs G and G' . Since the elements of $R_{\mathcal{H}}$ and $R_{\mathcal{H}'}$ correspond to cliques in G and G' respectively, a homomorphism from $(V_{\mathcal{H}}, R_{\mathcal{H}})$ to $(V_{\mathcal{H}'}, R_{\mathcal{H}'})$ naturally induces a homomorphism from G to G' . However the converse does not necessarily hold. For instance G may be n -colourable (as the conjecture claims) and G' may contain a n -clique C which does not correspond to any hyperedge of \mathcal{H}' . Identifying the elements of C with the colours in a n -colouring of G defines a homomorphism from G to G' which does not correspond to any homomorphism from $(V_{\mathcal{H}}, R_{\mathcal{H}})$ to $(V_{\mathcal{H}'}, R_{\mathcal{H}'})$. Now the partial homomorphisms from $(V_{\mathcal{H}}, R_{\mathcal{H}})$ to $(V_{\mathcal{H}'}, R_{\mathcal{H}'})$ need not even induce partial homomorphisms from G to G' . For instance every hyperedge of \mathcal{H} contains a vertex which does not belong to any other hyperedge. Removing such a vertex from every hyperedge yields a subset X of $V_{\mathcal{H}}$ which does not contain any hyperedge. Thus *any* map from X to $V_{\mathcal{H}'}$ is a partial homomorphism from $(V_{\mathcal{H}}, R_{\mathcal{H}})$ to $(V_{\mathcal{H}'}, R_{\mathcal{H}'})$, while the partial homomorphisms from G to G' with domain X need to preserve the edges of the subgraph induced by X . Thus from the point of view of partial homomorphisms, the n -ary relational structures induced by instances of the Erdős-Faber-Lovász conjecture do not behave like the corresponding graphs.

Given an integer m , the m -th power $(V, R)^m$ of a n -ary relational structure (V, R) is the n -ary relational structure (V^m, R') , where

$$R' = \{(X_1, \dots, X_n) \in (V^m)^n : (\text{pr}_i(X_1), \dots, \text{pr}_i(X_n)) \in R, i = 1, \dots, m\}$$

(where pr_i is the projection given by $\text{pr}_i(a_1, \dots, a_m) = a_i$). A partial function from a power of V to V is called a *partial operation* on V . Let $pPol(V, R)$ denote the set of partial operations on V which are partial homomorphisms from some power of (V, R) to (V, R) . Note that $pPol(V, R)$ contains all the projections, and is closed under the following composition:

If ϕ_1, \dots, ϕ_k are partial homomorphisms from $(V, R)^m$ to (V, R) and ψ a partial homomorphism from $(V, R)^k$ to (V, R) , then $\psi(\phi_1, \dots, \phi_k)$ is the partial homomorphism ψ' given by

$$\psi'(X_1, \dots, X_m) = \psi(\phi_1(X_1, \dots, X_m), \dots, \phi_k(X_1, \dots, X_m)),$$

on the domain consisting of the m -tuples (X_1, \dots, X_m) for which the above expression is well defined. A set of partial operations which contains all projections and is closed under composition is called a *partial clone*. The family of partial clones on a set V , ordered by inclusion, forms a dually atomic lattice. A partial clone is called *maximal* if it is not properly contained in any other partial clone, apart from the set of all partial operations on V . In other words, a partial clone \mathcal{C} on a set V is maximal if for any two partial operations f, g on V , neither one in \mathcal{C} , g can be obtained by composition using only f and elements of \mathcal{C} . The characterization of maximal partial clones on a finite set resembles that of maximal clones and generating sets of boolean operations, though the list is far more complex in the partial case. (see [2, 3, 4, 5, 6, 7]). In particular, the following is known:

Theorem 2 ([5]). *Let (V, R) be an n -ary reflexive, totally symmetric relational structure. Then $pPol(V, R)$ is a maximal partial clone if and only if (V, R) admits a strong n -colouring.*

Here, a *strong n -colouring* of (V, R) is just a strong n -colouring of the hypergraph whose hyperedges are the sets $\{x_1, \dots, x_n\}$ such that $(x_1, \dots, x_n) \in R$. Thus we have the following.

Corollary 3. *Let \mathcal{H} be an instance of the Erdős-Faber-Lovász conjecture. Then \mathcal{H} admits a strong n -colouring if and only if $pPol(V_{\mathcal{H}}, R_{\mathcal{H}})$ is a maximal partial clone.*

Theorem 1 follows directly from this corollary. Note that although this partial clone theoretic formulation uses the context of the Erdős-Faber-Lovász conjecture, the hypothesis that the hyperedges be almost disjoint is not necessary for the correspondence. It would be interesting to see whether this hypothesis can be incorporated to a partial clone-theoretic treatment of the subject.

References

- [1] P. Erdős, *On the Combinatorial problems which I would most like to see solved*, *Combinatorica* **1** (1981) 25–42.
- [2] L. Haddad, *Le treillis des clones partiels sur un univers fini et ses coatomes* (Ph. D. thesis, Université de Montréal, 1986).
- [3] L. Haddad and I.G. Rosenberg, *Critère général de complétude pour les algèbres partielles finies*, *C.R. Acad. Sci. Paris, tome 304, Série I*, **17** (1987) 507–509.
- [4] L. Haddad, *Maximal partial clones determined by quasi-diagonal relations*, *J. Inf. Process. Cybern. EIK* **24** (1988) 7/8 355–366.
- [5] L. Haddad and I.G. Rosenberg, *Maximal partial clones determined by areflexive relations*, *Discrete Appl. Math.* **24** (1989) 133–143.
- [6] L. Haddad, I.G. Rosenberg and D. Schweigert, *A maximal partial clone and a Skupecki-type criterion*, *Acta Sci. Math.* **54** (1990) 89–98.
- [7] L. Haddad and I.G. Rosenberg, *Completeness theory for finite partial algebras*, *Algebra Universalis* **29** (1992) 378–401.
- [8] T.R. Jensen and B. Toft, *Graph Coloring Problems* (Wiley-Interscience series in discrete mathematics and optimization, John Wiley & Sons Inc., 1995).
- [9] J. Kahn, *Coloring nearly-disjoint hypergraphs with $n + o(n)$ colors*, *J. Combin. Theory (A)* **59** (1992) 31–39.

Received 12 July 2004