

EDGE MAXIMAL C_{2k+1} -EDGE DISJOINT FREE GRAPHS

M.S.A. BATAINEH

Department of Mathematics
Yarmouk University
Irbid-Jordan

e-mail: bataineh71@hotmail.com

AND

M.M.M. JARADAT

Yarmouk University
Department of Mathematics
Irbid-Jordan
Department of Mathematics, Physics and Statistics
Qatar University
Doha-Qatar

e-mail: mmjst4@yu.edu.jo; mmjst4@qu.edu.qa

Abstract

For two positive integers r and s , $\mathcal{G}(n; r, s)$ denotes to the class of graphs on n vertices containing no r of s -edge disjoint cycles and $f(n; r, s) = \max\{|\mathcal{E}(G)| : G \in \mathcal{G}(n; r, s)\}$. In this paper, for integers $r \geq 2$ and $k \geq 1$, we determine $f(n; r, 2k + 1)$ and characterize the edge maximal members in $\mathcal{G}(n; r, 2k + 1)$.

Keywords: extremal graphs, edge disjoint, cycles.

2010 Mathematics Subject Classification: 05C38, 05C35.

1. INTRODUCTION

The graphs considered in this paper are finite, undirected and have no loops or multiple edges. Most of the notations that follow can be found in [5]. For a given graph G , we denote the vertex set of a graph G by $V(G)$ and the edge set by $E(G)$. The cardinalities of these sets are denoted by $\nu(G)$ and $\mathcal{E}(G)$, respectively. The cycle on n vertices is denoted by C_n .

Let G_1 and G_2 be graphs. The union of G_1 and G_2 is a graph with vertex set $V(G_1) \cup V(G_2)$ and edge set $E(G_1) \cup E(G_2)$. Two graphs G_1 and G_2 are vertex disjoint if and only if $V(G_1) \cap V(G_2) = \emptyset$; G_1 and G_2 are edge disjoint if $E(G_1) \cap E(G_2) = \emptyset$. If G_1 and G_2 are vertex disjoint, we denote their union by $G_1 + G_2$. The intersection $G_1 \cap G_2$ of graphs G_1 and G_2 is defined similarly, but in this case we need to assume that $V(G_1) \cap V(G_2) \neq \emptyset$. The join $G \vee H$ of two vertex disjoint graphs G and H is the graph obtained from $G + H$ by joining each vertex of G to each vertex of H . For two vertex disjoint subgraphs H_1 and H_2 of G , we let $E_G(H_1, H_2) = \{xy \in E(G) : x \in V(H_1), y \in V(H_2)\}$ and $\mathcal{E}_G(H_1, H_2) = |E_G(H_1, H_2)|$.

In this paper we consider the Turán-type extremal problem with the odd edge disjoint cycles being the forbidden subgraph. Since a bipartite graph contains no odd cycles, the non-bipartite graphs have been considered by some authors. First, we recall some notations and terminologies. For a positive integer n and a set of graphs \mathcal{F} , let $\mathcal{G}(n; \mathcal{F})$ denote the class of non-bipartite \mathcal{F} -free graphs on n vertices, and

$$f(n; \mathcal{F}) = \max\{\mathcal{E}(G) : G \in \mathcal{G}(n; \mathcal{F})\}.$$

For simplicity, in the case when \mathcal{F} consists only of one member C_s , where s is an odd integer, we write $\mathcal{G}(n; s) = \mathcal{G}(n; \mathcal{F})$ and $f(n; s) = f(n; \mathcal{F})$.

An important problem in extremal graph theory is that of determining the values of the function $f(n; \mathcal{F})$. Further, characterize the extremal graphs $\mathcal{G}(n; \mathcal{F})$ where $f(n; \mathcal{F})$ is attained. For a given r , the edge maximal graphs of $\mathcal{G}(n; r)$ have been studied by a number of authors [1, 2, 3, 7, 8, 9, 10, 12]. In 1998, Jia [11] proved the following result:




Figure 1. (a) The figure represents a member of $\mathcal{G}^*(n)$.
 (b) The figure represents a member of $\Omega(n, 2)$.

Theorem 1 (Jia). *Let $G \in \mathcal{G}(n; 5)$, $n \geq 10$. Then $\mathcal{E}(G) \leq \lfloor (n-2)^2/4 \rfloor + 3$. Furthermore, equality holds if and only if $G \in \mathcal{G}^*(n)$ where $\mathcal{G}^*(n)$ is the class of graphs obtained by adding a triangle, two vertices of which are new, to the complete bipartite graph $K_{\lfloor (n-2)/2 \rfloor, \lceil (n-2)/2 \rceil}$. Figure 1(a) displays a member of $\mathcal{G}^*(n)$.*

Jia, also conjectured that $f(n; 2k + 1) \leq \lfloor (n - 2)^2/4 \rfloor + 3$ for all $n \geq 4k + 2$. In 2007, Bataineh, confirmed positively the conjecture. In fact, he proved the following result:

Theorem 2 (Bataineh). *Let $k \geq 3$ be a positive integer and $G \in \mathcal{G}(n; 2k + 1)$. Then for large n , $\mathcal{E}(G) \leq \lfloor (n - 2)^2/4 \rfloor + 3$.*

Furthermore, equality holds if and only if $G \in \mathcal{G}^(n)$ where $\mathcal{G}^*(n)$ is as above.*

Let $\mathcal{G}(n; r, s)$ denote to the class of graphs on n vertices containing no r of s -edge disjoint cycles and

$$f(n; r, s) = \max\{\mathcal{E}(G) : G \in \mathcal{G}(n; r, s)\}.$$

Note that

$$\mathcal{G}(n; 2, s) \subseteq \mathcal{G}(n; 3, s) \subseteq \cdots \subseteq \mathcal{G}(n; r, s).$$

Let $\Omega(n, r)$ denote to the class of graphs obtained by adding $r - 1$ edges to the complete bipartite graphs $K_{\lfloor \frac{n}{2} \rfloor, \lceil \frac{n}{2} \rceil}$. Figure 1(b) displays a member of $\Omega(n, 2)$.

The Turán-type extremal problem with two odd edge disjoint cycles being the forbidden subgraph, was studied by Bataineh and Jaradat [2]. In fact, they only established partial results by proving the following:

Theorem 3 (Bataineh and Jaradat). *Let $k = 1, 2$ and $G \in \mathcal{G}(n; 2, 2k + 1)$. Then for large n ,*

$$\mathcal{E}(G) \leq \lfloor n^2/4 \rfloor + 1.$$

Furthermore, equality holds if and only if $G \in \Omega(n, 2)$.

In this paper, we continue the work initiated in [2] by generalizing and extending the above theorem. In fact, we determine $f(n; r, 2k + 1)$ and characterize the edge maximal members in $\mathcal{G}(n; r, 2k + 1)$. Now, we state a number of results, which play an important role in proving our result.

Lemma 4 (Bondy and Murty). *Let G be a graph on n vertices. If $\mathcal{E}(G) > n^2/4$, then G contains a cycle of length r for each $3 \leq r \leq \lfloor (n + 3)/2 \rfloor$.*

Theorem 5 (Brandt). *Let G be a non-bipartite graph with n vertices and more than $\lfloor (n - 1)^2/4 \rfloor + 1$ edges. Then G contains all cycles of length between 3 and the length of the longest cycle.*

In the rest of this paper, $N_G(u)$ stands for the set of neighbors of u in the graph G . Moreover, $G[X]$ denotes to the subgraph induced by X in G .

2. EDGE-MAXIMAL C_{2k+1} -EDGE DISJOINT FREE GRAPHS

In this section, we determine $f(n; r, 2k + 1)$ and characterize the edge maximal members in $\mathcal{G}(n; r, 2k + 1)$. Observe that $\Omega(n, r) \subseteq \mathcal{G}(n; r, 2k + 1)$ and every graph in $\Omega(n, r)$ contains $\lfloor n^2/4 \rfloor + r - 1$ edges. Thus, we have established that

$$(1) \quad f(n; r, 2k + 1) \geq \lfloor n^2/4 \rfloor + r - 1.$$

In the following work, we establish that equality holds. Further we characterize the edge maximal members in $\mathcal{G}(n; r, 2k + 1)$.

Theorem 6. *Let $k \geq 1, r \geq 2$ be two positive integers and $G \in \mathcal{G}(n; r, 2k + 1)$. For large n ,*

$$\mathcal{E}(G) \leq \lfloor n^2/4 \rfloor + r - 1.$$

Furthermore, equality holds if and only if $G \in \Omega(n, r)$.

Proof. We prove the theorem using induction on r .

Step 1. We show the result for $r = 2$. Note that by Theorem 3, it is enough to prove the result for $k \geq 3$. Let $G \in \mathcal{G}(n, 2, 2k + 1)$. If G does not have a cycle of length $2k + 1$, then by Lemma 4, $\mathcal{E}(G) \leq \lfloor n^2/4 \rfloor$. Thus, $\mathcal{E}(G) < \lfloor n^2/4 \rfloor + 1$. So, we need to consider the case when G has cycles of length $2k + 1$. Assume $C = x_1x_2 \dots x_{2k+1}x_1$ be a cycle of length $2k + 1$ in G . Consider $H = G - \{e_1 = x_1x_2, e_2 = x_2x_3, \dots, e_{2k+1} = x_{2k+1}x_1\}$. Observe that H cannot have $2k + 1$ -cycle as otherwise G would have two edge disjoint $2k + 1$ -cycles. We now consider two cases according to H :

Case 1. H is not a bipartite graph. If $k \geq 2$, then by Theorems 1 and 2

$$\mathcal{E}(H) \leq \lfloor (n - 2)^2/4 \rfloor + 3.$$

But, $\mathcal{E}(G) = \mathcal{E}(H) + 2k + 1 \leq \lfloor \frac{(n-2)^2}{4} \rfloor + 2k + 4 \leq \lfloor \frac{n^2}{4} \rfloor - n + 2k + 5$. Thus, for $n \geq 2k + 5$, we have $\mathcal{E}(G) < \lfloor \frac{n^2}{4} \rfloor + 1$. If $k = 1$, then by Theorems 5 $\mathcal{E}(H) \leq \lfloor (n - 1)^2/4 \rfloor + 1$. And so, by using the same argument as in the above, we get that for $n \geq 7$,

$$\mathcal{E}(G) < \left\lfloor \frac{n^2}{4} \right\rfloor + 1.$$

Case 2. H is a bipartite graph. Let X and Y be the partition of $V(H)$. Thus, $\mathcal{E}(H) \leq |X||Y|$. Observe $|X| + |Y| = n$. The maximum of the above is when $|X| = \lfloor \frac{n}{2} \rfloor$ and $|Y| = \lceil \frac{n}{2} \rceil$. Thus, $\mathcal{E}(H) \leq \lfloor \frac{n^2}{4} \rfloor$. Restore the edges of the cycle C . We now consider the following subcases:

(2.1). One of X and Y contains two edges of the cycle, say e_i and e_j belong to X . Let y_1, y_2, \dots, y_{k-1} be a set of vertices in $X - \{x_i, x_{i+1}, x_j, x_{j+1}\}$. We split this subcase into two subcases:

(2.1.1). i and j are not consecutive. Then $|N_Y(x_i) \cap N_Y(x_{i+1}) \cap N_Y(x_j) \cap N_Y(x_{j+1}) \cap N_Y(y_1) \cap N_Y(y_2) \cap \dots \cap N_Y(y_{k-1})| \leq k + 2$, as otherwise G contains two edge disjoint $2k + 1$ -cycles. Thus,

$$\mathcal{E}_G(\{x_i, x_{i+1}, x_j, x_{j+1}, y_1, y_2, \dots, y_{k-1}\}, Y) \leq (k + 2)|Y| + k + 2.$$

So,

$$\begin{aligned} \mathcal{E}(G) &= \mathcal{E}_G(X - \{x_i, x_{i+1}, x_j, x_{j+1}, y_1, y_2, \dots, y_{k-1}\}, Y) \\ &\quad + \mathcal{E}_G(\{x_i, x_{i+1}, x_j, x_{j+1}, y_1, y_2, \dots, y_{k-1}\}, Y) + \mathcal{E}(G[X]) + \mathcal{E}(G[Y]) \\ &\leq (|X| - k - 3)|Y| + (k + 2)|Y| + k + 2 + 2k + 1 \\ &\leq |X||Y| - |Y| + 3k + 3 \leq (|X| - 1)|Y| + 3k + 3. \end{aligned}$$

Observe that $|X| + |Y| = n$. The maximum of the above equation is when $|Y| = \lceil \frac{n-1}{2} \rceil$ and $|X| - 1 = \lfloor \frac{n-1}{2} \rfloor$. Thus,

$$\mathcal{E}(G) \leq \left\lfloor \frac{(n-1)^2}{4} \right\rfloor + 3k + 3.$$

Hence, for $n \geq 6k + 7$, we have $\mathcal{E}(G) < \lfloor \frac{n^2}{4} \rfloor + 1$.

(2.1.2). i and j are consecutive, say $j = i + 1$. Then by following, word by word, the same arguments as in (2.1.1) and by taking into the account that $|N_Y(x_i) \cap N_Y(x_{i+1}) \cap N_Y(x_{i+2}) \cap N_Y(y_1) \cap N_Y(y_2) \cap \dots \cap N_Y(y_{k-1})| \leq k + 1$ and so $\mathcal{E}(\{x_i, x_{i+1}, x_{i+2}, y_1, y_2, \dots, y_{k-1}\}, Y) \leq (k + 1)|Y| + k + 1$, we get the same inequality.

(2.2). $\mathcal{E}(G[X]) = 1$ and $\mathcal{E}(G[Y]) = 0$ or $\mathcal{E}(G[X]) = 0$ and $\mathcal{E}(G[Y]) = 1$, say $e_1 \in E(G[X])$. Then $G - e_1$ is a bipartite graph and so as in the above $\mathcal{E}(G - e_1) \leq \lfloor \frac{n^2}{4} \rfloor$. Thus, $\mathcal{E}(G) = \mathcal{E}(G - e_1) + 1 \leq \lfloor \frac{n^2}{4} \rfloor + 1$.

One can observe from the above arguments that for $r = 2$ the only time we have equality is in case when G is obtained by adding an edge to the complete bipartite graph $K_{\lfloor \frac{n}{2} \rfloor, \lceil \frac{n}{2} \rceil}$. This gives rise to the class $\Omega(n, 2)$.

Step 2. Assume that the result is true for $r - 1$. We now show the result is true for $r \geq 3$. To accomplish that we use similar arguments to those in Step 1. Let $G \in \mathcal{G}(n; r, 2k + 1)$. If G contains no $r - 1$ edge disjoint cycles of length $2k + 1$, then by the inductive step $\mathcal{E}(G) \leq \lfloor n^2/4 \rfloor + r - 2$. Thus, $\mathcal{E}(G) < \lfloor n^2/4 \rfloor + r - 1$. So, we need to consider the case when G has $r - 1$ edge disjoint cycles of length $2k + 1$. Assume that $\{C^i = x_{i1}, x_{i2}, \dots, x_{i2k+1}, x_{i1}\}_{i=1}^{r-1}$ be the set of cycles of length $2k + 1$. Consider $H = G - \cup_{i=1}^{r-1} E(C^i)$. Observe that H cannot have $2k + 1$ -cycles as otherwise G would have r of edges disjoint $2k + 1$ -cycles. As in Step 1, we consider two cases:

Case I. H is not a bipartite graph. If $k \geq 2$, then by Theorems 1 and 2 $\mathcal{E}(H) \leq \lfloor (n-2)^2/4 \rfloor + 3$. Thus, $\mathcal{E}(G) = \mathcal{E}(H) + (r-1)(2k+1) \leq \lfloor \frac{n^2}{4} \rfloor + (r-1) - n + 4 + 2k(r-1)$. Hence, for $n > 4 + 2k(r-1)$, we have $\mathcal{E}(G) < \lfloor \frac{n^2}{4} \rfloor + r - 1$. If $k = 1$, then by Theorems 5 $\mathcal{E}(H) \leq \lfloor (n-1)^2/4 \rfloor + 1$.

By using the same argument as in the above, we get that for $n \geq 4(r-1) + 1$,

$$\mathcal{E}(G) < \left\lfloor \frac{n^2}{4} \right\rfloor + 1.$$

Case II. H is a bipartite graph. Let X and Y be the partition of $V(H)$. Thus, $\mathcal{E}(H) \leq |X||Y|$. Observe $|X| + |Y| = n$. The maximum of the above is when $|X| = \lfloor \frac{n}{2} \rfloor$ and $|Y| = \lceil \frac{n}{2} \rceil$. Thus, $\mathcal{E}(H) \leq \lfloor \frac{n^2}{4} \rfloor$. Now, we consider the following two subcases:

(II.I). There is $1 \leq m \leq r-1$ such that C^m contains at least two edges, say $e_i = x_{mi}x_{m(i+1)}$ and $e_j = x_{mj}x_{m(j+1)}$, joining vertices of one of X and Y , say X . Let y_1, y_2, \dots, y_{k-1} be a set of vertices in $X - \{x_{mi}, x_{m(i+1)}, x_{mj}, x_{m(j+1)}\}$. To this end we have two subcases:

(II.I.I). i and j are not consecutive. Then $|N_Y(x_{mi}) \cap N_Y(x_{m(i+1)}) \cap N_Y(x_{mj}) \cap N_Y(x_{m(j+1)}) \cap N_Y(y_1) \cap N_Y(y_2) \cap \dots \cap N_Y(y_{k-1})| \leq k+2$, as otherwise $H \cup \{e_i, e_j\}$ contains two edges disjoint $2k+1$ -cycles and so G contains r edge disjoint cycles of length $2k+1$. Thus, as in (2.1.1) of Step 1,

$$\mathcal{E}_H(\{x_{mi}, x_{m(i+1)}, x_{mj}, x_{m(j+1)}, y_1, y_2, \dots, y_{k-1}\}, Y) \leq (k+2)|Y| + k + 2.$$

And so,

$$\begin{aligned} \mathcal{E}(G) &= \mathcal{E}(H) + |\cup_{i=1}^{r-1} E(C^i)| \\ &= \mathcal{E}_H(X - \{x_{mi}, x_{m(i+1)}, x_{mj}, x_{m(j+1)}, y_1, y_2, \dots, y_{k-1}\}, Y) \\ &\quad + \mathcal{E}_H(\{x_{mi}, x_{m(i+1)}, x_{mj}, x_{m(j+1)}, y_1, y_2, \dots, y_{k-1}\}, Y) + |\cup_{i=1}^{r-1} E(C^i)| \\ &\leq (|X| - k - 3)|Y| + (k+2)|Y| + k + 2 + (r-1)(2k+1) \\ &= (|X| - 1)|Y| + k + 2 + (r-1)(2k+1). \end{aligned}$$

Moreover, the maximum of the above inequality is when $|Y| = \lceil \frac{n-1}{2} \rceil$ and $|X| - 1 = \lfloor \frac{n-1}{2} \rfloor$. Thus,

$$\mathcal{E}(G) \leq \left\lfloor \frac{(n-1)^2}{4} \right\rfloor + k + 2 + (r-1)(2k+1).$$

Hence, for $n \geq 6k(r-1) + 7$, we have $\mathcal{E}(G) < \lfloor \frac{n^2}{4} \rfloor + (r-1)$.

(II.I.II). i and j are consecutive, say $j = i + 1$. Then by following word by word the same arguments as in (2.1.2) of Step 1 and (II.I.I) of Step 2, we get the same inequality

$$\mathcal{E}(G) < \left\lfloor \frac{n^2}{4} \right\rfloor + (r - 1).$$

(II.II). Each $1 \leq m \leq r - 1$, C^m has exactly one edge belonging to one of X and Y . Let e be the edge of C^1 that belongs to one of X and Y . Then $G - e \in \mathcal{G}(n; r - 1, 2k + 1)$ and so by inductive step, $\mathcal{E}(G) = \mathcal{E}(G - e) + 1 \leq \left\lfloor \frac{n^2}{4} \right\rfloor + r - 2 + 1 = \left\lfloor \frac{n^2}{4} \right\rfloor + r - 1$.

We now characterize the extremal graphs. Throughout the proof, we notice that the only time we have equality is in case when G obtained by adding $r - 1$ edges to the complete bipartite graph $K_{\lfloor \frac{n}{2} \rfloor, \lceil \frac{n}{2} \rceil}$. This gives rise to the class $\Omega(n, r)$. This completes the proof of the theorem. ■

From Theorem 6 and the inequality (1), we get that for $k \geq 1$, $r \geq 2$ and large n , $f(n; r, 2k + 1) = \left\lfloor \frac{n^2}{4} \right\rfloor + r - 1$.

REFERENCES

- [1] M.S. Bataineh, Some Extremal Problems in Graph Theory, Ph.D Thesis, Curtin University of Technology (Australia, 2007).
- [2] M.S. Bataineh and M.M.M. Jaradat, *Edge maximal C_3 and C_5 -edge disjoint free graphs*, International J. Math. Combin. **1** (2011) 82–87.
- [3] J. Bondy, *Large cycle in graphs*, Discrete Math. **1** (1971) 121–132.
doi:10.1016/0012-365X(71)90019-7
- [4] J. Bondy, *Pancyclic graphs*, J. Combin. Theory (B) **11** (1971) 80–84.
doi:10.1016/0095-8956(71)90016-5
- [5] J. Bondy and U. Murty, Graph Theory with Applications (The MacMillan Press, London, 1976).
- [6] S. Brandt, *A sufficient condition for all short cycles*, Discrete Appl. Math. **79** (1997) 63–66.
doi:10.1016/S0166-218X(97)00032-2
- [7] L. Caccetta, *A problem in extremal graph theory*, Ars Combin. **2** (1976) 33–56.
- [8] L. Caccetta and R. Jia, Edge maximal non-bipartite Hamiltonian graphs without cycles of length 5, Technical Report.14/97. School of Mathematics and Statistics, Curtin University of Technology (Australia, 1997).
- [9] L. Caccetta and R. Jia, *Edge maximal non-bipartite graphs without odd cycles of prescribed length*, Graphs and Combin. **18** (2002) 75–92.
doi:10.1007/s003730200004

- [10] Z. Füredi, *On the number of edges of quadrilateral-free graphs*, J. Combin. Theory (B) **68** (1996) 1–6.
doi:10.1006/jctb.1996.0052
- [11] R. Jia, *Some Extremal Problems in Graph Theory*, Ph.D Thesis, Curtin University of Technology (Australia, 1998).
- [12] P. Turán, *On a problem in graph theory*, Mat. Fiz. Lapok **48** (1941) 436–452.

Received 27 August 2010

Revised 15 March 2011

Accepted 12 May 2011