

ALMOST SELF-COMPLEMENTARY 3-UNIFORM HYPERGRAPHS

LATA N. KAMBLE

Department of Mathematics
Abasaheb Garware College
Karve Road, Pune-411004

e-mail: lata7429@gmail.com

CHARUSHEELA M. DESHPANDE

AND

BHAGYASHREE Y. BAM

Department of Mathematics
College of Engineering Pune
Pune-411006

e-mail: dcm.maths@coep.ac.in
bpa.maths@coep.ac.in

Abstract

It is known that self-complementary 3-uniform hypergraphs on n vertices exist if and only if n is congruent to 0, 1 or 2 modulo 4. In this paper we define an almost self-complementary 3-uniform hypergraph on n vertices and prove that it exists if and only if n is congruent to 3 modulo 4. The structure of corresponding complementing permutation is also analyzed. Further, we prove that there does not exist a regular almost self-complementary 3-uniform hypergraph on n vertices where n is congruent to 3 modulo 4, and it is proved that there exist a quasi regular almost self-complementary 3-uniform hypergraph on n vertices where n is congruent to 3 modulo 4.

Keywords: uniform hypergraph, self-complementary hypergraph, almost complete 3-uniform hypergraph, almost self-complementary hypergraph, quasi regular hypergraph.

2010 Mathematics Subject Classification: 05C65.

1. INTRODUCTION

The study of self-complementary graphs was initiated by Sachs and Ringel, independently. Ringel [11] and Sachs [12] have both proved the results concerning order and cycle structure of a complementing permutation of self-complementary graphs. Das [3] introduced the concept of almost self-complementary graphs which is similar to graphs self-complementary in $K_n - e$ introduced by Clapham [1]. They proved similar results on order and cycle structure of complementing permutation of almost self-complementary graphs. Kocay [9] extended the results of self-complementary graphs to self-complementary 3-uniform hypergraphs. He has analysed the cycle structure of complementing permutation of self-complementary 3-uniform hypergraphs. Szymański and Wojda [13] have characterized n and k for which there exist k -uniform self-complementary hypergraphs and gave the structure of corresponding self-complementing permutations. Gosselin [5] has characterized all n and k for which there exists a regular k -uniform self-complementary hypergraph of order n .

Potočník and Šajana [10] raised the following question strengthening Hartman's conjecture [2, 6] about existence of large sets of (not necessarily isomorphic) designs.

Question [10]. *Is it true that for every triple of integers $t < k < n$ such that $\binom{n-i}{k-i}$ is even for all $i = 0, \dots, t$, there exists a self-complementary t -subset-regular k -uniform hypergraph of order n ?*

The answer to the above question is affirmative for $k = 2$ and $t = 1$ (see [12]). The answer was proved affirmative also for the case $k = 3$ and $t = 1$ (see [10]). And in [8] it is shown that the answer to the question above is affirmative for the remaining case of 3-uniform hypergraph, namely for the case $k = 3$, $t = 2$.

It is clear that if the number of triples in the complete design (K_n^3) is odd, then there does not exist a self-complementary t -subset-regular 3-uniform hypergraph of order n . In this case one may modify the problem by "Does there exist a partition of $K_n^3 - e$ into two isomorphic t -subset-regular 3-uniform hypergraphs of order n ?" Das and Rosa [4] proved that there exists a partition of Steiner triple system (STS) into two isomorphic 3-uniform hypergraphs of order n , if $n \equiv 3$ or $7 \pmod{12}$. In this paper we prove that there does not exist a partition of $K_n^3 - e$ into two isomorphic 1-subset-regular 3-uniform hypergraphs of order n , if $n \equiv 3 \pmod{4}$ and some partial answers to the above question are given.

In Section 2, we define almost self-complementary 3-uniform hypergraph on n vertices and further prove that such a hypergraph exists if and only if n is congruent to 3 modulo 4.

In Section 3, the structure of a complementing permutation of such an almost self-complementary 3-uniform hypergraph is analyzed.

In Section 4, we prove that there does not exist a regular almost self complementary 3-uniform hypergraph on n vertices where n is congruent to 3 modulo 4. Further, we prove that there exists a quasi regular almost self-complementary 3-uniform hypergraph on n vertices where n is congruent to 3 modulo 4.

2. NECESSARY AND SUFFICIENT CONDITION FOR EXISTENCE OF ALMOST SELF-COMPLEMENTARY 3-UNIFORM HYPERGRAPH

Suppose H is a 3-uniform hypergraph with vertex set V and edge set E . A partition of $E = \bigcup_{i=1}^s E_i$ is called a factorization of H and the 3-uniform hypergraph $H_i(V, E_i)$ is called a factor of H for $i = 1, 2, \dots, s$. A factorization in which all factors are isomorphic is called an isomorphic factorization.

A factor in a factorization of complete 3-uniform hypergraph K_n^3 with only two isomorphic factors is nothing but a self complementary 3-uniform hypergraph. A partitioning of the edge set of K_n^3 into two isomorphic factors is not possible when K_n^3 has an odd number of edges, i.e., when n is congruent to 3 modulo 4. However, after deleting some odd number of edges from K_n^3 the remaining 3-uniform hypergraph may be partitioned into two isomorphic factors.

In this paper we delete one edge from K_n^3 and define an almost self-complementary 3-uniform hypergraph. We always denote by e the edge deleted from K_n^3 , call it the missing edge and the corresponding vertices of e the special vertices.

Definition. The hypergraph $\tilde{K}_n^3 = K_n^3 - e$ is called an almost complete 3-uniform hypergraph.

Definition. A 3-uniform hypergraph H with n vertices is almost self-complementary if it is isomorphic with its complement \bar{H} with respect to \tilde{K}_n^3 .

This means that a 3-uniform hypergraph H with n vertices is almost self-complementary if \tilde{K}_n^3 can be decomposed into two isomorphic factors with H as one factor.

Since \tilde{K}_n^3 has $\binom{n}{3} - 1$ edges, such a factorization is possible only if this number is divisible by 2. Thus it is necessary that $n \equiv 3 \pmod{4}$. We can compare this with the fact that isomorphic factorizations of K_n^3 , into 2 factors, i.e., self-complementary 3-uniform hypergraphs, exist only if $n \equiv 0, 1$ or $2 \pmod{4}$. Almost self-complementary 3-uniform hypergraphs in a sense fill the gap where self-complementary 3-uniform hypergraphs do not exist.

Following theorem gives a necessary and sufficient condition on the order of an almost self-complementary 3-uniform hypergraph.

Theorem 1. *There exists an almost self-complementary 3-uniform hypergraph on n vertices if and only if $n \equiv 3 \pmod{4}$.*




Figure 1. The types of triples making up the edge set of an almost self-complementary 3-uniform hypergraph on $n = 4m + 3$ vertices.

Proof. Necessity is obvious from the above discussions. To prove sufficiency, we construct a 3-uniform hypergraph which is self-complementary in \tilde{K}_n^3 on n vertices with $n \equiv 3 \pmod{4}$. Denote the missing edge by $e = \{x, y, z\}$.

Let m be a positive integer such that $n = 4m + 3$ and $V = V_0 \cup V_1 \cup V_2 \cup V_3 \cup \{x, y, z\}$, where $V_i = \{v_j^i : j \in \mathbb{Z}_m\}$ for all $i \in \mathbb{Z}_4$.

For pairwise distinct $i, i', i'' \in \mathbb{Z}_4$ we consider the following partition of edges of \tilde{K}_n^3 .

$$\begin{aligned}
 E_i &= V_i^{(3)} = \text{all 3-subsets of } V_i, \\
 E_{i,i'} &= \{\{v_{j_1}^i, v_{j_2}^i, v_{j'}^{i'}\} : j_1, j_2, j' \in \mathbb{Z}_m, j_1 \neq j_2\}, \\
 E_{i,i',i''} &= \{\{v_j^i, v_{j'}^{i'}, v_{j''}^{i''}\} : j, j', j'' \in \mathbb{Z}_m\}, \\
 E_i^x &= \{\{x, v_{j_1}^i, v_{j_2}^i\} : j_1, j_2 \in \mathbb{Z}_m, j_1 \neq j_2\}, \\
 E_i^y &= \{\{y, v_{j_1}^i, v_{j_2}^i\} : j_1, j_2 \in \mathbb{Z}_m, j_1 \neq j_2\}, \\
 E_i^z &= \{\{z, v_{j_1}^i, v_{j_2}^i\} : j_1, j_2 \in \mathbb{Z}_m, j_1 \neq j_2\}, \\
 E_{i,i'}^x &= \{\{x, v_j^i, v_{j'}^{i'}\} : j, j' \in \mathbb{Z}_m\}, \\
 E_{i,i'}^y &= \{\{y, v_j^i, v_{j'}^{i'}\} : j, j' \in \mathbb{Z}_m\}, \\
 E_{i,i'}^z &= \{\{z, v_j^i, v_{j'}^{i'}\} : j, j' \in \mathbb{Z}_m\},
 \end{aligned}$$

$$\begin{aligned} E_i^{(x,y)} &= \{\{x, y, v_j^i\} : j \in \mathbb{Z}_m\}, \\ E_i^{(x,z)} &= \{\{x, z, v_j^i\} : j \in \mathbb{Z}_m\}, \\ E_i^{(y,z)} &= \{\{y, z, v_j^i\} : j \in \mathbb{Z}_m\}. \end{aligned}$$

Let

$$\begin{aligned} E = \bigcup_{k=0,1} & \left(E_k \cup E_{2,k} \cup E_{3,k} \cup E_{k,2,3} \cup E_k^x \cup E_k^y \cup E_k^z \cup E_k^{(x,y)} \cup E_k^{(x,z)} \cup E_k^{(y,z)} \right) \\ & \cup E_{0,1} \cup E_{1,0} \cup E_{0,1}^x \cup E_{1,2}^x \cup E_{0,3}^x \cup \bigcup_{k=1,2,3} \left(E_{k,k+1}^y \cup E_{k,k+1}^z \right). \end{aligned}$$

Let H be the 3-uniform hypergraph with vertex set V and edge set E . Figure 1 gives a diagrammatic construction of H .

To prove that H is almost self-complementary, we define a permutation $\phi : V \rightarrow V$ by $\phi(x) = x, \phi(y) = y, \phi(z) = z, \phi(v_j^0) = v_j^3, \phi(v_j^3) = v_j^1, \phi(v_j^1) = v_j^2,$ and $\phi(v_j^2) = v_j^0,$ for all $j \in \mathbb{Z}_m$. It is checked easily that ϕ is a complementing permutation of H and therefore H is almost self-complementary. ■

3. THE COMPLEMENTING PERMUTATION

It is known (see [9]) that for a 3-uniform self-complementary (s.c.) hypergraph, if τ is a complementing permutation of the vertices that maps H onto its complement \bar{H} , then

- (i) every cycle of τ has even length, or
- (ii) τ has 1 or 2 fixed points, and the length of all other cycles is a multiple of 4.

We prove similar results for the complementing permutation of an almost self-complementary 3-uniform hypergraph.

Given an almost self-complementary 3-uniform hypergraph H , let the edges of H be coloured red and the remaining edges of \tilde{K}_n^3 be coloured green. Since the 2 factors are isomorphic, there is a permutation τ of the vertices of \tilde{K}_n^3 that induces a mapping of the red edges onto the green edges. We consider τ as a permutation of the vertices of K_n^3 , and denote by τ' the corresponding mapping induced on the set of edges of K_n^3 . Thus τ' maps each red edge onto a green edge. However, the mapping τ' need not necessarily map each green edge onto a red edge. This would be so if τ' mapped e onto itself, but it may be that τ' maps e onto a red edge and some green edge onto e . Such a τ (which, for definiteness we shall always assume induces a mapping from red to green) will (as for s.c. 3-uniform hypergraphs) be called a complementing permutation. It will be useful to consider the cycles of the induced mapping τ' .

The following remarks regarding the cycles of induced mapping τ' will be used to prove a number of results about the structure of complementing permutation τ .

Remark 2. A cycle of τ' that does not include e must be of even length, consisting of edges alternately red and green.

Remark 3. The cycle of τ' that includes e has odd length, consisting of e followed by red and green edges alternately. Further, this length equals 1 when τ' maps e onto itself.

Lemma 4. *If the special vertices x, y, z occur in different cycles of τ , then they must be three fixed points $(x)(y)(z)$.*

Proof. Suppose that x occurs in a cycle of length L_1 , y occurs in a cycle of length L_2 and z occurs in a cycle of length L_3 .

Consider the cycle of τ' including e . The number of edges in this cycle is the least common multiple of L_1, L_2 and L_3 . By Remark 3 this number must be odd. If $L_1 \geq 3$ then any triple $\{i, j, k\}$ of this cycle gives rise to a sequence of triples $\{i, j, k\}, \tau\{i, j, k\}, \tau^2\{i, j, k\}, \dots$ etc. These must be alternately edges of H and \bar{H} . This is possible only if L_1 is even. Hence $L_1 = 1$. Similarly $L_2 = L_3 = 1$. ■

Lemma 5. *If all special vertices x, y, z occur in the same cycle C of τ , then C has length 3.*

Proof. Suppose all the special vertices x, y, z occur in a cycle C of length L . Consideration of the cycle of τ' including e shows that L must be odd.

If $L > 3$, one finds that there is another cycle of τ' not including e , of odd length, which is a contradiction to Remark 2. Thus $L = 3$. ■

Lemma 6. *If any two of the special vertices, say x, y , occur in the same cycle C_1 of τ , then C_1 has length $4h + 2, h \geq 0$, with $\tau^{2h+1}(x) = y$ and special vertex z fixed.*

Proof. Suppose special vertices x, y occur in the same cycle C_1 of length L_1 . Let the remaining special vertex z occur in a cycle C_2 of length L_2 . Clearly $L_1 \geq 2$ and $L_2 \geq 1$. Since $L_1 \geq 2$, it must be even as argued in Lemma 4. If $L_2 > 1$ we get a contradiction to Remark 3. Hence $L_2 = 1$.

Let $\tau^m(x) = y$, therefore $\tau^{L_1-m}(y) = x$. Consider the sequence of triples $\{x, y, z\}, \tau\{x, y, z\}, \dots$. This cycle has length either L_1 or m if $m = \frac{L_1}{2}$ and it must be odd. Since L_1 is even the only possibility is that the length is m and m is odd. Hence $L_1 = 2m = 4h + 2, h \geq 0$. ■

Lemma 7. *The cycles of τ that do not include the special vertices are of length multiple of 4.*

Proof. If a cycle $(u_1u_2 \cdots u_L)$ does not involve the special vertices, then by Remark 2 L is even. We get the following cases depending on occurrence of special vertices in τ .

Case (i) Special vertices x, y, z are fixed points.

Case (ii) Special vertices x, y, z occur in the cycle of length 3, say $C' = (x, y, z)$.

Case (iii) Two special vertices say x, y occur in the same cycle and z is fixed.

In all these cases consider the cycle of τ' including the edge $\{u_1, u_{(\frac{L}{2}+1)}, z\}$ which is of length $\frac{L}{2}$. From Remark 2 we find that $\frac{L}{2}$ must be even. Thus L must be a multiple of 4. ■

Complete description of complementing permutation of almost self complementary 3-uniform hypergraph is given below.

Theorem 8. *Let τ be a complementing permutation of an almost self-complementary 3-uniform hypergraph on $n \geq 3$ vertices and $e = \{x, y, z\}$ be the deleted edge. Then $n \equiv 3 \pmod{4}$. Further*

- (a) τ consists of 3 fixed special vertices and all other cycles of length multiple of 4, or
- (b) τ consists of a cycle of length $4h + 2$, $h \geq 0$, including two special vertices x and y with $\tau^{2h+1}(x) = y$, one fixed special vertex z and all other cycles of length multiple of 4, or
- (c) τ consists of the cycle (x, y, z) and all other cycles are of length multiple of 4.

4. REGULAR AND QUASI REGULAR ALMOST SELF-COMPLEMENTARY 3-UNIFORM HYPERGRAPH

It is known that (see [10]) a regular self-complementary 3-uniform hypergraph on n vertices exists if and only if $n \geq 5$ and n is congruent to 1 or 2 modulo 4. In the next theorem we prove that there does not exist a regular almost self-complementary 3-uniform hypergraph on n vertices where n is congruent to 3 modulo 4.

Theorem 9. *There does not exist a regular almost self-complementary 3-uniform hypergraph on n vertices where n is congruent to 3 modulo 4.*

Proof. Suppose there exists a regular almost self-complementary 3-uniform hypergraph, say H of regular degree r . Then the total number of edges in H is $\frac{1}{2}(\binom{n}{3} - 1) = \frac{n(n-1)(n-2)-6}{12}$. Since H is regular we get $rn = 3 \times$ number of edges

in H , i.e., $rn = \frac{3(n(n-1)(n-2)-6)}{12}$. Hence $r = \frac{n(n-1)(n-2)-6}{4n}$ which is not an integer for any n congruent to 3 modulo 4, a contradiction. Hence, there does not exist a regular almost self-complementary 3-uniform hypergraph on n vertices where n is congruent to 3 modulo 4. ■

A hypergraph H is said to be quasi regular if the degree of every vertex is either r or $r - 1$ for some positive integer r . In [7], it is proved that there exists a quasi regular self-complementary 3-uniform hypergraph on n vertices if and only if n is congruent to 0 modulo 4. In the following theorem we prove that there exists a quasi regular almost self-complementary 3-uniform hypergraph on n vertices where n is congruent to 3 modulo 4.

Theorem 10. *There exists a quasi regular almost self complementary 3-uniform hypergraph on n vertices where n is congruent to 3 modulo 4.*

Proof. H constructed in Theorem 1 is already shown to be almost self-complementary 3-uniform hypergraph. We show that H is quasi regular. Considering the same notation as in proof of Theorem 1, take any vertex v_j^i .

Case (i) If $i \in \{0, 1\}$ then, for fixed $i', i'' \in \mathbb{Z}_4$ distinct from i , the vertex v_j^i lies in $\binom{m-1}{2}$ triples of E_i , $3\binom{m}{2}$ triples of $E_{i',i}$, $(m-1)m$ triples of $E_{i,i'}$, m^2 triples of $E_{i,i',i''}$, $(m-1)$ triples of each E_i^x, E_i^y, E_i^z , $4m$ triples of $E_{i,i'}^x, E_{i,i'}^y, E_{i,i'}^z$ and 1 triple of each $E_i^{(x,y)}, E_i^{(x,z)}, E_i^{(y,z)}$. Hence, for every vertex v_j^i in H with $i \in \{0, 1\}$, we have

$$\deg(v_j^i) = \binom{m-1}{2} + 3\binom{m}{2} + m(m-1) + m^2 + 3(m-1) + 4m + 3 = 4m^2 + 3m + 1.$$

Case (ii) If $i \in \{2, 3\}$ then the vertex v_j^i lies in $2(m-1)m$ triples of $E_{i,i'}$, $2m^2$ triples of $E_{i,i',i''}$, $5m$ triples of $E_{i,i'}^x, E_{i,i'}^y, E_{i,i'}^z$. Hence, for every vertex v_j^i in H with $i \in \{2, 3\}$, we obtain

$$\deg(v_j^i) = 2(m-1)m + 2m^2 + 5m = 4m^2 + 3m.$$

Case (iii) x lies in $2\binom{m}{2}$ triples of E_i^x , m triples of each $E_i^{(x,y)}, E_i^{(x,z)}$ and $3m^2$ triples of $E_{i,i'}^x$. Hence

$$\deg(x) = 2\binom{m}{2} + 4m + 3m^2 = 4m^2 + 3m.$$

y lies in $2\binom{m}{2}$ triples of E_i^y , $4m$ triples of $E_i^{(x,y)}, E_i^{(y,z)}$ and $3m^2$ triples of $E_{i,i'}^y$. Hence

$$\deg(y) = 2\binom{m}{2} + 4m + 3m^2 = 4m^2 + 3m.$$

Similarly,

$$\deg(z) = 2 \binom{m}{2} + 4m + 3m^2 = 4m^2 + 3m.$$

Hence, H is quasi regular with degrees $r = 4m^2 + 3m + 1$ and $r - 1 = 4m^2 + 3m$. ■

Acknowledgment

Authors wish to thank Professor N.S. Bhave for fruitful discussions.

REFERENCES

- [1] C.R.J. Clapham, *Graphs self-complementary in K_n -e*, Discrete Math. **81** (1990) 229–235.
doi:10.1016/0012-365X(90)90062-M
- [2] C.J. Colbourn and J.H. Dinitz, *The CRC Handbook of Combinatorial Designs* (CRC Press, Boca Raton, 1996).
- [3] P.K. Das, *Almost self-complementary graphs 1*, Ars Combin. **31** (1991) 267–276.
- [4] P.K. Das and A. Rosa, *Halving Steiner triple systems*, Discrete Math. **109** (1992) 59–67.
doi:10.1016/0012-365X(92)90278-N
- [5] S. Gosselin, *Constructing regular self-complementary uniform hypergraphs*, Combin. Designs **16** (2011) 439–454.
doi:10.1002/jcd.20286
- [6] A. Hartman, *Halving the complete design*, Ann. Discrete Math. **34** (1987) 207–224.
doi:10.1016/s0304-0208(08)72888-3
- [7] L.N. Kamble, C.M. Deshpande and B.Y. Bam, *The existence of quasi regular and bi-regular self-complementary 3-uniform hypergraphs*, Discuss. Math. Graph Theory **36** (2016) 419–426.
doi:10.7151/dmgt.1862
- [8] M. Knor and P. Potočník, *A note on 2-subset-regular self-complementary 3-uniform hypergraphs*, Ars Combin. **11** (2013) 33–36.
- [9] W. Kocay, *Reconstructing graphs as subsumed graphs of hypergraphs, and some self-complementary triple systems*, Graphs Combin. **8** (1992) 259–276.
doi:10.1007/BF02349963
- [10] P. Potočník and M. Šajana, *The existence of regular self-complementary 3-uniform hypergraphs*, Discrete Math. **309** (2009) 950–954.
doi:10.1016/j.disc.2008.01.026
- [11] G. Ringel, *Über Selbstkomplementäre Graphen*, Arch. Math. **14** (1963) 354–358.
doi:10.1007/BF01234967

- [12] H. Sachs, *Über Selbstkomplementäre Graphen*, Publ. Math. Debrecen **9** (1962) 270–288.
- [13] A. Szymański and A.P. Wojda, *A note on k -uniform self-complementary hypergraphs of given order*, Discuss. Math. Graph Theory **29** (2009) 199–202.
doi:10.7151/dmgt.1440

Received 30 September 2015

Revised 23 February 2016

Accepted 3 March 2016