

$(k - 1)$ -KERNELS IN STRONG k -TRANSITIVE DIGRAPHS

RUIXIA WANG

School of Mathematical Sciences
Shanxi University
Taiyuan Shanxi, 030006, PR China

e-mail: wangrx@sxu.edu.cn

Abstract

Let $D = (V(D), A(D))$ be a digraph and $k \geq 2$ be an integer. A subset N of $V(D)$ is k -independent if for every pair of vertices $u, v \in N$, we have $d(u, v) \geq k$; it is l -absorbent if for every $u \in V(D) - N$, there exists $v \in N$ such that $d(u, v) \leq l$. A (k, l) -kernel of D is a k -independent and l -absorbent subset of $V(D)$. A k -kernel is a $(k, k - 1)$ -kernel.

A digraph D is k -transitive if for any path $x_0x_1 \cdots x_k$ of length k , x_0 dominates x_k . Hernández-Cruz [3-*transitive digraphs*, Discuss. Math. Graph Theory **32** (2012) 205–219] proved that a 3-transitive digraph has a 2-kernel if and only if it has no terminal strong component isomorphic to a 3-cycle. In this paper, we generalize the result to strong k -transitive digraphs and prove that a strong k -transitive digraph with $k \geq 4$ has a $(k - 1)$ -kernel if and only if it is not isomorphic to a k -cycle.

Keywords: digraph, transitive digraph, k -transitive digraph, k -kernel.

2010 Mathematics Subject Classification: 05C20.

REFERENCES

- [1] J. Bang-Jensen and G. Gutin, *Digraphs: Theory, Algorithms and Applications* (Springer, London, 2000).
- [2] E. Boros and V. Gurvich, *Perfect graphs, kernels, and cores of cooperative games*, Discrete Math. **306** (2006) 2336–2354.
doi:10.1016/j.disc.2005.12.031
- [3] V. Chvátal, *On the computational complexity of finding a kernel*, Report No. CRM-300, Centre de Recherches Mathématiques, Université de Montréal (1973).

- [4] C. Hernández-Cruz and H. Galeana-Sánchez, *k-kernels in k-transitive and k-quasi-transitive digraphs*, Discrete Math. **312** (2012) 2522–2530.
doi:10.1016/j.disc.2012.05.005
- [5] C. Hernández-Cruz, *3-transitive digraphs*, Discuss. Math. Graph Theory **32** (2012) 205–219.
doi:10.7151/dmgt.1613
- [6] C. Hernández-Cruz, *4-transitive digraphs I: the structure of strong transitive digraphs*, Discuss. Math. Graph Theory **33** (2013) 247–260.
doi:10.7151/dmgt.1645
- [7] C. Hernández-Cruz and J.J. Montellano-Ballesteros, *Some remarks on the structure of strong k-transitive digraphs*, Discuss. Math. Graph Theory **34** (2014) 651–671.
doi:10.7151/dmgt.1765
- [8] H. Galeana-Sánchez, C. Hernández-Cruz and M.A. Juárez-Camacho, *On the existence and number of (k+1)-kings in k-quasi-transitive digraphs*, Discrete Math. **313** (2013) 2582–2591.
doi:10.1016/j.disc.2013.08.007
- [9] M. Kwaśnik, *On (k, l)-kernels on graphs and their products*, Doctoral Dissertation, Technical University of Wrocław, Wrocław, 1980.

Received 7 January 2014

Revised 15 May 2014

Accepted 19 May 2014