

THE FLOWER CONJECTURE IN SPECIAL CLASSES OF GRAPHS

ZDENĚK RYJÁČEK¹

*Department of Mathematics, University of West Bohemia
Americká 42, 306 14 Plzeň, Czech Republic*

and

INGO SCHIERMEYER¹

*Lehrstuhl C für Mathematik, Rhein.-Westf. Techn. Hochschule
Templergraben 55, D-52062 Aachen, Germany*

Abstract

We say that a spanning eulerian subgraph $F \subset G$ is a *flower* in a graph G if there is a vertex $u \in V(G)$ (called the center of F) such that all vertices of G except u are of the degree exactly 2 in F . A graph G has the *flower property* if every vertex of G is a center of a flower.

Kaneko conjectured that G has the flower property if and only if G is hamiltonian. In the present paper we prove this conjecture in several special classes of graphs, among others in squares and in a certain subclass of claw-free graphs.

Keywords: hamiltonian graphs, flower conjecture, square, claw-free graphs.

1991 Mathematics Subject Classification: 05C45.

1. INTRODUCTION

We consider only finite undirected graphs without loops and multiple edges. For terminology and notation not defined here we refer to [1].

If $x \in V(G)$, then by $d_G(x)$ we denote the degree of x and by $N_G(x)$ (or simply $N(x)$) we denote the set of all vertices of G that are adjacent

¹This research was carried out partly while the second author was visiting the University of West Bohemia and partly while the first author was visiting RWTH Aachen. Research supported by EC-grant No 927.

to x . Unlike in [1], we denote the induced subgraph on a set $M \subset V(G)$ by $\langle M \rangle$. If for every $x \in V(G)$, $\langle N(x) \rangle$ has a property P , then we say that G is *locally P* .

The *square* of a connected graph H is the graph $G = H^2$ such that $V(G) = V(H)$ and two vertices x, y are adjacent in G if and only if x, y are at distance at most 2 in H . If G and G' are graphs, then we say that G is *G' -free* if G contains no induced subgraph isomorphic to G' . Specifically, in the case that $G' = K_{1,3}$ we say that G is *claw-free* and the star $K_{1,3}$ will be also referred to as the *claw*.

Let G be a graph of order $n \geq 3$ and $u \in V(G)$. If there is a spanning eulerian subgraph F of G such that $d_F(u) \geq 2$ and $d_F(v) = 2$ for all $v \in V(G)$, $v \neq u$, then F is called a *flower at u* and the vertex u is called the *center* of F . If F is a flower at u then the components of the graph $F - u$ will be called the *leaves* of F . Since $1 \leq d_{F-u}(x) \leq 2$ for every $x \neq u$, every leaf of F is a path.

We say that a graph G has the *flower property* if G has a flower at u for every $u \in V(G)$.

Obviously, every hamiltonian cycle of G is a flower and hence every hamiltonian graph has the flower property. Kaneko [4] conjectured that these properties are equivalent.

Conjecture [4] (The Flower Conjecture). *A graph G has the flower property if and only if G is hamiltonian.*

Kaneko and Ota [5] proved that if G has the flower property, then G is 1-tough and has a 2-factor.

In the present paper we prove the flower conjecture in several special classes of graphs.

2. OBSERVATIONS

Proposition 1. *Let G be a graph with a minimum degree $\delta(G) \leq 3$. Then G has the flower property if and only if G is hamiltonian.*

Proof. If $x \in V(G)$ is a vertex such that $d_G(x) \leq 3$ then every flower at x is a hamiltonian cycle. ■

Proposition 2. *Let G be a graph with connectivity $\kappa(G) \leq 2$. Then G has the flower property if and only if G is hamiltonian.*

Proof. If $\kappa(G) = 1$ then G is neither hamiltonian nor has the flower property and thus we can assume that $\kappa(G) = 2$. Suppose that G has

the flower property. Let $\{x, y\}$ be a 2-vertex cut set of G . By the result of Kaneko and Ota [5], G is 1-tough and hence $G - \{x, y\}$ has two components H_1, H_2 . Choose $z_i \in H_i$ and let F_i be a flower of G at z_i , $i = 1, 2$. Then $P_1 = F_1 - H_1$ is a hamiltonian $\{x, y\}$ -path in $G - H_1$ and, similarly, $P_2 = F_2 - H_2$ is a hamiltonian $\{y, x\}$ -path in $G - H_2$. But then the cycle $C = xP_1yP_2x$ is a hamiltonian cycle in G . ■

Proposition 3. *Let G be a bipartite graph. Then G has the flower property if and only if G is hamiltonian.*

Proof. Let (X, Y) be the bipartition of G . If F is a flower at $u \in X$, then $\sum_{x \in X} d_F(x) = |E(F)| = \sum_{y \in Y} d_F(y)$, from which

$$d_F(u) + 2|X - \{u\}| = 2|Y|,$$

or, equivalently,

$$d_F(u) - 2 + 2|X| = 2|Y|,$$

which implies $|X| \leq |Y|$. Taking a flower F' at $v \in Y$, we get analogously $|X| \geq |Y|$ and hence $|X| = |Y|$. This implies $d_F(u) = 2$ and hence F is a hamiltonian cycle. ■

Proposition 4. *Let G be a graph and let $x \in V(G)$ be such that $\langle N(x) \rangle$ is a complete graph. Then G has the flower property if and only if G is hamiltonian.*

Proof. Suppose that G has the flower property and let F be a flower at x such that $d_F(x)$ is minimum. Suppose that $d_F(x) > 2$ and let z_1, z_2 be end vertices of two different leaves of F . Then, deleting from F the edges xz_1, xz_2 and adding z_1z_2 , we get a flower F' with $d_{F'}(x) < d_F(x)$, which contradicts the minimality of F . Thus, $d_F(x) = 2$ and F is a hamiltonian cycle. ■

3. SQUARES

Fleischner [2] proved the following theorem.

Theorem A. [2] *If H is a 2-connected graph and $G = H^2$, then G is hamiltonian.*

The following statement is also due to Fleischner and follows from Theorem 3 of [3].

Theorem B. [3] *Let y be an arbitrary vertex of a 2-connected graph H . Then the graph $G = H^2$ contains a hamiltonian cycle C such that both edges of C containing y are in $E(H)$.*

Using these two theorems, we can prove the following.

Theorem 5. *Let H be a graph and $G = H^2$. Then G has the flower property if and only if G is hamiltonian.*

Proof. Suppose that $G = H^2$ and G has the flower property.

If H is 2-connected, then G is hamiltonian by Theorem A. Hence $\kappa(H) = 1$.

If H has a vertex x with $d_H(x) = 1$, then $\langle N_G(x) \rangle$ is a complete graph and G is hamiltonian by Proposition 4. Hence $\delta(H) \geq 2$.

If H has a cut edge (i.e. an edge which is a block) $xy \in E(H)$, then, since $\delta(H) \geq 2$, $\{x, y\}$ is a 2-vertex cut set of G and G is hamiltonian by Proposition 2.

Hence we can assume that H has connectivity $\kappa(H) = 1$, minimum degree $\delta(H) \geq 2$ and every block of H has at least three vertices.

Let H_1 be an end block (i.e. a block containing exactly one cut vertex) of H and let x be the cut vertex of H in H_1 . By Theorem B, there is a hamiltonian cycle C_1 in H_1^2 such that $xx^- \in E(H)$ and $xx^+ \in E(H)$ (here we denote by x^- and x^+ the predecessor and successor of x on C).

Put $H_2 = H - (H_1 - x)$, choose a vertex $y \in N_{H_1}(x)$ and let F be a flower in G at y . We consider the subgraph $F' = F - (H_1 - x)$. Since $1 \leq d_{F'}(v) \leq 2$ for every $v \in V(H_2)$ and $d_{F'}(v) = 1$ if and only if $v = x$ or $v \in N(x)$, F' is a collection of paths P_i , $i = 1, \dots, \ell$, with end vertices $a_i, b_i \in N(x) \cup \{x\}$, $i = 1, \dots, \ell$.

If all the vertices a_i, b_i , $i = 1, \dots, \ell$, are distinct from x , then, since $\langle N(x) \cup \{x\} \rangle$ is a clique in G , $C' = xa_1P_1b_1a_2P_2b_2 \dots a_\ell P_\ell b_\ell x^+ Cx$ is a hamiltonian cycle in G . Hence there is an i_0 such that $x = a_{i_0}$ (or, similarly, $x = b_{i_0}$). We can assume without loss of generality that $x = a_1$ and then analogously $C' = xP_1b_1a_2P_2b_2 \dots a_\ell P_\ell b_\ell x^+ Cx$ is a hamiltonian cycle in G . ■

4. CLAW-FREE GRAPHS

Theorem 6. *Let G be a graph and let $x \in V(G)$ be such that $\langle N(x) \rangle$ is connected and x is not a vertex of an induced claw in G . Then G has the flower property if and only if G is hamiltonian.*

Proof. Suppose that G has the flower property but is not hamiltonian and let F be a flower at x such that $d_F(x)$ is minimum. Let $P_1, \dots, P_\ell$ be the leaves of F and denote by x_i^1, x_i^2 the end vertices of P_i , $i = 1, \dots, \ell$. If some end vertices $x_{i_1}^{j_1}, x_{i_2}^{j_2}$ ($i_1 \neq i_2$) of two different leaves P_{i_1}, P_{i_2} are adjacent, then, deleting from F the edges $xx_{i_1}^{j_1}, xx_{i_2}^{j_2}$ and adding $x_{i_1}^{j_1}x_{i_2}^{j_2}$, we get a flower F' with $d_{F'}(x) < d_F(x)$. Hence, no end vertices of two different leaves of F can be adjacent. This implies that $\ell = 2$ since otherwise $\langle x, x_1^1, x_2^1, x_3^1 \rangle$ is an induced claw centred at x . Moreover, $x_1^1x_1^2 \in E(G)$ (since otherwise $\langle x, x_1^1, x_1^2, x_2^1 \rangle$ is an induced claw centred at x) and, similarly, $x_2^1x_2^2 \in E(G)$. Denote $x_i^1x_i^2 = e_i$, $i = 1, 2$.

Since $\langle N(x) \rangle$ is connected, there is a path P in $\langle N(x) \rangle$ joining e_1 to e_2 . Suppose that the flower F and the path P are chosen such that, among all flowers F at x with minimum $d_F(x)$, the $\{e_1, e_2\}$ -path P is the shortest possible. We can assume without loss of generality that P is an $\{x_1^1, x_2^1\}$ -path. Let $x_1^1 = z_0, z_1, \dots, z_k = x_2^1$ be the vertices of P .

Suppose first that there is an integer i , $1 \leq i \leq k$, such that $z_{i-1}z_i \in E(F)$. If $z_{i-1}z_i \in E(P_1)$, then, deleting from F the edges $z_{i-1}z_i$, xx_1^1 and xx_2^1 and adding the edges $x_1^1x_2^1$, xz_{i-1} and xz_i (not excluding the possible case $i = 1$), we get a contradiction with the minimality of P . Similarly we show that $z_{i-1}z_i \notin E(P_2)$ and hence $z_{i-1}z_i \notin E(F)$ for any i , $1 \leq i \leq k$, i.e., no two consecutive vertices of P are consecutive on F .

We now consider the subgraph $\langle z_1, x, z_1^-, z_1^+ \rangle$, where z_1^-, z_1^+ are the predecessor and successor of z_1 on F . If $z_1^-z_1^+ \in E(G)$, then, deleting from F the edges $z_1z_1^-, z_1z_1^+$ and xz_0 and adding the edges z_0z_1 , z_1x and $z_1^-z_1^+$, we get a flower that contradicts the minimality of P . Hence, $z_1^-z_1^+ \notin E(G)$. Since $\langle z_1, x, z_1^-, z_1^+ \rangle$ cannot be an induced claw centred at z_1 , we have $xz_1^- \in E(G)$ or $xz_1^+ \in E(G)$. We distinguish the following four cases.

Case	Deleted edges	Added edges
$xz_1^- \in E(G), z_1 \in V(P_1)$	$z_1z_1^-, xx_1^1, xx_1^2$	$xz_1^-, xz_1, x_1^1x_1^2$
$xz_1^- \in E(G), z_1 \in V(P_2)$	$z_1z_1^-, xx_2^1, xx_2^2$	$xz_1^-, xz_1, x_2^1x_2^2$
$xz_1^+ \in E(G), z_1 \in V(P_1)$	$z_1z_1^+, xx_1^1, xx_1^2$	$xz_1^+, xz_1, x_1^1x_1^2$
$xz_1^+ \in E(G), z_1 \in V(P_2)$	$z_1z_1^+, xx_2^1, xx_2^2$	$xz_1^+, xz_1, x_2^1x_2^2$

In each of these cases we get a contradiction with the minimality of P . ■

Corollary 7. *Let G be a claw-free graph which is not locally disconnected. Then G has the flower property if and only if G is hamiltonian.*

Proof. Follows immediately from Theorem 6.

Remark 8. It is easy to observe that if G is a locally disconnected claw-free graph, then, for every $x \in V(G)$, $\langle N(x) \rangle$ consists of two vertex disjoint cliques and hence G is a line graph. Moreover, if $G = L(H)$, then G is locally disconnected if and only if H is triangle-free. Thus, according to Theorem 6, for the proof of the flower conjecture in claw-free graphs, it remains to prove it in the case that G is a line graph of a triangle-free graph. Hence we have the following corollary.

Corollary 9. *Let G be a claw-free graph that is not a line graph of a triangle-free graph. Then G has the flower property if and only if G is hamiltonian.*

REFERENCES

- [1] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications (Macmillan, London and Elsevier, New York, 1976).
- [2] H. Fleischner, *The square of every two-connected graph is hamiltonian*, J. Combin. Theory (B) **16** (1974) 29–34.
- [3] H. Fleischner, *In the squares of graphs, hamiltonicity and pancyclicity, hamiltonian connectedness and panconnectedness are equivalent concepts*, Monatshefte für Math. **82** (1976) 125–149.
- [4] A. Kaneko, *Research problem*, Discrete Math., (to appear).
- [5] A. Kaneko and K. Ota, *The flower property implies 1-toughness and the existence of a 2-factor*, Manuscript (unpublished).

Received 28 November 1994