

## ON-LINE RANKING NUMBER FOR CYCLES AND PATHS

ERIK BRUOTH AND MIRKO HORŇÁK

*Department of Geometry and Algebra*  
*P.J. Šafárik University, Jesenná 5*  
*041 54 Košice, Slovakia*

**e-mail:** ebruoth@duro.upjs.sk

**e-mail:** hornak@turing.upjs.sk

### Abstract

A  $k$ -ranking of a graph  $G$  is a colouring  $\varphi : V(G) \rightarrow \{1, \dots, k\}$  such that any path in  $G$  with endvertices  $x, y$  fulfilling  $\varphi(x) = \varphi(y)$  contains an internal vertex  $z$  with  $\varphi(z) > \varphi(x)$ . On-line ranking number  $\chi_r^*(G)$  of a graph  $G$  is a minimum  $k$  such that  $G$  has a  $k$ -ranking constructed step by step if vertices of  $G$  are coming and coloured one by one in an arbitrary order; when colouring a vertex, only edges between already present vertices are known. Schiermeyer, Tuza and Voigt proved that  $\chi_r^*(P_n) < 3 \log_2 n$  for  $n \geq 2$ . Here we show that  $\chi_r^*(P_n) \leq 2 \lfloor \log_2 n \rfloor + 1$ . The same upper bound is obtained for  $\chi_r^*(C_n), n \geq 3$ .

**Keywords:** ranking number, on-line vertex colouring, cycle, path.

**1991 Mathematics Subject Classification:** 05C15.

### 1 INTRODUCTION

In this article we deal with simple finite undirected graphs. For formal reasons we also use the empty graph  $K_0 = (\emptyset, \emptyset)$ . A  $k$ -ranking of a graph  $G$  is a vertex colouring of  $G$  which takes as colours integers  $1, \dots, k$  in such a way that, whenever a path of  $G$  has endvertices of the same colour, it contains an internal vertex with a greater colour. If  $k$  is not specified, we speak simply about a *ranking*. Evidently, a ranking is a proper vertex colouring and a  $k$ -ranking of a connected graph uses  $k$  at most once. Rankings are important in the parallel Cholesky factorization of matrices (Liu [3]) and also in VLSI layout (Leiserson [2]).

*Ranking number*  $\chi_r(G)$  of a graph  $G$  is a minimum  $k$  such that  $G$  has a  $k$ -ranking. The problem of finding the ranking number of an arbitrary graph is NP-complete, see Llewelyn et al. [4]. Katchalski et al. [1] proved, among other results on trees, that  $\chi_r(P_n) = \lfloor \log_2 n \rfloor + 1$  for  $n \geq 1$ . They have also an upper bound for the ranking number of a planar graph  $G$ , namely  $\chi_r(G) \leq 3(\sqrt{6} + 2)\sqrt{|V(G)|}$ .

In an *on-line* version of the problem vertices of a graph  $G$  are coming in an arbitrary order. They are coloured one by one in such a way that only a local information concerning edges between already present vertices is known in a moment when a colour for a vertex is to be chosen. Schiermeyer et al. [5] showed that, for  $n \geq 2$ , there is an on-line algorithm providing a ranking of  $n$ -vertex path, for which the maximum used number is smaller than  $3 \log_2 n$ , independently from arriving order of vertices. Our main aim is to show that this number is  $\leq 2 \lfloor \log_2 n \rfloor + 1$ .

For a graph  $G$  and a set  $W \subseteq V(G)$  let  $G\langle W \rangle$  be the subgraph of  $G$  induced by  $W$ . The notation  $C_n$  and  $P_n$  is used for  $n$ -vertex cycle and  $n$ -vertex path, respectively.

For integers  $p, q$  we denote by  $[p, q]$  the set of all integers  $r$  with  $p \leq r \leq q$ , and by  $[p, \infty)$  the set of all integers  $r$  with  $p \leq r$ .

The *length* of a finite sequence  $A$  (i.e., the number of terms of  $A$ ), is denoted by  $|A|$ . For finite sequences  $A = (a_1, \dots, a_m)$  and  $B = (b_1, \dots, b_n)$  let  $AB = (a_1, \dots, a_m, b_1, \dots, b_n)$  be the *concatenation* of  $A$  and  $B$  (in this order); the concatenation can be generalized to any finite number of finite sequences. The concatenation is, clearly, associative, and we will use  $\Pi_{i=1}^k A_i$  for the concatenation of finite sequences  $A_1, \dots, A_k$  (in this order).

Now, let us describe our on-line version of the ranking problem more precisely. An *input sequence* for a graph  $G$  is any sequence of vertices of  $G$  containing all vertices of  $G$  exactly once. Let  $\text{Is}(G)$  be the set of all input sequences for  $G$  and let  $Y = \Pi_{i=1}^n (y_i) \in \text{Is}(G)$ . Vertices  $y_1, \dots, y_n$  are coloured in this order one by one in the following way: We denote by  $G(Y, y_i)$  the graph  $G\langle \{y_j : j \in [1, i]\} \rangle$  induced by all vertices that come in  $Y$  not later than  $y_i$  does,  $i \in [1, n]$ . We colour  $y_1$  with an arbitrary positive integer. In the moment when  $y_i$ ,  $i \in [2, n]$ , is to be coloured, only the graph  $G(Y, y_i)$  and a ranking of  $G(Y, y_{i-1})$  is known; the colour of  $y_i$  has to be chosen in such a way that a ranking of  $G(Y, y_i)$  results (without altering “old” colours).

We would like to analyze all possibilities of forming a ranking of a graph  $G$  in the above on-line fashion. To that aim, we denote by  $\mathcal{Q}$  the set of all quadruples  $(G, H, \varphi, x)$  such that  $G$  is a non-empty graph,  $H$  is an induced subgraph of  $G$  with  $|V(H)| = |V(G)| - 1$ ,  $\varphi$  is a ranking of  $H$

and  $\{x\} = V(G) - V(H)$ . We say that two quadruples  $(G, H, \varphi, x)$  and  $(G', H', \varphi', x')$  are *equivalent* (and we do not distinguish them in  $\mathcal{Q}$ ) if there is an isomorphism  $\iota$  between  $G$  and  $G'$  which maps  $H$  onto  $H'$  (so that  $\iota(x) = x'$ ) and an automorphism  $\alpha'$  of  $H'$  such that for any  $y \in V(H)$  it holds  $\varphi(y) = \varphi'(\alpha'(\iota(y)))$ . A *ranking algorithm* is a mapping  $\mathcal{A} : \mathcal{Q} \rightarrow [1, \infty)$  such that, for any  $(G, H, \varphi, x) \in \mathcal{Q}$ ,  $\varphi \cup \{(x, \mathcal{A}(G, H, \varphi, x))\}$  is a ranking of  $G$ .

Let  $\mathcal{A}$  be a ranking algorithm, let  $G$  be a graph and let  $Y = \Pi_{i=1}^n(y_i) \in \text{Is}(G)$ . The algorithm  $\mathcal{A}$  provides a ranking  $\text{rank}(\mathcal{A}, G, Y, y_i)$  of the graph  $G(Y, y_i), i \in [1, n]$ , recurrently as follows:

$$\text{rank}(\mathcal{A}, G, Y, y_1) := \{(y_1, \mathcal{A}(K_1, K_0, \emptyset, y_1))\},$$

$$\text{rank}(\mathcal{A}, G, Y, y_i) := \text{rank}(\mathcal{A}, G, Y, y_{i-1})$$

$$\cup \{(y_i, \mathcal{A}(G(Y, y_i), G(Y, y_{i-1}), \text{rank}(\mathcal{A}, G, Y, y_{i-1}), y_i))\}, \quad i \in [2, n].$$

We denote by  $\text{rank}(\mathcal{A}, G, Y)$  the ranking  $\text{rank}(\mathcal{A}, G, Y, y_n)$  of the graph  $G(Y, y_n) = G$  provided by the algorithm  $\mathcal{A}$  if the vertices of  $G$  are coming in the input sequence  $Y$ . Clearly, the ranking  $\text{rank}(\mathcal{A}, G, Y, y_i)$  is a restriction of the ranking  $\text{rank}(\mathcal{A}, G, Y)$  to the graph  $G(Y, y_i), i \in [1, n]$ . By  $\max(\mathcal{A}, G, Y)$  we will denote the maximum number attributed to a vertex of  $G$  by  $\text{rank}(\mathcal{A}, G, Y)$  and by  $\max(\mathcal{A}, G)$  the maximum of  $\max(\mathcal{A}, G, Y)$  over all  $Y \in \text{Is}(G)$ . The *on-line ranking number*  $\chi_r^*(G)$  of the graph  $G$  is the minimum of  $\max(\mathcal{A}, G)$  over all ranking algorithms  $\mathcal{A}$ . Evidently, for any graph  $G$  and any ranking algorithm  $\mathcal{A}$  we have

$$\chi_r(G) \leq \chi_r^*(G) \leq \max(\mathcal{A}, G).$$

**Proposition 1.** *If  $G_1$  is an induced subgraph of  $G_2$  and  $\mathcal{A}$  is a ranking algorithm, then  $\max(\mathcal{A}, G_1) \leq \max(\mathcal{A}, G_2)$ .*

**Proof.** Consider an input sequence  $Y_1 = \Pi_{i=1}^n(y_i) \in \text{Is}(G_1)$  such that  $\max(\mathcal{A}, G_1, Y_1) = \max(\mathcal{A}, G_1)$  and an arbitrary input sequence  $Y_2$  of the graph  $G_2 \setminus (V(G_2) - V(G_1))$ . Then  $Y_1 Y_2 \in \text{Is}(G_2)$ , and we have  $\text{rank}(\mathcal{A}, G_2, Y_1 Y_2, y_n) = \text{rank}(\mathcal{A}, G_1, Y_1)$ , so that  $\max(\mathcal{A}, G_2) \geq \max(\mathcal{A}, G_2, Y_1 Y_2) \geq \max(\mathcal{A}, G_1, Y_1) = \max(\mathcal{A}, G_1)$ . ■

**Corollary 2.** *If  $G_1$  is an induced subgraph of  $G_2$ , then  $\chi_r^*(G_1) \leq \chi_r^*(G_2)$ .* ■

## 2 REDUCTION

A natural *greedy* algorithm  $\mathcal{G}$  (called also First Fit Algorithm) is determined by the requirement that, for any  $(G, H, \varphi, x) \in \mathcal{Q}$ ,  $\mathcal{G}(G, H, \varphi, x)$  is the minimum positive integer  $k$  such that  $\varphi \cup \{(x, k)\}$  is a ranking of  $G$ . In other words, we can describe  $\mathcal{G}$  as follows: A colour  $l \in [1, \infty)$  is *forbidden* for  $x$  if the colouring  $\psi = \varphi \cup \{(x, l)\}$  produces a  $(u, v)$ -path  $P$  in  $G$  with  $\psi(u) = \psi(v) = \max\{\psi(y) : y \in V(P)\}$  (clearly,  $x \in V(P)$ ). The greedy algorithm colours  $x$  with the smallest colour that is not forbidden for  $x$ . Evidently, the colour  $\max\{\varphi(y) : y \in V(H)\} + 1$  is not forbidden for  $x$ . That is why, we know that for any graph  $G$  and any input sequence  $Y \in \text{Is}(G)$  the ranking  $\text{rank}(\mathcal{G}, G, Y)$  of  $G$  uses every integer from the interval  $[1, \max(\mathcal{G}, G, Y)]$  at least once.

Now we are going to analyze how  $\mathcal{G}$  works for cycles and paths. For that purpose suppose that  $G = C_n$ ,  $n \in [3, \infty)$ , or  $G = P_n$ ,  $n \in [1, \infty)$ , with  $V(G) = \{x_i : i \in [1, n]\}$  and  $E(G) \supseteq \{x_i x_{i+1} : i \in [1, n-1]\}$  (there is an equality in this inclusion if  $G = P_n$ , and, if  $G = C_n$ , there is an additional edge  $x_n x_1$ ). Sometimes it will be necessary to use for indices arithmetics modulo  $n$ , i.e.,  $x_{i-n} = x_i = x_{i+n}$  for any  $i \in [1, n]$ .

As an example, consider the input sequence  $Y = (x_6, x_7, x_3, x_5, x_2, x_4, x_1) \in \text{Is}(C_7) = \text{Is}(P_7)$ . We have  $\text{rank}(\mathcal{G}, C_7, Y) = \{(x_6, 1), (x_7, 2), (x_3, 1), (x_5, 3), (x_2, 2), (x_4, 4), (x_1, 5)\}$  and  $\text{rank}(\mathcal{G}, P_7, Y)$  differs from  $\text{rank}(\mathcal{G}, C_7, Y)$  only by attributing 1 to  $x_1$ .

An important role in our analysis is played by the following reduction process: We suppose that  $G = C_n$ ,  $n \in [5, \infty)$ , or  $G = P_n$ ,  $n \in [2, \infty)$ ,  $Y \in \text{Is}(G)$  and  $\varphi = \text{rank}(\mathcal{G}, G, Y)$ . A vertex  $x_i \in V(G)$  is said to be a *survivor* of  $G$  (with respect to the input sequence  $Y$ ) if  $\varphi(x_i) \geq 2$ ; if  $\varphi(x_i) = 1$ , it is a *non-survivor*. We transform  $G$  into a non-empty graph  $R(G, Y)$  homeomorphic to  $G$  as follows: We delete from  $G$  all non-survivors and we join by a new edge any two survivors having a non-survivor as a common neighbour (i.e., we delete all non-survivors of degree 1 and we “smooth out” all non-survivors of degree 2). We can do this because it is easy to see that the number of survivors is always positive and, in the case  $G = C_n$ , it is  $\geq 3$ . The input sequence  $Y$  induces in a natural way an input sequence  $R(Y, G)$  for the graph  $R(G, Y)$  – we simply delete from  $Y$  all non-survivors.

If  $Y \in \text{Is}(C_7)$  is as above, then  $R(C_7, Y) = C_5$ ,  $R(Y, C_7) = (x_7, x_5, x_2, x_4, x_1)$  and  $R(P_7, Y) = P_4$ ,  $R(Y, P_7) = (x_7, x_5, x_2, x_4)$ .

**Lemma 3.** Let  $G = C_n$ ,  $n \in [5, \infty)$ , or  $G = P_n$ ,  $n \in [2, \infty)$ , let  $Y \in \text{Is}(G)$ ,  $\varphi = \text{rank}(\mathcal{G}, G, Y)$ ,  $\dot{G} = R(G, Y)$ ,  $\dot{Y} = R(Y, G)$  and  $\dot{\varphi} = \text{rank}(\mathcal{G}, \dot{G}, \dot{Y})$ . Then, for any survivor  $x_i$  of  $G$  with respect to  $Y$ , it holds  $\dot{\varphi}(x_i) = \varphi(x_i) - 1$ .

**Proof.** Consider a sequence  $Y' \in \text{Is}(G)$  in which all non-survivors (with respect to  $Y$ ) come first (in an arbitrary order) and then all survivors (with respect to  $Y$ ) come in the order induced by that of  $Y$ . It is easy to see that  $\varphi = \text{rank}(\mathcal{G}, G, Y')$ .

Let  $Y' = \Pi_{i=1}^n(y_i)$  and let  $y_s$  be the first survivor with respect to  $Y'$  (and  $Y$  as well). We are going to show by induction on  $i$  that  $\dot{\varphi}(y_i) = \varphi(y_i) - 1$  for any  $i \in [s, n]$ . Obviously,  $\dot{\varphi}(y_s) = 1 = 2 - 1 = \varphi(y_s) - 1$ .

Now suppose that  $i \in [s + 1, n]$  and that  $\dot{\varphi}(y_j) = \varphi(y_j) - 1$  for every  $j \in [s, i - 1]$ . Note that survivors  $y_j, y_k$  with  $j, k \in [s, i]$ ,  $j \neq k$ , are joined by a path  $P$  in  $G(Y', y_i)$  if and only if they are joined in  $\dot{G}(\dot{Y}, y_i)$  by the path  $\dot{P}$  such that  $V(\dot{P}) = V(P) - \{y_l : l \in [1, s - 1]\}$ . Hence, by the induction hypothesis and the fact that  $\varphi(y_l) = 1$  for any  $l \in [1, s - 1]$ , a colour  $a \in [2, \infty)$  is forbidden for  $y_i$  in  $G(Y, y_i)$  by a path  $P$  if and only if the colour  $a - 1$  is forbidden for  $y_i$  in  $\dot{G}(\dot{Y}, y_i)$  by the corresponding path  $\dot{P}$ . Since  $\varphi(y_i) \geq 2$ , we obtain  $\dot{\varphi}(y_i) = \varphi(y_i) - 1$ , as necessary. ■

We define a *section* of our graph  $G$  as follows: A section of  $P_n$  is any sequence  $\Pi_{i=j}^k(x_i)$  of vertices of  $P_n$  with  $j, k \in [1, n]$  and  $j \leq k$ . A section of  $C_n$  is any sequence  $\Pi_{i=j}^k(x_i)$  of vertices of  $C_n$  with  $j, k \in [1 - n, 2n]$  and  $j \leq k \leq j - 1 + n$ . From the definition we see that a section  $\Pi_{i=j}^k(x_i)$  consists of  $k + 1 - j \leq n$  distinct vertices of  $G$  and that  $x_i x_{i+1}$  is an edge of  $G$  for every  $i \in [j, k - 1]$ . An *endsection* of  $P_n$  is any section of  $P_n$  containing an endvertex of  $P_n$ . The *type* of a section  $\Pi_{i=j}^k(x_i)$  (with respect to the ranking  $\varphi = \text{rank}(\mathcal{G}, G, Y)$ ) is the sequence formed from  $\Pi_{i=j}^k(\varphi(x_i))$  by replacing any term  $\varphi(x_i)$  fulfilling  $\varphi(x_i) \geq 3$  with  $3+$ . The ranking  $\varphi = \text{rank}(\mathcal{G}, G, Y)$  determines two types of vertices in  $G$ : a vertex  $x \in V(G)$  is *high* (with respect to  $\varphi$ ), if  $\varphi(x) \geq 3$ , otherwise it is *low*. A section of  $G$  containing only high [low] vertices, which is maximal (non-extendable with respect to this property), is called a *high* [low] section of  $G$ . The *defect* of a section  $S$  of  $G$  is the difference  $\text{def}(S)$  between the number of low vertices in  $S$  and the number of high vertices in  $S$ . The *defect* of a graph  $G$  is the difference  $\text{def}(G)$  between the number of low vertices in  $V(G)$  and the number of high vertices in  $V(G)$ , i.e., the defect of (any) section  $S$  of  $G$  with  $|S| = |V(G)|$ .

**Lemma 4.** Let  $G = C_n$ ,  $n \in [3, \infty)$ , or  $G = P_n$ ,  $n \in [1, \infty)$ , let  $Y \in \text{Is}(G)$ ,  $\varphi = \text{rank}(\mathcal{G}, G, Y)$  and  $q \in [1, n]$ .

1. If  $\Pi_{i=q}^{q+3}(x_i)$  is a section of  $G$ , then there are  $j, k \in [q, q+3]$  such that  $\varphi(x_j) = 1$  and  $\varphi(x_k) \geq 3$ .
2. If  $\Pi_{i=q}^{q+2}(x_i)$  is such a section of  $G$  that  $\varphi(x_{q+1}) = 2$ , then  $\min\{\varphi(x_q), \varphi(x_{q+2})\} = 1$ .
3. If  $G = P_n$  and  $\varphi(x_1) \geq 2$ , then  $n \geq 2$  and  $\varphi(x_2) = 1$ .
4. If  $G = P_n$  and  $\varphi(x_1) \geq 3$ , then  $n \geq 3$ ,  $\varphi(x_2) = 1$  and  $\varphi(x_3) = 2$ .
5. If  $G = P_n$  and  $\varphi(x_n) \geq 2$ , then  $n \geq 2$  and  $\varphi(x_{n-1}) = 1$ .
6. If  $G = P_n$  and  $\varphi(x_n) \geq 3$ , then  $n \geq 3$ ,  $\varphi(x_{n-1}) = 1$  and  $\varphi(x_{n-2}) = 2$ .
7. If  $\Pi_{i=q}^{q+2}(x_i)$  is a section of  $G$  of type  $(3+, 3+, 3+)$ , then  $\Pi_{i=q-2}^{q+4}(x_i)$  also is a section of  $G$  and it is of type  $(2, 1, 3+, 3+, 3+, 1, 2)$ .
8. If  $\Pi_{i=q}^{q+3}(x_i)$  is a section of  $G$  of type  $(3+, 3+, 1, 3+)$ , then  $\Pi_{i=q-2}^{q+5}(x_i)$  also is a section of  $G$  and it is of type  $(2, 1, 3+, 3+, 1, 3+, 1, 2)$  or  $(2, 1, 3+, 3+, 1, 3+, 2, 1)$ .
9. If  $\Pi_{i=q}^{q+3}(x_i)$  is a section of  $G$  of type  $(3+, 1, 3+, 3+)$ , then  $\Pi_{i=q-2}^{q+5}(x_i)$  also is a section of  $G$  and it is of type  $(1, 2, 3+, 1, 3+, 3+, 1, 2)$  or  $(2, 1, 3+, 1, 3+, 3+, 1, 2)$ .
10. If  $G = P_n$ ,  $n \geq 3$ ,  $\varphi(x_1) = 1$  and  $\varphi(x_3) \geq 3$ , then  $\varphi(x_2) = 2$ .
11. If  $G = P_n$ ,  $n \geq 3$ ,  $\varphi(x_n) = 1$  and  $\varphi(x_{n-2}) \geq 3$ , then  $\varphi(x_{n-1}) = 2$ .
12. If  $G = P_n$  and  $\Pi_{i=q}^{q+1}(x_i)$  is a section of  $G$  of type  $(3+, 3+)$ , then  $n \geq 6$  and  $q \in [3, n-3]$ .
13. If  $G = P_n$  and  $\Pi_{i=q}^{q+2}(x_i)$  is a section of  $G$  of type  $(3+, 1, 3+)$ , then  $n \geq 7$  and  $q \in [3, n-4]$ .
14. If  $G = P_n$  and  $\Pi_{i=q}^{q+2}(x_i)$  is a section of  $G$  of type  $(3+, 3+, 2)$ , then  $n \geq 7$  and  $q \in [3, n-4]$ .
15. If  $G = P_n$  and  $\Pi_{i=q}^{q+2}(x_i)$  is a section of  $G$  of type  $(2, 3+, 3+)$ , then  $n \geq 7$  and  $q \in [3, n-4]$ .

**Proof.** 1. The existence of  $k$  follows immediately from the definition of a ranking. As concerns the existence of  $j$ , we may suppose that  $\min\{\varphi(x_q), \varphi(x_{q+3})\} \geq 2$  – otherwise we are done. Let  $x_j$  be that vertex from among  $x_{q+1}, x_{q+2}$ , which comes sooner in  $Y$ . Then, clearly,  $\varphi(x_j) = 1$ .

2. Suppose that  $\varphi(x_q) \geq 3$  and  $\varphi(x_{q+2}) \geq 3$ . We have  $\varphi(x_{q+1}) \neq 1$ , hence the colour 1 is forbidden for  $x_{q+1}$  because of an  $(x_s, x_t)$ -path with  $\varphi(x_s) = \varphi(x_t) = a$  containing  $x_{q+1}$  as an internal vertex. Clearly,  $\min\{\varphi(x_s), \varphi(x_t)\} \geq 3$  implies  $a \geq 3$ . Then, however, the colour 2 is forbidden for  $x_{q+1}$ , too, a contradiction.

3. The inequality  $n \geq 2$  is immediate. Also, we cannot have  $\varphi(x_2) \geq 2$ , because then  $\varphi(x_1) = 1$ .

4. Since  $\varphi$  uses each colour from  $[1, \max(\mathcal{G}, G, Y)]$  at least once, we have  $n \geq 3$ . From 3 we know that  $\varphi(x_2) = 1$ . The assumption  $\varphi(x_3) \geq 3$  then would lead to  $\varphi(x_1) = 2$ .

5,6. The situation is symmetric with that of 3 and 4.

7. Since, clearly,  $n \geq 5$  (1 and 2 are used at least once), the reduction process applies and yields  $\dot{G} = R(G, Y)$ ,  $\dot{Y} = R(Y, G)$ ,  $\dot{\varphi} = \text{rank}(\mathcal{G}, \dot{G}, \dot{Y})$ .

Suppose first that  $G = P_n$ . From 4 and 6 it follows that  $\Pi_{i=q-1}^{q+3}(x_i)$  is a section of  $G$  and from 1 we obtain  $\varphi(x_{q-1}) = \varphi(x_{q+3}) = 1$ . From Lemma 3 we know that  $\dot{\varphi}(x_i) = \varphi(x_i) - 1 \geq 2$  for  $i = q, q+1, q+2$ ; then, from 3 and 5 (applied to the ranking  $\dot{\varphi}$  of  $\dot{G}$ ) we see that  $x_q$  and  $x_{q+2}$  are not endvertices of  $\dot{G}$ , which (since  $x_{q-1}$  and  $x_{q+3}$  as non-survivors are not in  $\dot{G}$ ) means that  $x_{q-2}, x_{q+4} \in V(\dot{G})$  and  $S = \Pi_{i=q-2}^{q+4}(x_i)$  is a section of  $G$ . Then, from 1 applied to  $\dot{\varphi}$ , we have  $\dot{\varphi}(x_{q-2}) = \dot{\varphi}(x_{q+4}) = 1$ , and, by Lemma 3 again,  $S$  is a section of  $G$  of type  $(2, 1, 3+, 3+, 3+, 1, 2)$ .

If  $G = C_n$ , then, by 1,  $\Pi_{i=q-1}^{q+3}(x_i)$  is a section of  $G$  of type  $(1, 3+, 3+, 3+, 1)$ , hence  $n \geq 6$  ( $\varphi$  as a ranking is a proper vertex colouring of  $G$ ). If  $n \geq 7$ , then, as in the case  $G = P_n$ , we conclude that  $S$  is a section of  $G$  of type  $(2, 1, 3+, 3+, 3+, 1, 2)$ . If  $n = 6$ ,  $\Pi_{i=q-2}^{q+3}(x_i)$  would be a section of  $G$  of type  $(2, 1, 3+, 3+, 3+, 1)$ . Then, however,  $\dot{G} = C_4$  and  $\dot{\varphi} = \text{rank}(\mathcal{G}, C_4, \dot{Y})$  uses 1 exactly once in contradiction with the following fact (which can be easily checked out):

(\*) For any input sequence  $\bar{Y} \in \text{Is}(C_4)$  the ranking  $\text{rank}(\mathcal{G}, C_4, \bar{Y})$  uses 1 exactly twice.

8. As in 7, we use the reduction process leading to  $\dot{G}, \dot{Y}$  and  $\dot{\varphi}$ . In the case  $G = P_n$ , we obtain from 4 and 6 that  $\Pi_{i=q-1}^{q+4}(x_i)$  is a section of  $G$ . Clearly, because of 7, we have  $\varphi(x_{q-1}) \leq 2$ . Then, the assumption  $q = 2$  would mean  $\varphi(x_q) \leq 2$ , a contradiction. Thus,  $q \geq 3$ . Suppose that  $\varphi(x_{q-1}) = 2$ . If  $x_q$  comes in  $Y$  before  $x_{q+1}$ , then  $\varphi(x_q) = 1$ , and, if  $x_{q+1}$  comes in  $Y$  before  $x_q$ , then  $\varphi(x_{q+1}) \leq 2$ , in both cases a contradiction. Thus,  $\varphi(x_{q-1}) = 1$ ; we cannot have  $\varphi(x_{q-2}) \geq 3$ , because in such a case, by Lemma 3,  $(x_{q-2}, x_q, x_{q+1}, x_{q+3})$  would be a section of  $\dot{G}$  contradicting 1 (applied to  $\dot{\varphi}$ ). The mentioned contradiction yields  $\varphi(x_{q-2}) = 2$ . If  $\varphi(x_{q+4}) \geq 3$ , considering the section  $(x_q, x_{q+1}, x_{q+3}, x_{q+4})$  of  $\dot{G}$  supplies an analogous contradiction. So, there are two possibilities for  $\varphi(x_{q+4})$ : If  $\varphi(x_{q+4}) = 1$ , then  $n \geq q + 5$ , as  $n = q + 4$  would imply  $\varphi(x_{q+3}) = 2$ , a contradiction; then, by 1 applied to  $\dot{\varphi}$ , we get  $\dot{\varphi}(x_{q+5}) = 1$  and  $\varphi(x_{q+5}) = 2$ .

The assumption  $\varphi(x_{q+4}) = 2$  excludes  $n = q + 4$ , by 5. Then, by 2,  $\varphi(x_{q+5}) \geq 3$  is impossible and  $\varphi(x_{q+5}) = 1$ , as necessary.

Now, consider the case  $G = C_n$ . Since  $\varphi$  must use 2, we have  $n \geq 5$ . However,  $n = 5$  is impossible, because then  $\dot{\varphi}$  would contradict (\*). Thus,  $n \geq 6$  and, just as in the case  $G = P_n$ , we can show that  $\varphi(x_{q-1}) = 1$  and  $\varphi(x_{q-2}) = 2$ . That is why,  $n = 6$  is impossible – use again (\*) for  $\dot{\varphi}$ . We cannot have  $\varphi(x_{q+4}) \geq 3$  from the same reason as applied for  $G = P_n$ . Then the assumption  $n = 7$  would lead to  $\varphi(x_{q+4}) = 1$  ( $\varphi$  is proper) and a contradiction involving once more (\*) for  $\dot{\varphi}$ . Finally, for  $n \geq 8$ , the reasoning for  $G = P_n$  can be repeated, and we are done.

9. Use the symmetry with the situation of 8.

10,11. The proof is immediate.

12. From 4 we see that  $q \geq 2$ . If  $\varphi(x_{q-1}) \geq 2$ , from 3 we obtain  $q \geq 3$ . If  $\varphi(x_{q-1}) = 1$ , then  $q \geq 3$ , since  $q = 2$  would lead to  $\varphi(x_q) = 2$ . Thus,  $q \geq 3$  in any case, and, because of the symmetry of the type  $(3+, 3+)$ , we have  $n \geq q + 3$ , too.

13. The proof is analogous to that of 12.

14. By 5 we have  $n \geq q + 3$ , so that 1 yields  $\varphi(x_{q+3}) = 1$ . Now,  $n = q + 3$  is impossible – this would mean that  $\varphi(x_{q+1}) = 1$ . To show that  $q \geq 3$ , proceed as in 12.

15. Symmetry with 14. ■

For a ranking algorithm  $\mathcal{A}$ , we will denote by  $f_i(\mathcal{A}, G, Y), i \in [1, \infty)$ , the number of vertices that are coloured with  $i$  by  $\text{rank}(\mathcal{A}, G, Y)$ .

**Lemma 5.** *Let  $G = C_n, n \in [3, \infty)$ , or  $G = P_n, n \in [1, \infty)$ , and let  $Y \in \text{Is}(G)$ . Then the sequence  $\{f_i(\mathcal{G}, G, Y)\}_{i=1}^\infty$  is non-increasing.*

**Proof.** We proceed by induction on  $n$ . First, it is straightforward to see that  $f_1(\mathcal{G}, P_1, Y) = 1$  for (the unique)  $Y \in \text{Is}(P_1)$ ,  $f_i(\mathcal{G}, C_3, Y) = 1, i = 1, 2, 3$ , for any  $Y \in \text{Is}(C_3)$ , and  $f_1(\mathcal{G}, C_4, Y) = 2$  (in fact, this is (\*)),  $f_i(\mathcal{G}, C_4, Y) = 1, i = 2, 3$ , for any  $Y \in \text{Is}(C_4)$ .

Now, suppose that  $n \geq 5$  (if  $G = C_n$ ) or  $n \geq 2$  (if  $G = P_n$ ) and that  $\{f_i(\mathcal{G}, G', Y')\}_{i=1}^\infty$  is a non-increasing sequence for any graph  $G'$  homeomorphic to  $G$  with  $|V(G')| < n$  and any input sequence  $Y' \in \text{Is}(G')$ . Let  $\varphi = \text{rank}(\mathcal{G}, G, Y)$ ,  $\dot{G} = R(G, Y), \dot{Y} = R(Y, G), \dot{\varphi} = \text{rank}(\mathcal{G}, \dot{G}, \dot{Y})$ . From Lemma 3 we know that, for any  $i \in [2, \infty)$ , we have  $f_{i-1}(\mathcal{G}, \dot{G}, \dot{Y}) = f_i(\mathcal{G}, G, Y)$  and, since  $|V(\dot{G})| < n$  (there are non-survivors of  $G$  with respect to  $Y$ , because  $\varphi$  uses 1 at least once), from the induction hypothesis we obtain  $f_i(\mathcal{G}, G, Y) = f_{i-1}(\mathcal{G}, \dot{G}, \dot{Y}) \geq f_i(\mathcal{G}, \dot{G}, \dot{Y}) = f_{i+1}(\mathcal{G}, G, Y)$ .



Put  $V_i = \{x \in V(G) : \varphi(x) = i\}$ ,  $i = 1, 2$ , and consider a mapping  $\alpha : V_2 \rightarrow V_1$  defined in such a way that  $x\alpha(x)$  is an edge of  $G$  for any  $x \in V_2$ . From Lemmas 4.2, 4.3 and 4.5 it follows that  $\alpha$  is well defined. Moreover, the definition of a ranking implies that  $\alpha$  is an injection; thus,  $f_1(\mathcal{G}, G, Y) = |V_1| \geq |V_2| = f_2(\mathcal{G}, G, Y)$ , which represents the last wanted inequality. ■

Suppose that  $G \in \{C_n, P_n\}$ ,  $n \in [4, \infty)$  and let  $\tilde{G}$  be the cycle defined as follows:  $\tilde{G} = G$  if  $G = C_n$ ,  $\tilde{G} = G + x_n x_1$  if  $G = P_n$ . The ranking  $\varphi$  of  $G$  is then also a vertex colouring of  $\tilde{G}$ , which, if  $G = P_n$ , in general is *not* a ranking of  $\tilde{G}$  (it may be even not proper). When working with  $\tilde{G}$ , types of vertices will be always related to this colouring “inherited” from the ranking  $\varphi$  of the “underlying” graph  $G$ . With respect to this colouring we define also high and low sections of  $\tilde{G}$ .

By Lemma 4.1, rotating around  $\tilde{G}$  we meet alternately high and low sections; their possible lengths are between 1 and 3 if  $G = C_n$ , and between 1 and 6 if  $G = P_n$  (and in this case, due to Lemmas 4.4 and 4.6, only one section, namely low, obtained by joining two low endsections of  $P_n$ , can be of length greater than 3). Let  $s$  be the number of high (and low as well) sections of  $\tilde{G}$ . We will denote those sections  $S_i, i \in [1, 2s]$ , in such a way that  $S_1$  is that high section of maximum length which contains a vertex  $x_t$  with minimum index  $t$ . Consider a (high) section  $S_{2i-1}, i \in [1, s]$ . Starting from it and rotating around  $\tilde{G}$  in the sense of the orientation of  $\tilde{G}$  given by the growing order of sections indices (modulo  $2s$ ) we take all sections until we arrive at the first high section not shorter than  $S_{2i-1}$  (maybe  $S_{2i-1}$  itself). The section which arises by the concatenation of those sections (in their natural “rotating” order) is called the *closure* of  $S_{2i-1}$  and is denoted by  $\text{cl}(S_{2i-1})$ . Thus,  $\text{cl}(S_{2i-1}) = \Pi_{k=2i-1}^{2j} S_k$ , where  $j \in [i, s]$  is (uniquely) chosen to fulfill the conditions  $|S_{2k-1}| < |S_{2i-1}|$  for each  $k \in [i+1, j]$  and  $|S_{2j+1}| \geq |S_{2i-1}|$  (note that  $j \leq s$  because  $S_1$  is the longest high section).

In our example we have  $S_1 = (x_4, x_5)$ ,  $\text{cl}(S_1) = S_1 S_2 = (x_4, x_5, x_6, x_7)$ ,  $S_3 = (x_1)$ ,  $\text{cl}(S_3) = S_3 S_4 = (x_1, x_2, x_3)$  (for  $G = C_7$ ) and  $S_1 = (x_4, x_5)$ ,  $\text{cl}(S_1) = S_1 S_2 = (x_4, x_5, x_6, x_7, x_1, x_2, x_3)$  (for  $G = P_7$ ).

**Lemma 6.** *The closure of any high section of  $\tilde{G}$  has a nonnegative defect.*

**Proof.** Let  $S_{2i-1}$  be a high section of  $\tilde{G}$  and suppose that  $\text{cl}(S_{2i-1}) = \Pi_{k=2i-1}^{2j} S_k$ .

1. If  $|S_{2i-1}| = 1$ , then  $\text{cl}(S_{2i-1}) = S_{2i-1} S_{2i}$  and  $\text{def}(\text{cl}(S_{2i-1})) = |S_{2i}| - 1 \geq 0$ .

2. Assume that  $|S_{2i-1}| = 2$ . Evidently, we have  $\text{def}(\text{cl}(S_{2i-1})) = \text{def}(S_{2i-1} S_{2i}) + \sum_{k=i+1}^j \text{def}(S_{2k-1} S_{2k})$ . Since  $2 = |S_{2i-1}| > |S_{2k-1}| = 1$

for each  $k \in [i+1, j]$ , the sum consists of nonnegative summands  $|S_{2k}| - 1$ . Thus, we are done if  $\text{def}(S_{2i-1}S_{2i}) \geq 0$ .

If  $\text{def}(S_{2i-1}S_{2i}) = |S_{2i}| - |S_{2i-1}| < 0$ , then, necessarily,  $|S_{2i}| = 1$ . From Lemmas 4.2, 4.3 and 4.5 we then see that  $S_{2i}$  is of type (1). Suppose that  $S_{2i-1}S_{2i} = \Pi_{k=q}^{q+2}(x_k)$ ,  $q \in [1, n]$ , and consider the section  $S = \Pi_{k=q}^{q+3}(x_k)$  of  $\tilde{G}$  of type  $(3+, 3+, 1, 3+)$ . If  $S$  is also a section of  $G$ , then, by Lemma 4.8,  $S_{2i+1}$  is of length 1 (so that  $j \geq i+1$ ) and  $\text{def}(S_{2i+1}S_{2i+2}) \geq 1$ , which implies  $\text{def}(\text{cl}(S_{2i-1})) \geq -1 + 1 + \sum_{k=i+2}^j (|S_{2k}| - 1) \geq 0$ . If  $S$  is not a section of  $G$ , then  $G = P_n$  and  $n \in [q, q+2]$ . However,  $n = q$  is impossible by Lemma 4.4,  $n = q+1$  by Lemma 4.5 and  $n = q+2$  by Lemma 4.11.

3. Now, let  $|S_{2i-1}| = 3$ . First we show that, for any  $l \in [i, j]$ , we have  $d_l = \text{def}(\Pi_{k=2i-1}^{2l} S_k) \geq -1$ , and, if  $d_k = -1$  for every  $k \in [i, l]$ , then either  $S_{2l}$  is of type (1,2) or  $S_{2l-1}S_{2l}$  is of type (3+,1). We proceed by induction on  $l$ . If  $l = i$  and  $S_{2i-1} = \Pi_{k=q}^{q+2}(x_k)$  with  $q \in [1, n]$ , we know that  $S_{2i-1}$  is a section of  $G$  (otherwise  $G = P_n$  and  $n \in [q, q+1]$ , which contradicts Lemma 4.3 or Lemma 4.5). Thus, we can use Lemma 4.7, from which it follows that  $d_i \geq -1$  and  $d_i = -1$  only if  $S_{2i}$  is of type (1,2).

Suppose that  $j > i$  and that our statement is true for some  $l \in [i, j-1]$  (so that  $|S_{2l+1}| \leq 2$ ). Since  $d_{l+1} = d_l + |S_{2l+2}| - |S_{2l+1}| \geq d_l + 1 - 2 = d_l - 1$ , to prove the statement for  $l+1$  it is sufficient to analyze the case  $d_l = -1$ . (If  $d_l \geq 0$ , then  $d_{l+1} \geq -1$  and it is not true that  $d_k = -1$  for any  $k \in [i, l+1]$ .) By the induction hypothesis, we have two possibilities:

a)  $S_{2l} = \Pi_{k=q}^{q+1}(x_k)$ , where  $q \in [1, n]$ , is of type (1,2). If  $|S_{2l+1}| = 2$ , then  $\Pi_{k=q}^{q+5}(x_k)$  is the section of the graph  $G$  ( $G = P_n$  and  $n \in [q, q+4]$ ) would be in contradiction with one of Lemmas 4.3, 4.5 and 4.11) and  $S_{2l+2}$  is neither of type (1,1) nor of type (2,2) (this would mean  $G = P_n$  and  $n = q+4$ ). Next, by Lemma 4.1,  $S_{2l+2}$  cannot be of type (2) or (2,1), and, by Lemma 4.8, of type (1); thus, either  $d_{l+1} = d_l = -1$  and  $S_{2l+2}$  is of type (1,2) (as necessary) or  $d_{l+1} \geq 0$  (and there is nothing more to prove). Let  $|S_{2l+1}| = 1$ . The only interesting case (in which  $d_{l+1} = -1$ ) is that with  $|S_{2l+2}| = 1$ . Then, because of Lemma 4.2 or 4.5,  $S_{2l+2}$  is not of type (2), and, consequently,  $S_{2l+1}S_{2l+2}$  is of type (3+,1), as needed.

b)  $S_{2l-1}S_{2l} = (x_q, x_{q+1})$ , where  $q \in [1, n]$ , is of type (3+,1). If  $|S_{2l+1}| = 2$ , then  $\Pi_{k=q}^{q+3}(x_k)$  is the section of the graph  $G$  ( $G = P_n$  and  $n \in [q, q+2]$ ) would be in contradiction with one of Lemmas 4.3, 4.6 and 4.10). Then, by Lemma 4.9,  $\varphi(x_{q+4}) = 1$  and  $\varphi(x_{q+5}) = 2$ , so that either  $d_{l+1} = -1$  and  $S_{2l+2}$  is of type (1,2) or  $d_{l+1} = 0$ ; in both cases we are done. Suppose  $|S_{2l+1}| = 1$ . It is sufficient to deal with the case  $d_{l+1} = -1$ , in which

$|S_{2l+2}| = 1$ . If  $S_{2l+1}S_{2l+2}$  is of type  $(3+, 1)$ , we are done. On the other hand, by Lemmas 4.2 and 4.5,  $S_{2l+2}$  cannot be of type (2) and our statement is completely proved.

Now, it is clear that we cannot have  $d_k = -1$  for each  $k \in [i, j]$ , because  $|S_{2j+1}| = 3$  and, by Lemma 4.7, the type of  $S_{2j}$  ends up with  $(2, 1)$ . Thus, there exists (uniquely determined)  $l \in [i, j]$  fulfilling  $d_l \geq 0$  and  $d_k = -1$  for any  $k \in [i, l-1]$ . If  $l = j$ , then  $\text{def}(\text{cl}(S_{2i-1})) = d_l \geq 0$ . Suppose therefore  $l < j$ . If  $|S_{2k-1}| = 1$  for any  $k \in [l+1, j]$ , then  $\text{def}(\text{cl}(S_{2i-1})) = d_l + \sum_{k=l+1}^j (|S_{2k}| - 1) \geq 0$ . If  $|S_{2m-1}| = 2$  for some  $m \in [l+1, j]$  and  $|S_{2k-1}| = 1$  for any  $k \in [l+1, m-1]$ , delete from the sequence  $\Pi_{k=m}^j(2k-1)$  all terms  $2k-1$  with  $|S_{2k-1}| = 1$  and denote by  $\Pi_{k=1}^q(p_k)$  the resulting sequence. Then it is easy to see directly from the definitions that  $\Pi_{k=2m-1}^{2j} S_k = \Pi_{k=1}^q \text{cl}(S_{p_k})$  and, as  $S_{p_k}$  is a high section of length 2, by 2 we have  $\text{def}(\text{cl}(S_{p_k})) \geq 0$  for each  $k \in [1, q]$ . That is why,  $\text{def}(\text{cl}(S_{2i-1})) = d_l + \sum_{k=l+1}^{m-1} (|S_{2k}| - 1) + \sum_{k=1}^q \text{def}(\text{cl}(S_{p_k})) \geq 0$ . ■

**Theorem 7.** Let  $G = C_n$ ,  $n \in [3, \infty)$ , or  $G = P_n$ ,  $n \in [1, \infty)$ , and let  $Y \in \text{Is}(G)$ . Then  $\sum_{i=1}^2 f_i(\mathcal{G}, G, Y) \geq \lceil n/2 \rceil$  and  $f_1(\mathcal{G}, G, Y) \geq \lceil \lceil n/2 \rceil / 2 \rceil$ .

**Proof.** The assertion is immediate if  $n \leq 3$ . If  $n \in [4, \infty)$ , consider the graph  $\tilde{G}$  and its high and low sections  $S_i, i \in [1, 2s]$ , as defined before Lemma 6. Let  $\Pi_{i=1}^m(l_i)$  be the increasing sequence of indices of all longest high sections of  $\tilde{G}$ . Then, obviously, the section  $\Pi_{i=1}^m \text{cl}(S_{l_i})$  contains all vertices of  $V(\tilde{G}) = V(G)$ , and so, by Lemma 6,  $\sum_{i=1}^2 f_i(\mathcal{G}, G, Y) - \sum_{i=3}^\infty f_i(\mathcal{G}, G, Y) = \text{def}(G) = \text{def}(\Pi_{i=1}^m \text{cl}(S_{l_i})) = \sum_{i=1}^m \text{def}(\text{cl}(S_{l_i})) \geq 0$ . Thus, we have  $n = \sum_{i=1}^2 f_i(\mathcal{G}, G, Y) + \sum_{i=3}^\infty f_i(\mathcal{G}, G, Y) \leq 2 \sum_{i=1}^2 f_i(\mathcal{G}, G, Y)$  and the first inequality follows. The remaining one comes from Lemma 5, since  $2f_1(\mathcal{G}, G, Y) \geq \sum_{i=1}^2 f_i(\mathcal{G}, G, Y) \geq \lceil n/2 \rceil$ . ■

**Proposition 8.** If  $k \in [1, \infty)$  and  $l \in [3, \infty)$ , there exist  $q \in [1, \infty)$  and  $r \in [3, \infty)$  such that  $\max(\mathcal{G}, P_q) = k$  and  $\max(\mathcal{G}, C_r) = l$ .

**Proof.** Suppose that there is no  $q \in [1, \infty)$  such that  $\max(\mathcal{G}, P_q) = k$ . Since, evidently,  $\max(\mathcal{G}, P_n) = n, n = 1, 2$ , we have  $k \geq 3$ . The sequence  $\{\chi_r(P_n)\}_{n=1}^\infty = \{\lfloor \log_2 n \rfloor + 1\}_{n=1}^\infty$  is unbounded and  $\max(\mathcal{G}, P_n) \geq \chi_r^*(P_n) \geq \chi_r(P_n)$ , hence there exists  $q \in [1, \infty)$  such that  $\max(\mathcal{G}, P_q) \geq k+1$ ; without loss of generality, we may suppose that  $q$  is minimum with this property, i.e.,  $\max(\mathcal{G}, P_n) \leq k-1$  for any  $n \in [1, q-1]$ . Consider such an input sequence  $Y \in \text{Is}(P_q)$  that  $\max(\mathcal{G}, P_q, Y) = \max(\mathcal{G}, P_q)$ . Clearly,  $q \geq k+1 \geq 4$ , so we may use our reduction process yielding  $\dot{G} = R(P_q, Y)$ ,  $\dot{Y} = R(Y, P_q)$ . We

have  $|V(\dot{G})| < q$ , which implies  $\max(\mathcal{G}, \dot{G}) \leq k - 1$ . On the other hand, by Lemma 3, the maximum number used by  $\dot{\varphi}$  is by 1 smaller than that used by  $\varphi$ , i.e.,  $\max(\mathcal{G}, \dot{G}, \dot{Y}) = \max(\mathcal{G}, P_q, Y) - 1 = \max(\mathcal{G}, P_q) - 1 \geq (k+1) - 1 = k$ , hence  $\max(\mathcal{G}, \dot{G}) \geq \max(\mathcal{G}, \dot{G}, \dot{Y}) \geq k$ , a contradiction.

For cycles we proceed analogously using the fact that  $\max(\mathcal{G}, C_3) = 3$  and that the reduction process applies if the number of vertices of  $C_n$  is at least 5. Note that also the sequence  $\{\chi_r(C_n)\}_{n=1}^\infty$  is unbounded, because  $P_{n-1}$  is an induced subgraph of  $C_n$ , and so (as can be easily seen)  $\chi_r(P_{n-1}) \leq \chi_r(C_n)$  for any  $n \in [3, \infty)$ . ■

From Proposition 8 we conclude that the numbers

$$\begin{aligned} f(k) &:= \min\{n \in [1, \infty) : \max(\mathcal{G}, P_n) = k\}, \quad k \in [1, \infty), \\ g(k) &:= \min\{n \in [3, \infty) : \max(\mathcal{G}, C_n) = k\}, \quad k \in [3, \infty) \end{aligned}$$

( $f(k)$  was introduced in [5]) are correctly defined. It is easily seen that  $f(k) = k$  for  $k = 1, 2, 3$  and  $g(3) = 3$ . Clearly, from Lemma 3 it follows that  $f(k) \neq f(l)$  and  $g(k) \neq g(l)$  for  $k \neq l$ . However, we can say more:

**Proposition 9.** *The sequences  $\{f(k)\}_{k=1}^\infty$  and  $\{g(k)\}_{k=3}^\infty$  are increasing.*

**Proof.** In the case of paths use simply Proposition 1 and the fact that  $P_m$  is an induced subgraph of  $P_n$  if  $m < n$ .

For cycles suppose that  $\{h(k)\}_{k=3}^\infty$  is the increasing sequence created by rearranging  $\{g(k)\}_{k=3}^\infty$ , that  $\{h(k)\} \neq \{g(k)\}$  and that  $k$  is the minimum index with  $h(k) \neq g(k)$ . Since  $g(3) = h(3) = 3$ , we have  $k \geq 4$  and  $h(k) = g(l) < g(k)$  with  $k < l$ . For  $n = g(l)$  take an input sequence  $Y \in \text{Is}(C_n)$  fulfilling  $\max(\mathcal{G}, C_n, Y) = l$ . As  $l \geq 5$ ,  $\dot{G} = R(C_n, Y)$  and  $\dot{Y} = R(Y, C_n)$  are well defined. Then, by Lemma 3,  $\max(\mathcal{G}, \dot{G}, \dot{Y}) = \max(\mathcal{G}, C_n, Y) - 1 = l - 1 \geq k$  and, since  $|V(\dot{G})| < |V(C_n)| = g(l)$ , we have  $g(l - 1) \leq |V(\dot{G})| < g(l) < g(k)$  and  $l - 1 > k$ . Now,  $g(l - 1) > g(k - 1)$  is in contradiction with  $h(k) = g(l)$  and  $g(l - 1) < g(k - 1)$  contradicts the minimality of  $k$ . ■

**Corollary 10.** *For any  $k, n \in [1, \infty)$  it holds  $\max(\mathcal{G}, P_n) = k$  if and only if  $n \in [f(k), f(k + 1) - 1]$ .*

**Proof.** A consequence of Propositions 1 and 9. ■

For cycles the situation is unclear, but we conjecture that, analogously, for any  $k, n \in [3, \infty)$ ,  $\max(\mathcal{G}, C_n) = k$  if and only if  $n \in [g(k), g(k + 1) - 1]$ .

Theorem 7 has an important consequence:

**Theorem 11.** *Let  $k \in [1, \infty)$ ,  $l \in [3, \infty)$ ,  $q \in [2, \infty)$  and  $r \in [7, \infty)$ .*

1. *If  $f(k) \geq q$ , then  $f(k + 2i) \geq q \cdot 2^i$  for any  $i \in [0, \infty)$ .*
2. *If  $g(k) \geq r$ , then  $g(k + 2i) \geq r \cdot 2^i$  for any  $i \in [0, \infty)$ .*

**Proof.** 1. We proceed by induction on  $i$ . For  $i = 0$  there is nothing to prove, so we suppose that  $i \in [1, \infty)$  and  $f(k + 2i - 2) \geq q \cdot 2^{i-1}$ . With respect to Proposition 9 it is sufficient to show that  $\max(\mathcal{G}, P_n, Y) \leq k + 2i - 1$  for any  $n \in [q \cdot 2^{i-1} + 2, q \cdot 2^i - 1]$  and any  $Y \in \text{Is}(P_n)$ . Since  $n \geq q \cdot 2^{i-1} + 2 \geq q + 2 \geq 4$ , the reduction process applied to  $P_n$  and  $Y$  yields  $\dot{G} = R(P_n, Y)$  and  $\dot{Y} = R(Y, P_n)$ . The ranking  $\text{rank}(\mathcal{G}, P_n, Y)$  is a proper vertex colouring of  $P_n$ , hence  $f_1(\mathcal{G}, P_n, Y) \leq \lceil n/2 \rceil$ ,  $|V(\dot{G})| = n - f_1(\mathcal{G}, P_n, Y) \geq n - \lceil n/2 \rceil = \lfloor n/2 \rfloor \geq 2$ , so that the reduction process applied to  $\dot{G}$  and  $\dot{Y}$  leads to  $\ddot{G} = R(\dot{G}, \dot{Y})$  and  $\ddot{Y} = R(\dot{Y}, \dot{G})$ . By a repeated use of Lemma 3 we see that  $|V(\ddot{G})| = n - \sum_{i=1}^2 f_i(\mathcal{G}, P_n, Y)$ , hence, by Theorem 7,  $|V(\ddot{G})| \leq n - \lceil n/2 \rceil = \lfloor n/2 \rfloor \leq q \cdot 2^{i-1} - 1$ , and, by the induction hypothesis,  $\max(\mathcal{G}, \ddot{G}, \ddot{Y}) \leq \max(\mathcal{G}, \ddot{G}) \leq k + 2i - 3$ . Using Lemma 3 twice then  $\max(\mathcal{G}, P_n, Y) = \max(\mathcal{G}, \dot{G}, \dot{Y}) + 1 = \max(\mathcal{G}, \ddot{G}, \ddot{Y}) + 2 \leq k + 2i - 1$ , as needed.

2. We proceed as in 1 and use the fact that  $f_1(\mathcal{G}, C_n, Y) \leq \lfloor n/2 \rfloor$ , so that  $|V(R(C_n, Y))| \geq n - \lfloor n/2 \rfloor = \lceil n/2 \rceil \geq 5$  for any  $n \in [r \cdot 2^{i-1} + 2, r \cdot 2^i - 1]$ ,  $i \in [1, \infty)$  and any  $Y \in \text{Is}(C_n)$ , which enables us to use the reduction process twice, as above. ■

### 3 INSERTION

Now we are going to show that, in some extent, our reduction process can be inverted. Let  $\mathcal{A}_{m,n}$ ,  $n \in [1, \infty)$ ,  $m \in [0, n]$ , be the set of all non-empty increasing sequences of integers from  $[m, n]$ .

We will analyze in detail the case  $G = P_n$ . For  $A = \Pi_{i=1}^l(a_i) \in \mathcal{A}_{0,n}$  we denote by  $I(P_n, A)$  the path with  $n + l$  vertices constructed as follows: Add to  $V(P_n) = \{x_i : i \in [1, n]\}$   $l$  new vertices (called *newcomers*)  $z_i$ ,  $i \in [1, l]$ . If  $i \in [1, l]$  is such that  $a_i \in [1, n - 1]$ , the newcomer  $z_i$  is inserted between vertices  $x_{a_i}$  and  $x_{a_i+1}$  (i.e., the edge  $x_{a_i}x_{a_i+1}$  is deleted and edges  $x_{a_i}z_i$  and  $z_ix_{a_i+1}$  are added). If  $a_1 = 0$ , the newcomer  $z_1$  is a new endvertex – the edge  $z_1x_1$  is added. Similarly, if  $a_l = n$ , the newcomer  $z_l$  is a new endvertex – the edge  $x_nz_l$  is added. Note that the set of newcomers is an independent set of vertices of  $I(P_n, A)$ . An input sequence  $Y \in \text{Is}(P_n)$  for the path  $P_n$  yields in a natural way an input sequence  $I(P_n, A, Y) = [\Pi_{i=1}^l(z_i)]Y$  for the path  $I(P_n, A)$  – newcomers are coming first ( $z_i$  comes as  $i$ -th,  $i \in [1, l]$ ) and

then vertices of  $P_n$  arrive in the order given by  $Y$ . Consider the ranking  $\varphi = \text{rank}(\mathcal{G}, P_n, Y)$ . An internal vertex  $x_i$  of  $P_n$ ,  $i \in [2, n-1]$ , is *Y-good*, if it comes in  $Y$  as the last from among  $x_{i-1}, x_i, x_{i+1}$ , and  $\varphi(x_{i-1}) = \varphi(x_{i+1})$ . A sequence  $A \in \mathcal{A}_{0,n}$  is *Y-proper*, if any vertex of  $P_n$ , that is not *Y-good*, has in  $I(P_n, A)$  at least one newcomer as a neighbour.

For example, if  $Y$  is the input sequence  $(x_3, x_2, x_5, x_6, x_4, x_1) \in \text{Is}(P_6)$ , there is only one *Y-good* vertex in  $P_6$ , namely  $x_4$  – we have  $\text{rank}(\mathcal{G}, P_6, Y) = \{(x_3, 1), (x_2, 2), (x_5, 1), (x_6, 2), (x_4, 3), (x_1, 1)\}$  ( $x_2$  is not *Y-good*, because it comes in  $Y$  before  $x_1$ ). Thus, the sequence  $A = (1, 2, 5) \in \mathcal{A}_{0,6}$  is *Y-proper* – vertices  $x_i$ ,  $i \in [1, 6] - \{5\}$ , that are not *Y-good*, are “dominated” by newcomers of the graph  $I(P_6, A) = P_9$  (its vertices are successively  $x_1, z_1, x_2, z_2, x_3, x_4, x_5, z_3, x_6$ ). The input sequence  $I(P_6, A, Y)$  is  $(z_1, z_2, z_3, x_3, x_2, x_5, x_6, x_4, x_1)$ .

**Lemma 12.** *Let  $n \in [1, \infty)$ ,  $Y \in \text{Is}(P_n)$ , let a sequence  $A \in \mathcal{A}_{0,n}$  be *Y-proper* and let  $\varphi = \text{rank}(\mathcal{G}, P_n, Y)$ ,  $\hat{G} = I(P_n, A)$ ,  $\hat{Y} = I(P_n, A, Y)$ ,  $\hat{\varphi} = \text{rank}(\mathcal{G}, \hat{G}, \hat{Y})$ . Then  $\hat{\varphi}(z_i) = 1$  for any newcomer  $z_i$ ,  $i \in [1, |A|]$ , and  $\hat{\varphi}(x_i) = \varphi(x_i) + 1$  for any  $i \in [1, n]$ .*

**Proof.** Newcomers of the graph  $\hat{G}$  are attributed 1 by  $\hat{\varphi}$  because they form an independent set of vertices in  $\hat{G}$  and they are coming at the beginning of  $\hat{Y}$ , before all remaining vertices of  $\hat{G}$ .

Let us prove by induction on  $i$  that  $\hat{\varphi}(y_i) = \varphi(y_i) + 1$  for every  $i \in [1, n]$ . The vertex  $y_1$ , clearly, is not *Y-good*, hence it has at least one newcomer as a neighbour and  $\hat{\varphi}(y_1) = 2 = \varphi(y_1) + 1$ .

Suppose that  $i \in [2, n]$  and that  $\hat{\varphi}(y_j) = \varphi(y_j) + 1$  for any  $j \in [1, i-1]$ . Vertices  $y_j, y_k$  with  $j, k \in [1, i], j \neq k$ , are joined by a path  $\hat{P}$  in  $\hat{G}(\hat{Y}, y_i)$  if and only if they are joined in  $G(Y, y_i)$  by the path  $P$  with  $V(P) = V(\hat{P}) - \{z_l : l \in [1, |A|]\}$ . Since  $\hat{\varphi}(z_l) = 1$  for any  $l \in [1, |A|]$ , using the induction hypothesis we see that a colour  $a \in [2, \infty)$  is forbidden for  $y_i$  in  $\hat{G}(\hat{Y}, y_i)$  because of a path  $\hat{P}$  if and only if the colour  $a-1$  is forbidden for  $y_i$  in  $G(Y, y_i)$  because of the corresponding path  $P$ . Moreover, the colour 1 is forbidden for  $y_i$  in  $\hat{G}(\hat{Y}, y_i)$ , too – either a neighbour of  $y_i$  is a newcomer (and so is coloured with 1 in  $\hat{G}(\hat{Y}, y_i)$ ) or both neighbours of  $y_i$  are coloured in  $\hat{G}(\hat{Y}, y_i)$  and they received the same colour. This means that  $\varphi(y_i) = \hat{\varphi}(y_i) - 1$  and we are done.  $\blacksquare$

In our illustrative example with  $n = 6$  we have  $\hat{\varphi} = \text{rank}(\mathcal{G}, P_9, I(P_6, A, Y)) = \{(z_1, 1), (z_2, 1), (z_3, 1), (x_3, 2), (x_2, 3), (x_5, 2), (x_6, 3), (x_4, 4), (x_1, 2)\}$ .

Put  $e_l := 3 \cdot 2^{l-1} - 1$  and  $o_l := 2^{l+1} - 1$ ,  $l \in [1, \infty)$ .

**Theorem 13.** *For any  $l \in [1, \infty)$  there exists*

1. *an input sequence  $Y_{2l} \in \text{Is}(P_{e_l})$  such that  $\max(\mathcal{G}, P_{e_l}, Y_{2l}) = 2l$  and the set of  $Y_{2l}$ -good vertices of the path  $P_{e_l}$  is  $\{x_{3i} : i \in [1, 2^{l-1} - 1]\}$ ;*
2. *an input sequence  $Y_{2l+1} \in \text{Is}(P_{o_l})$  such that  $\max(\mathcal{G}, P_{o_l}, Y_{2l+1}) = 2l + 1$  and the set of  $Y_{2l+1}$ -good vertices of the path  $P_{o_l}$  is  $\{x_{4i} : i \in [1, 2^{l-1} - 1]\}$ .*

**Proof.** Evidently, for  $l = 1$  any input sequence  $Y_2 \in \text{Is}(P_2)$  has all the properties required by 1 (no vertex of  $P_2$  is  $Y_2$ -good). We are going to show that for any  $l \in [1, \infty)$  the existence of  $Y_{2l}$  implies that of  $Y_{2l+1}$  and the existence of  $Y_{2l+1}$  implies that of  $Y_{2l+2}$ . So, suppose that there is an input sequence  $Y_{2l} \in \text{Is}(P_{e_l})$  with properties given by 1. The sequence  $A_{2l} := \Pi_{i=1}^{2^{l-1}} (3i - 2) \in \mathcal{A}_{0, e_l}$  is  $Y_{2l}$ -proper – note that vertices of  $P_{e_l}$ , that are not  $Y_{2l}$ -good, are in pairs  $x_{3i-2}, x_{3i-1}$ , and an “old” edge  $x_{3i-2}x_{3i-1}$  is subdivided by the newcomer  $z_i$ ,  $i \in [1, 2^{l-1}]$ . The graph  $I(P_{e_l}, A_{2l})$  is a path with  $e_l + 2^{l-1} = o_l$  vertices and, if we define  $Y_{2l+1} := I(P_{e_l}, A_{2l}, Y_{2l})$ , then, by Lemma 12,  $\max(\mathcal{G}, P_{o_l}, Y_{2l+1}) = \max(\mathcal{G}, P_{e_l}, Y_{2l}) + 1 = 2l + 1$ . Moreover, any  $Y_{2l}$ -good vertex  $x_{3i}$ ,  $i \in [1, 2^{l-1} - 1]$ , is  $Y_{2l+1}$ -good. There are no other  $Y_{2l+1}$ -good vertices, because, by Lemma 12, any vertex of the path  $P_{e_l}$ , that is  $Y_{2l+1}$ -good and not  $Y_{2l}$ -good, must have two newcomers as neighbours (and the distance between any two newcomers in  $I(P_{o_l}, A_{2l})$  is at least 3). Now, if we rename vertices of  $I(P_{e_l}, A_{2l}) = P_{o_l}$  in our ordinary way (i.e., they will be  $x_i$ ,  $i \in [1, o_l]$ ), then  $x_{3i}$  becomes  $x_{4i}$ ,  $i \in [1, 2^{l-1} - 1]$ , and the set of  $Y_{2l+1}$ -good vertices of  $P_{o_l}$  is  $\{x_{4i} : i \in [1, 2^{l-1} - 1]\}$ .

The sequence  $A_{2l+1} := \Pi_{i=1}^{2^{l-1}} (4i - 3, 4i - 2) \in \mathcal{A}_{0, o_l}$  is  $Y_{2l+1}$ -proper, because vertices of  $P_{o_l}$ , that are not  $Y_{2l}$ -good, occur in triples  $x_{4i-3}, x_{4i-2}, x_{4i-1}$ , which are “dominated” by newcomers  $z_{2i-1}$  and  $z_{2i}$ ,  $i \in [1, 2^{l-1}]$ . The graph  $I(P_{o_l}, A_{2l+1})$  is a path with  $o_l + 2 \cdot 2^{l-1} = e_{l+1}$  vertices and, for  $Y_{2l+2} := I(P_{o_l}, A_{2l+1}, Y_{2l+1})$ , we have, by Lemma 12,  $\max(\mathcal{G}, P_{e_{l+1}}, Y_{2l+2}) = \max(\mathcal{G}, P_{o_l}, Y_{2l+1}) + 1 = 2l + 2$ . Any  $Y_{2l+1}$ -good vertex  $x_{4i}$ ,  $i \in [1, 2^{l-1}]$ , is  $Y_{2l+2}$ -good. Moreover, the vertex  $x_{4i-2}$ ,  $i \in [1, 2^{l-1}]$ , is  $Y_{2l+2}$ -good, too (it has two newcomers as neighbours). There are no other  $Y_{2l+2}$ -good vertices, because there are no more pairs of newcomers which are at the distance 2 apart. Thus, after renaming vertices of  $I(P_{o_l}, A_{2l+1}) = P_{e_{l+1}}$  in our ordinary way (so that  $x_{4i}$  becomes  $x_{6i}$ ,  $i \in [1, 2^{l-1} - 1]$ , and  $x_{4i-2}$  becomes  $x_{6i-3}$ ,  $i \in [1, 2^{l-1}]$ ), the set of  $Y_{2l+2}$ -good vertices of  $P_{e_{l+1}}$  is  $\{x_{3i} : i \in [1, 2^l - 1]\}$ . ■

**Corollary 14.** *For any  $l \in [1, \infty)$ ,  $f(2l) \leq e_l$  and  $f(2l + 1) \leq o_l$ .* ■

Evidently, the reduction process can also be (partially) inverted for cycles. In this case the sequence  $A = \Pi_{i=1}^l(a_i)$ , characterizing positions of newcomers, is from the set  $\mathcal{A}_{1,n}$  (if the original cycle is  $C_n$ ), a newcomer  $z_i$  subdivides the edge  $x_{a_i}x_{a_i+1}$ ,  $i \in [1, l]$ , and there is no restriction on index of a  $Y$ -good vertex. (Recall that, for paths, endvertices are not  $Y$ -good.) Thus, an analogue of Lemma 12 is presented without proof (no new idea is necessary).

**Lemma 15.** *Let  $n \in [3, \infty)$ ,  $Y \in \text{Is}(C_n)$ , let a sequence  $A \in \mathcal{A}_{1,n}$  be  $Y$ -proper and let  $\varphi = \text{rank}(\mathcal{G}, C_n, Y)$ ,  $\hat{G} = I(C_n, A)$ ,  $\hat{Y} = I(C_n, A, Y)$ ,  $\hat{\varphi} = \text{rank}(\mathcal{G}, \hat{G}, \hat{Y})$ . Then  $\hat{\varphi}(z_i) = 1$  for any newcomer  $z_i$ ,  $i \in [1, |A|]$  and  $\hat{\varphi}(x_i) = \varphi(x_i) + 1$  for any  $i \in [1, n]$ . ■*

#### 4 MAIN RESULTS

Now we are able to analyze First Fit Algorithm for cycles and paths in a detailed way.

**Proposition 16.**  $g(4) \leq 5$ ,  $g(5) \leq 7$ ,  $g(6) \leq 10$  and  $g(7) \leq 15$ .

**Proof.** It is easy to check that the sequences  $\hat{A}_3 = (1, 2)$ ,  $\hat{A}_4 = (1, 4)$ ,  $\hat{A}_5 = (2, 5, 7)$  and  $\hat{A}_6 = (1, 3, 5, 7, 9)$  are such that  $\hat{A}_n$  is  $\hat{Y}_n$ -proper,  $n \in [3, 6]$ , if the graph  $\hat{G}_n$  and the input sequence  $\hat{Y}_n$  for  $\hat{G}_n$ ,  $n \in [3, 7]$ , are defined by the following recurrence:  $\hat{G}_3 := C_3$ ,  $\hat{Y}_3 := (x_1, x_2, x_3)$  and  $\hat{G}_{n+1} := I(\hat{G}_n, \hat{A}_n)$ ,  $\hat{Y}_{n+1} := I(\hat{G}_n, \hat{A}_n, \hat{Y}_n)$ ,  $n \in [3, 6]$ . Since  $\max(\mathcal{G}, \hat{G}_3, \hat{Y}_3) = 3$ ,  $\hat{G}_4 = C_5$ ,  $\hat{G}_5 = C_7$ ,  $\hat{G}_6 = C_{10}$ ,  $\hat{G}_7 = C_{15}$  and, by Lemma 15,  $\max(\mathcal{G}, \hat{G}_{n+1}, \hat{Y}_{n+1}) = \max(\mathcal{G}, \hat{G}_n, \hat{Y}_n) + 1$  for  $n \in [3, 6]$ , the proof follows. ■

**Proposition 17.** *If  $k \in [3, \infty)$ , then*

1.  $f(k+1) \geq \min\{n \in [f(k)+1, \infty) : n - \lceil \lceil n/2 \rceil / 2 \rceil \geq f(k)\}$ ;
2.  $g(k+1) \geq \min\{n \in [g(k)+1, \infty) : n - \lceil \lceil n/2 \rceil / 2 \rceil \geq g(k)\}$ .

**Proof.** 1. Suppose that  $f(k+1) = n$ ; by Proposition 9 then  $n \geq f(k) + 1$ . Take an input sequence  $Y \in \text{Is}(P_n)$  such that  $\max(\mathcal{G}, P_n, Y) = k+1$  and put  $\dot{G} = R(P_n, Y)$ ,  $\dot{Y} = R(Y, P_n)$ . For the path  $\dot{G}$  we have, by Theorem 7,  $|V(\dot{G})| = n - f_1(\mathcal{G}, P_n, Y) \leq n - \lceil \lceil n/2 \rceil / 2 \rceil$ , and, by Lemma 3,  $\max(\mathcal{G}, \dot{G}, \dot{Y}) = \max(\mathcal{G}, P_n, Y) - 1 = k$ . Since  $|V(\dot{G})| < n = f(k+1)$ , due to Proposition 9 we obtain  $\max(\mathcal{G}, \dot{G}) = \max(\mathcal{G}, \dot{G}, \dot{Y}) = k$ . Thus,  $|V(\dot{G})| \geq f(k)$  and we see that  $n - \lceil \lceil n/2 \rceil / 2 \rceil \geq f(k)$ .

2. The proof is completely analogous to that of 1. ■



**Theorem 18.**  $f(4) = g(4) = 5, f(5) = g(5) = 7, f(6) = 11, g(6) = 10, f(7) = 15$  and  $14 \leq g(7) \leq 15$ .

**Proof.** Take  $k \in [4, 7]$ . The upper bounds for  $f(k)$  come from Corollary 14 and those for  $g(k)$  from Proposition 16. On the other hand, by Theorem 1 and Lemma 7 of [5],  $f(4) \geq 5$  and  $g(4) \geq 5$ , so that  $f(4) = g(4) = 5$ . Now, by Proposition 17,  $f(5) \geq 7$  and  $g(5) \geq 7$ , which implies  $f(5) = g(5) = 7$ . By Proposition 17 again, we get  $f(6) \geq 10$  and  $g(6) \geq 10$ , yielding  $g(6) = 10$  and, consequently,  $g(7) \geq 14$ .

Suppose that there is an input sequence  $Y \in \text{Is}(P_{10})$  such that  $\max(\mathcal{G}, P_{10}, Y) = 6$  and put  $\varphi = \text{rank}(\mathcal{G}, P_{10}, Y)$ . Since  $f(4) = 5$ , from Lemma 3 (used twice) we see that  $\sum_{i=1}^2 f_i(\mathcal{G}, P_{10}, Y) \leq 5$ . So, with help of Theorem 7,  $\sum_{i=1}^2 f_i(\mathcal{G}, P_{10}, Y) = \sum_{i=3}^6 f_i(\mathcal{G}, P_{10}, Y) = 5$ , and, by Lemma 5,  $f_1(\mathcal{G}, P_{10}, Y) = 3, f_2(\mathcal{G}, P_{10}, Y) = 2$ . Consider the cycle  $\tilde{P}_{10} = C_{10}$  introduced before Lemma 6 and its high and low sections. First we show that there is no high section of  $\tilde{P}_{10}$  of length 3. Suppose there is one; by Lemmas 4.4 and 4.6, this section  $\Pi_{i=q}^{q+2}(x_i)$  must also be a section of  $P_{10}$ . Then, by Lemma 4.7,  $\Pi_{i=q-2}^{q+4}(x_i)$  is a section of  $P_{10}$  of type  $(2, 1, 3+, 3+, 3+, 1, 2)$ . The remaining three vertices of  $P_{10}$  do not form a section of  $P_{10}$ , because two of them are high (otherwise we would obtain a contradiction with one of Lemmas 4.4, 4.6, 4.10 and 4.11). Thus, they form two nonempty endsections of  $P_{10}$ . That containing only one vertex cannot be of type  $(3+)$  ( $P_{10}$  would have an endsection of type  $(3+, 2)$  or  $(2, 3+)$  in contradiction with Lemmas 4.4 and 4.6), hence that of length 2 is of type  $(3+, 3+)$ , which contradicts again Lemmas 4.4 and 4.6.

Denote the number of low sections of  $P_{10}$  and  $\tilde{P}_{10}$  by  $l$  and  $\tilde{l}$ , respectively. Clearly,  $\tilde{l} \geq 3$ , since for  $\tilde{l} = 2$  one of two high sections of  $\tilde{P}_{10}$  would be of length 3. By Lemmas 4.2, 4.3 and 4.5, any low section of  $P_{10}$  contains a vertex coloured with 1, hence  $l \leq 3$ . On the other hand,  $\tilde{l} \leq l$ , and we get  $l = \tilde{l} = 3$ . Thus,  $\tilde{P}_{10}$  has two low sections of type  $(1, 2)$  or  $(2, 1)$ , one low section of type  $(1)$ , two high sections of length 2 and one high section of length 1.

A high section of  $\tilde{P}_{10}$  of length 2 must be a section of  $P_{10}$ , too – otherwise, by Lemmas 4.4 and 4.6,  $\Pi_{i=1}^3(x_i)$  is of type  $(3+, 1, 2)$  and  $\Pi_{i=8}^{10}(x_i)$  is of type  $(2, 1, 3+)$ , so that  $\Pi_{i=4}^7(x_i)$  is of type  $(3+, 3+, 1, 3+)$  or  $(3+, 1, 3+, 3+)$ , which contradicts Lemma 4.8 or Lemma 4.9. Thus, two high sections of  $P_{10}$  of length 2 are, by Lemmas 4.8 and 4.9, separated by a low section of  $P_{10}$  of length 2; let  $\Pi_{i=q}^{q+5}(x_i)$  be the corresponding section of  $P_{10}$  with  $\min\{\varphi(x_i) : i \in \{q, q+1, q+4, q+5\}\} \geq 3$ . Then  $q = 1$  is impossible by

Lemma 4.4,  $q = 2$  by Lemmas 4.3 and 4.10 and, symmetrically,  $q = 4$  by Lemmas 4.5 and 4.11,  $q = 5$  by Lemma 4.6. If  $q = 3$ , one endvertex of  $P_{10}$  is high, which contradicts Lemma 4.4 or Lemma 4.6.

So, we conclude that  $f(6) = 11$ , and then Proposition 17 yields  $f(7) = 15$ . ■

**Corollary 19.** *For  $n = 5, 6$ ,  $\chi_r^*(C_n) = \chi_r^*(P_n) = 4$ .*

**Proof.** Those on-line ranking numbers must be at least 4, by Theorem 1 of [5]. On the other hand, due to Theorem 18,  $\max(\mathcal{G}, C_n) = \max(\mathcal{G}, P_n) = 4$ . ■

Note that, by Theorem 1 of [5], it holds  $\chi_r^*(C_4) = \chi_r^*(P_4) = 3$ . The value of on-line ranking number for simplest cycles and paths (with at most three vertices) is evidently equal to the corresponding number of vertices.

For an input sequence  $Y = \Pi_{i=1}^n(y_i) \in \text{Is}(C_n)$  and  $j \in [0, n-1]$  let  $Y^{+j}$  be the input sequence for the graph  $C_n$  defined by  $Y^{+j} := \Pi_{i=1}^n(y_{i+j})$ .

**Lemma 20.** *If  $n \in [3, \infty)$ ,  $j \in [0, n-1]$  and  $Y \in \text{Is}(C_n)$ , then  $\max(\mathcal{G}, C_n, Y^{+j}) = \max(\mathcal{G}, C_n, Y)$ .*

**Proof.** Evidently,  $V(C_n(Y^{+j}, x_i)) = \{x_{k+j} : x_k \in V(C_n(Y, x_i))\}$  for any  $i \in [1, n]$ . If  $i \in [1, n]$  and  $x_k \in V(C_n(Y, x_i))$ , the ranking  $\text{rank}(\mathcal{G}, C_n, Y^{+j}, x_{i+j})$  attributes to the vertex  $x_{k+j}$  the same colour as the ranking  $\text{rank}(\mathcal{G}, C_n, Y, x_i)$  does to the vertex  $x_k$ , hence the proof follows. ■

**Proposition 21.** *If  $n \in [2, \infty)$ , then  $\max(\mathcal{G}, P_n) \leq \max(\mathcal{G}, C_{n+1}) \leq \max(\mathcal{G}, P_n) + 1$ .*

**Proof.** The first inequality comes from Proposition 1, because  $P_n$  is an induced subgraph of  $C_{n+1}$ .

Take an input sequence  $Y = \Pi_{i=1}^{n+1}(y_i) \in \text{Is}(C_{n+1})$  such that  $\max(\mathcal{G}, C_{n+1}, Y) = \max(\mathcal{G}, C_{n+1})$ . Since  $C_{n+1}(Y, y_n)$  is a path with  $n$  vertices, with respect to Lemma 20 we may suppose that  $V(C_{n+1}(Y, y_n)) = \{x_i : i \in [1, n]\}$ . Then, for the input sequence  $Y^- = \Pi_{i=1}^n(y_i) \in \text{Is}(P_n)$ , we have  $\text{rank}(\mathcal{G}, P_n, Y^-) = \text{rank}(\mathcal{G}, C_{n+1}, Y, y_n)$ . That is why,  $\max(\mathcal{G}, P_n, Y^-) \geq \max(\mathcal{G}, C_{n+1}, Y) - 1 = \max(\mathcal{G}, C_{n+1}) - 1$  (the arrival of  $y_{n+1}$ , the last vertex of  $Y$ , can increase the number of used colours only by 1) and  $\max(\mathcal{G}, C_{n+1}) \leq \max(\mathcal{G}, P_n, Y^-) + 1 \leq \max(\mathcal{G}, P_n) + 1$ . ■

**Corollary 22.** *If  $k \in [3, \infty)$ , then  $g(k) \leq f(k) + 1$ .*

**Proof.** Suppose that  $f(k) = n$ . As  $n \geq k \geq 3$ , Proposition 21 implies  $\max(\mathcal{G}, C_{n+1}) \geq \max(\mathcal{G}, P_n) = k$ , and so, by Proposition 9,  $g(k) \leq n + 1 = f(k) + 1$ . ■

**Theorem 23.** *Let  $i$  be a nonnegative integer. Then*

1.  $11 \cdot 2^i \leq f(2i + 6) \leq 12 \cdot 2^i - 1$ .
2.  $15 \cdot 2^i \leq f(2i + 7) \leq 16 \cdot 2^i - 1$ .
3.  $10 \cdot 2^i \leq g(2i + 6) \leq 12 \cdot 2^i$ .
4.  $14 \cdot 2^i \leq g(2i + 7) \leq 16 \cdot 2^i$ .

**Proof.** Lower bounds come from Theorems 11 and 18. The upper bounds in 1 and 2 follow from Corollary 14, and then those in 3 and 4 from Corollary 22. ■

**Theorem 24.** *Let  $i$  be a nonnegative integer.*

1. *If  $n \in [12 \cdot 2^i - 1, 15 \cdot 2^i - 1]$ , then  $\max(\mathcal{G}, P_n) = 2i + 6$ .*
2. *If  $n \in [15 \cdot 2^i, 16 \cdot 2^i - 2]$ , then  $2i + 6 \leq \max(\mathcal{G}, P_n) \leq 2i + 7$ .*
3. *If  $n \in [16 \cdot 2^i - 1, 22 \cdot 2^i - 1]$ , then  $\max(\mathcal{G}, P_n) = 2i + 7$ .*
4. *If  $n \in [22 \cdot 2^i, 24 \cdot 2^i - 2]$ , then  $2i + 7 \leq \max(\mathcal{G}, P_n) \leq 2i + 8$ .*
5. *If  $n \in [12 \cdot 2^i, 14 \cdot 2^i - 1]$ , then  $\max(\mathcal{G}, C_n) = 2i + 6$ .*
6. *If  $n \in [14 \cdot 2^i, 16 \cdot 2^i - 1]$ , then  $2i + 6 \leq \max(\mathcal{G}, C_n) \leq 2i + 7$ .*
7. *If  $n \in [16 \cdot 2^i, 20 \cdot 2^i - 1]$ , then  $\max(\mathcal{G}, C_n) = 2i + 7$ .*
8. *If  $n \in [20 \cdot 2^i, 24 \cdot 2^i - 1]$ , then  $2i + 7 \leq \max(\mathcal{G}, C_n) \leq 2i + 8$ .*

**Proof.** Because of Proposition 1, the statements 1–4 follow from Theorems 23.1 and 23.2.

If  $n \in [12 \cdot 2^i, \infty)$ , then  $\max(\mathcal{G}, C_n) \geq 2i + 6$ , since otherwise, by Proposition 21,  $\max(\mathcal{G}, P_{n-1}) \leq \max(\mathcal{G}, C_n) \leq 2i + 5$ , which contradicts Theorem 23.1 (with respect to Proposition 1). Thus, 5 and 6 follow from Theorems 23.3 and 23.4. The remaining two statements are proved analogously. ■

**Theorem 25.** *Let  $i$  be a nonnegative integer.*

1. *If  $n \in [12 \cdot 2^i - 1, 15 \cdot 2^i - 1]$ , then  $\chi_r^*(P_n) \leq 2\lfloor \log_2 n \rfloor$ .*
2. *If  $n \in [15 \cdot 2^i, 16 \cdot 2^i - 1]$ , then  $\chi_r^*(P_n) \leq 2\lfloor \log_2 n \rfloor + 1$ .*
3. *If  $n \in [16 \cdot 2^i, 22 \cdot 2^i - 1]$ , then  $\chi_r^*(P_n) \leq 2\lfloor \log_2 n \rfloor - 1$ .*
4. *If  $n \in [22 \cdot 2^i, 24 \cdot 2^i - 2]$ , then  $\chi_r^*(P_n) \leq 2\lfloor \log_2 n \rfloor$ .*

5. If  $n \in [12 \cdot 2^i, 14 \cdot 2^i - 1]$ , then  $\chi_r^*(C_n) \leq 2\lfloor \log_2 n \rfloor$ .
6. If  $n \in [14 \cdot 2^i, 16 \cdot 2^i - 1]$ , then  $\chi_r^*(C_n) \leq 2\lfloor \log_2 n \rfloor + 1$ .
7. If  $n \in [16 \cdot 2^i, 20 \cdot 2^i - 1]$ , then  $\chi_r^*(C_n) \leq 2\lfloor \log_2 n \rfloor - 1$ .
8. If  $n \in [20 \cdot 2^i, 24 \cdot 2^i - 1]$ , then  $\chi_r^*(C_n) \leq 2\lfloor \log_2 n \rfloor$ .

**Proof.** If  $n \in [12 \cdot 2^i - 1, 15 \cdot 2^i - 1]$ , then  $\lfloor \log_2 n \rfloor = i + 3$ , and, by Theorem 24.1,  $\chi_r^*(P_n) \leq \max(\mathcal{G}, P_n) = 2i + 6 = 2\lfloor \log_2 n \rfloor$ , which represents 1. The remaining assertions follow from Theorem 24, too. ■

**Theorem 26.** For any  $n \in [3, \infty)$ ,  $\chi_r(C_n) = \lfloor \log_2(n - 1) \rfloor + 2$ .

**Proof.** First we show that  $\chi_r(C_n) \geq 1 + \chi_r(P_{n-1})$ . Suppose, on the contrary, that  $\chi_r(C_n) = l \leq \chi_r(P_{n-1})$ , and consider an  $l$ -ranking  $\varphi$  of  $C_n$ . If  $x$  is the (only) vertex of  $C_n$  coloured with  $l$ , then  $\varphi - \{(x, l)\}$  is an  $(l - 1)$ -ranking of the path  $P_{n-1} = C_n - x$ , and so  $\chi_r(P_{n-1}) \leq l - 1$ , a contradiction. Thus, according to [1], we have  $\chi_r(C_n) \geq 1 + \lfloor \log_2(n - 1) \rfloor + 1 = \lfloor \log_2(n - 1) \rfloor + 2$ .

Now, take  $k \in [1, \infty)$ ,  $m \in [1, 2^k - 1]$  and  $n = 2^k + m$ . From Lemma 2.1 of [1] it is easy to see that  $\chi_r(P_{2^k}) = k + 1$  and  $\chi_r(P_m) = \lfloor \log_2 m \rfloor + 1 = l(m) \leq k$ . Let  $\varphi_1$  be a  $(k + 1)$ -ranking of  $P_{2^k}$  with  $V(P_{2^k}) = \{x_i : i \in [1, 2^k]\}$  and endvertices  $x_1, x_{2^k}$ , and let  $\varphi_2$  be an  $l(m)$ -ranking of  $P_m$  with  $V(P_m) = \{u_i : i \in [1, m]\}$ , with endvertices  $u_1, u_m$  and with  $V(P_{2^k}) \cap V(P_m) = \emptyset$ . Without loss of generality, by Proposition 2.1 of [1], we may suppose that  $\varphi_1(x_1) = k + 1$ . Let  $C_{2^k+m}$  be the cycle formed from  $P_{2^k} \cup P_m$  by adding the edges  $x_1 u_m$  and  $x_{2^k} u_1$ . The colouring  $\varphi$  of  $C_{2^k+m}$  defined by  $\varphi(x_i) := \varphi_1(x_i)$ ,  $i \in [1, 2^k]$ ,  $\varphi(u_1) = k + 2$  and  $\varphi(u_i) = \varphi_2(u_i)$ ,  $i \in [2, m]$ , is easily seen to be a  $(k + 2)$ -ranking. Thus,  $\chi_r(C_n) \leq k + 2 = \lfloor \log_2(n - 1) \rfloor + 2$ .

For  $k \in [1, \infty)$  let  $\varphi'$  be such a  $(k + 2)$ -ranking of  $P_{2^{k+1}}$  that the (unique) appearance of the colour  $k + 2$  is at an endvertex of  $P_{2^{k+1}}$ . Then,  $\varphi'$  is also a  $(k + 2)$ -ranking of the cycle  $C_{2^{k+1}}$ , which is created from  $P_{2^{k+1}}$  by joining its endvertices by a new edge, and, for  $n = 2^k + 2^k = 2^{k+1}$ , we have  $\chi_r(C_n) \leq k + 2 = \lfloor \log_2(n - 1) \rfloor + 2$ .

So,  $\chi_r(C_n) \leq \lfloor \log_2(n - 1) \rfloor + 2$  for any  $n \in [2^k + 1, 2^{k+1}]$  and any  $k \in [1, \infty)$ , and the desired result follows. ■

**Theorem 27.**

1. For any  $n \in [1, \infty)$ ,  $\lfloor \log_2 n \rfloor + 1 \leq \chi_r^*(P_n) \leq 2\lfloor \log_2 n \rfloor + 1$ .
2. For any  $n \in [3, \infty)$ ,  $\lfloor \log_2(n - 1) \rfloor + 2 \leq \chi_r^*(C_n) \leq 2\lfloor \log_2 n \rfloor + 1$ .

**Proof.** Lower bounds come from the values of  $\chi_r(P_n)$  and  $\chi_r(C_n)$  due to [1] and Theorem 26.

As concerns upper bounds, for  $n \in [12, \infty)$  see Theorem 25; for  $n \leq 11$  use Theorem 18 and the fact that  $f(i) = i, i = 1, 2, 3$ , and  $g(3) = 3$ . ■

First Fit Algorithm is not necessarily optimal when computing  $\chi_r^*(P_n)$ , as shows our next statement.

**Theorem 28.**  $\chi_r^*(P_7) = 4 < 5 = \max(\mathcal{G}, P_7)$ .

**Proof.** According to Theorem 1 of [5], we have  $\chi_r^*(P_7) \geq 4$ . Consider the ranking algorithm  $\mathcal{G}'$  functioning just as  $\mathcal{G}$  does with the only exception: If  $G = P_5$ ,  $H = 2K_2$ ,  $\{x\} = V(G) - V(H)$  and  $\varphi$  is a ranking of  $H$  such that both neighbours of  $x$  (in  $G$ ) are coloured with 2, then  $\mathcal{G}'(G, H, \varphi, x) = 4$  (and not 3, as required by  $\mathcal{G}$ ). We are going to show that  $m' = \max(\mathcal{G}', P_7, Y) \leq 4$  for any  $Y \in \text{Is}(P_7)$ .

First suppose that  $Y = \Pi_{i=1}^7(y_i)$  is such that  $\varphi' = \text{rank}(\mathcal{G}', P_7, Y) \neq \text{rank}(\mathcal{G}, P_7, Y) = \varphi$ . Then  $P_7(Y, y_5) = P_5$  and it is easy to see that any neighbour of (a vertex of)  $P_7(Y, y_5)$  is coloured with 3 and any non-neighbour (at most one) of  $P_7(Y, y_5)$  is coloured with 1 by  $\varphi'$ ; thus,  $m' = 4$ .

Now, assume that  $\varphi' = \varphi$ . If  $y_7 \in \{x_3, x_4, x_5\}$ , then  $P_7(Y, y_6) = P_i \cup P_{6-i}, i \in \{2, 3\}$ . Clearly, the maximum colour used by  $\varphi'_6 = \text{rank}(\mathcal{G}', P_7, Y, y_6)$  is not greater than  $\max\{\max(\mathcal{G}, P_i), \max(\mathcal{G}, P_{6-i})\}$ ; this maximum is equal to 3, by Proposition 1 and  $f(3) = 3$ ,  $f(4) = 5$  (Theorem 18), hence  $m' \leq 4$ .

If  $y_7 \in \{x_1, x_2\}$ , we may suppose that  $\varphi'_6$  uses colour 4 – otherwise we are done.

If  $y_7 = x_2$ , then  $P_7(Y, y_6) = P_1 \cup P_5$  and 4 is used by  $\varphi'_6$  for a vertex of  $P_5$ -component of  $P_7(Y, y_6)$ . If one of  $x_3, x_4$  is coloured with a colour  $\geq 3$ , then, using Lemma 4.3,  $\varphi'(x_2) = 2$ . On the other hand,  $\{\varphi'_6(x_3), \varphi'_6(x_4)\} \neq \{1, 2\}$ , because otherwise at least two vertices from among  $x_5, x_6, x_7$  would be coloured with a colour  $\geq 3$  (3 is used at least once by  $\varphi'_6$ ) in contradiction with one of Lemmas 4.2, 4.7, 4.12 and 4.13.

If  $y_7 = x_1$ , then  $P_7(Y, y_6) = P_6$ . We may assume that  $\varphi'_6(x_2) = 1$  and  $\varphi'_6(x_3) = 2$ , since if not, we would have  $\varphi'(x_1) \leq 2$ . Because of Lemmas 4.1, 4.7, 4.8 and 4.9, exactly two vertices from among  $x_4, x_5, x_6, x_7$  are coloured with a colour  $\geq 3$ . From Lemmas 4.2, 4.12, 4.13 and 4.15 it follows that these are  $x_4$  and  $x_7$ . If  $\varphi'_6(x_4) = 4$ , then  $\varphi'(x_1) = 3$ . Finally, suppose that  $\varphi'_6(x_4) = 3$  and  $\varphi'_6(x_7) = 4$ . Then  $\varphi'_6(x_6) = 1$  and  $\varphi'_6(x_5) = 2$  (by Lemma 4.6),  $x_4$  comes in  $Y$  before  $x_7$  (otherwise  $\varphi'_6(x_7) \leq 3$ ),  $x_4$  comes in  $Y$  after each of  $x_i, i \in \{2, 3, 5, 6\}$  (otherwise  $\varphi'_6(x_4) = 1$ ), which means that  $P_7(Y, y_4) = 2K_2$  and that the vertex  $y_5 = x_4$  has in  $P_7(Y, y_5)$  both

neighbours coloured with 2. This, however, is a contradiction, because in such a case  $4 = \varphi'(y_5) = \varphi'_6(y_5)$ .

The last possibility,  $y_7 \in \{x_6, x_7\}$ , leads to a situation which is symmetric with that of  $y_7 \in \{x_1, x_2\}$ .

Now, to conclude the proof, we use Theorem 18, from which it follows that  $\max(\mathcal{G}, P_7) = 5$ .  $\blacksquare$

**Theorem 29.**  $\chi_r^*(C_7) = 5$ .

**Proof.** By Theorem 1 of [5], we have  $\chi_r^*(C_7) \geq 4$ . We are going to show by the way of contradiction, that  $\chi_r^*(C_7) \geq 5$ ; this, together with  $\max(\mathcal{G}, C_7) = 5$  (Theorem 18), will mean that  $\chi_r^*(C_7) = 5$ .

We know from Theorem 26 that  $\chi_r(C_7) = 4$ . Let  $\varphi$  be a 4-ranking of  $C_7$ . It can be immediately seen that  $\varphi$  uses 3 and 4 exactly once, say, for vertices  $x_i$  and  $x_j$ . Since  $\chi_r(P_4) = 3 = \chi_r(P_5)$ , no component of  $H = C_7 - \{x_i, x_j\}$  can have more than 3 vertices, so that  $H = P_2 \cup P_3$ . Clearly,  $\varphi$  restricted to  $P_3$ -component of  $H$  uses 2 just once, for the internal vertex of that  $P_3$ . Also,  $\varphi$  restricted to  $P_2$ -component of  $H$ , uses 2 once. Thus,  $\varphi$  colours two vertices of  $C_7$  with 2 and two vertices with a colour  $\geq 3$ ; the mutual distance of vertices in those two pairs is 3.

Now, suppose that there is a ranking algorithm  $\mathcal{A}$  such that  $\max(\mathcal{A}, C_7) = 4$ . Consider an input sequence  $Y = \Pi_{i=1}^7(y_i) \in \text{Is}(P_7)$  and the ranking  $\varphi = \text{rank}(\mathcal{A}, C_7, Y)$ . As  $\chi_r(C_7) = 4$ ,  $\varphi$  is a 4-ranking of  $C_7$ . If  $C_7(Y, y_2) = P_2$ , the ranking  $\text{rank}(\mathcal{A}, C_7, Y, y_2)$  must use colours 1 and 2. To see this suppose that a colour  $i \in \{3, 4\}$  is used for a vertex  $y_j$  of  $C_7(Y, y_2)$ . Assume, moreover, that  $C_7(Y, y_k) = P_2 \cup P_{k-2}$ ,  $k = 3, 4, 5$  (we cannot avoid this situation). We have  $\varphi(y_k) \neq i$ ,  $k = 3, 4, 5$ , hence it may happen that  $\varphi(y_k) = 7 - i$  for some  $k \in [3, 5]$  and an endvertex  $y_k$  of  $C_7(Y, y_k)$  – if  $\{\varphi(y_3), \varphi(y_4)\} = \{1, 2\}$ ,  $y_5$  may be an endvertex of  $C_7(Y, y_5)$  with the neighbour coloured with 1. Then, however, the distance between  $y_j$  and  $y_k$ , the vertices coloured with 3 and 4, may be 2 in contradiction with the structure of a 4-ranking of  $C_7$ .

If  $C_7(Y, y_2) = P_2$ ,  $C_7(Y, y_3) = P_3$  and the neighbour of  $y_3$  in  $C_7(Y, y_3)$  is coloured with 1, we have  $\varphi(y_3) = i \in \{3, 4\}$ . It may happen that  $C_7(Y, y_5) = P_3 \cup P_2$ . For vertices of  $P_2$ -component of  $C_7(Y, y_5)$  two from among colours 1, 2 and  $7 - i$  are used. If 2 is used, it may happen that there are two vertices coloured with 2 by  $\varphi$ , whose distance is 2, a contradiction. On the other hand, the presence of  $7 - i$  could yield two vertices of distance 2, coloured with 3 and 4 by  $\varphi$ , a contradiction again.  $\blacksquare$

## 5 OPEN PROBLEMS

There are several open problems which naturally arise from our analysis.

1. Find nontrivial lower bounds for  $\chi_r^*(C_n)$  and  $\chi_r^*(P_n)$ .
2. Which is the minimum  $n$  such that  $\chi_r^*(P_n) = 5$ ?
3. Does there exist  $n \in [8, \infty)$  such that  $\chi_r^*(C_n) < \max(\mathcal{G}, C_n)$ ? If so, which is the minimum  $n$  in such an inequality?
4. Determine  $g(7)$ . (We conjecture that  $g(7) = 15$ .)
5. Prove or disprove that the sequence  $\{\max(\mathcal{G}, C_n)\}_{n=3}^\infty$  is non-decreasing.

**Acknowledgements**

The authors are indebted to an anonymous referee whose hints helped to shorten proofs of Lemmas 3 and 12.

A support of the Slovak VEGA grant 1/4377/97 for the work of the second author is gratefully acknowledged.

## REFERENCES

- [1] M. Katchalski, W. McCuaig and S. Seager, *Ordered colourings*, Discrete Math. **142** (1995) 141–154.
- [2] C.E. Leiserson, *Area-efficient graph layouts (for VLSI)*, in: Proc. 21st Annu. IEEE Symp. on Foundations of Computer Science (1980) 270–281.
- [3] J.W.H. Liu, *The role of elimination trees in sparse factorization*, SIAM J. Matrix Analysis and Appl. **11** (1990) 134–172.
- [4] D.C. Llewelyn, C. Tovey and M. Trick, *Local optimization on graphs*, Discrete Appl. Math. **23** (1989) 157–178.
- [5] I. Schiermeyer, Zs. Tuza and M. Voigt, *On-line rankings of graphs*, submitted.

Received 22 February 1999

Revised 29 October 1999