

## NOTE ON THE WEIGHT OF PATHS IN PLANE TRIANGULATIONS OF MINIMUM DEGREE 4 AND 5

TOMÁŠ MADARAS

*Department of Geometry and Algebra*  
*P.J. Šafárik University*  
*Jesenná 5, 041 54 Košice, Slovak Republic*  
**e-mail:** madaras@duro.upjs.sk

### Abstract

The weight of a path in a graph is defined to be the sum of degrees of its vertices in entire graph. It is proved that each plane triangulation of minimum degree 5 contains a path  $P_5$  on 5 vertices of weight at most 29, the bound being precise, and each plane triangulation of minimum degree 4 contains a path  $P_4$  on 4 vertices of weight at most 31.

**Keywords:** weight of path, plane graph, triangulation.

**2000 Mathematics Subject Classification:** 05C10, 05C38, 52B10.

Throughout this paper we consider connected graphs without loops or multiple edges. Let  $P_r$  ( $C_r$ ) denote a path (cycle) on  $r$  vertices (an  $r$ -path and  $r$ -cycle, in the sequel). A vertex of degree  $m$  is called an  $m$ -vertex, a vertex of degree at least (at most)  $m$  is called a  $+m$ -vertex ( $-m$ -vertex).

The *weight* of the subgraph  $H$  in the graph  $G$  is defined to be the sum of the degrees of the vertices of  $H$  in  $G$ ,  $w(H) = \sum_{v \in V(H)} \deg_G(v)$ . For a family  $\mathcal{G}$  of graphs having a subgraph isomorphic to  $H$ , define the number  $w(H, \mathcal{G}) = \max_{G \in \mathcal{G}} \min_{H \subseteq G} w(H)$ .

The exact value of  $w(H, \mathcal{G})$  is known only for a few graphs and families of graphs. For  $\mathcal{G}(3)$  the family of all 3-connected plane graphs, Ando, Iwasaki and Kaneko [1] proved that  $w(P_3, \mathcal{G}(3)) = 21$ . From the result of Fabrici and Jendroľ [5] it follows that  $w(P_k, \mathcal{G}(3)) \leq 5k^2$  for  $k \geq 1$ ; also, they gave a lower bound for this number as a function of order  $O(k \log(k))$ , see [6]. Recently, the upper bound  $5k^2$  was improved to  $\frac{5}{2}k(k+1)$  for  $k \geq 4$ , see [11]. For  $PHam$  the class of all hamiltonian plane graphs, Mohar [12] proved the exact

value  $w(P_k, PHam) = 6k - 1$ . For  $\mathcal{G}(5)$  and  $\mathcal{T}(5)$  the families of all connected plane graphs/triangulations of minimum degree 5 and subgraphs other than a path, the known exact values are  $w(C_3, \mathcal{G}(5)) = 17$  ([2]),  $w(K_{1,3}, \mathcal{G}(5)) = 23$  ([9]),  $w(K_{1,4}, \mathcal{G}(5)) = 30$ ,  $w(C_4, \mathcal{T}(5)) = 25$ ,  $w(C_5, \mathcal{T}(5)) = 30$  ([4]).

In the following we deal with the weight of paths  $P_k$  in the graphs of the families  $\mathcal{T}(4)$  and  $\mathcal{T}(5)$  (plane triangulations of minimum degree 4 and 5). It is known that  $w(P_2, \mathcal{G}(5)) = 11$  ([13]),  $w(P_3, \mathcal{G}(5)) = 17$  ([8]),  $w(P_4, \mathcal{G}(5)) = 23$  ([9]),  $w(P_3, \mathcal{G}(4)) = 17$  ([1, 3]),  $w(P_4, \mathcal{T}(4)) \leq 4 \cdot 15 = 60$  ([7]). The aim of this paper is to improve the best known upper bound for  $w(P_k, \mathcal{T}(4)), w(P_k, \mathcal{T}(5))$  for small values of  $k$ , showing the following

**Theorem 1.**  $w(P_5, \mathcal{T}(5)) = 29$ .

**Theorem 2.**  $27 \leq w(P_4, \mathcal{T}(4)) \leq 31$ .

**Proof of Theorem 1.** To prove first the inequality  $w(P_5, \mathcal{T}(5)) \leq 29$  suppose that there exists a graph  $G \in \mathcal{T}(5)$  in which every path  $P_5$  has a weight  $w(P_5) > 29$ . We will use the Discharging method. According to the consequence of the Euler formula,

$$\sum_{x \in V(G)} (\deg_G(x) - 6) = -12$$

assign to each vertex  $x \in V(G)$  the initial charge  $\varphi(x) = \deg_G(x) - 6$ . Thus  $\sum_{x \in V(G)} \varphi(x) = -12$ .

Now, we define a local redistribution of charges in a way such that the sum of the charges after redistribution remains the same. This redistribution is performed by the following

**Rule.** Each  $k$ -vertex  $x$ ,  $k \geq 6$ , sends the charge  $\frac{k-6}{m(x)}$  to each adjacent 5-vertex, where  $m(x)$  is the number of 5-vertices adjacent to  $x$ . If  $m(x) = 0$ , no charge is transferred.

**Proposition.** Each +8-vertex sends at least  $\frac{1}{2}$  to each adjacent 5-vertex; each 7-vertex sends at least  $\frac{1}{4}$  to each adjacent 5-vertex.

**Proof.** Consider a 7-vertex  $x$ . Then  $x$  is adjacent to at most four 5-vertices (otherwise two pairs of adjacent 5-vertices are found in the neighbourhood of  $x$ , hence there exists a path  $P_5$  of weight 27, a contradiction). From the similar reason, a 8-vertex (9-vertex) is adjacent to at most four (five) 5-vertices. Since none five consecutive vertices in the neighbourhood of

a  $k$ -vertex,  $k \geq 6$ , can be 5-vertices, every 10-vertex and every 11-vertex is adjacent to at most eight 5-vertices. Then computing  $\frac{k-6}{m(x)}$  yields the desired values of charge. A +12-vertex always sends at least  $\frac{1}{2}$ . ■

We will show that, after redistribution of charges, the new charges  $\tilde{\varphi}(x)$  are non-negative for all  $x \in V(G)$ . This will contradict the fact that  $\sum_{x \in V(G)} \tilde{\varphi}(x) = \sum_{x \in V(G)} \varphi(x) = -12$ . To this end, several cases have to be considered.

*Case 1.*  $x$  is a 5-vertex. Then  $x$  is adjacent to at least two +7-vertices (otherwise, it is adjacent to at least four −6-vertices and there exists a path  $P_5$  with  $w(P_5) \leq 5 + 4 \cdot 6 = 29$ , a contradiction); denote them  $u, v$ . If  $u, v$  are both +8-vertices, then  $\tilde{\varphi}(x) \geq -1 + 2 \cdot \frac{1}{2} = 0$  by Proposition. Otherwise consider the following possibilities:

*Case 1a.*  $u$  is a +8-vertex,  $v$  is a 7-vertex, all other neighbours are 6-vertices. Observe that  $x$  is the only 5-neighbour of  $v$  (otherwise, a 5-path of weight at most  $2 \cdot 5 + 2 \cdot 6 + 7 = 29$  is found). Thus  $\tilde{\varphi}(x) \geq -1 + 1 + \frac{1}{2} > 0$ .


*Case 1b.*  $u, v$  are 7-vertices, all other neighbours are 6-vertices. As above,  $x$  is the only 5-neighbour of  $u, v$ , thus  $\tilde{\varphi}(x) \geq -1 + 2 \cdot 1 > 0$ .

*Case 1c.* Three of the neighbours of  $x$  are 7-vertices, the other ones are −6-vertices. Observe that, for at least one 7-vertex,  $x$  is its only 5-neighbour; thus  $\tilde{\varphi}(x) \geq -1 + 1 + 2 \cdot \frac{1}{4} > 0$ .

*Case 1d.* At least four of the neighbours of  $x$  are 7-vertices. Then  $\tilde{\varphi}(x) \geq -1 + 4 \cdot \frac{1}{4} = 0$ .

*Case 2.*  $x$  is a  $k$ -vertex,  $k \geq 6$ . If  $x$  is adjacent to a 5-vertex, then  $\tilde{\varphi}(x) = k - 6 - m(x) \cdot \frac{k-6}{m(x)} = 0$ ; otherwise  $\tilde{\varphi}(x) = \varphi(x) = k - 6 \geq 0$ .

To prove that the upper bound is best possible consider the so called *edge-hexagon substitution* by which a given plane map  $G$  is transformed into the following plane map  $G'$ : Let every  $x \in V(G)$  be also a vertex of  $G'$ . Assign to every incident pair  $(x, \alpha)$  of a vertex  $x$  and a face  $\alpha$  of  $G$  a new vertex of  $G'$ . Connect two vertices  $x'_1, x'_2 \in V(G')$  by an edge iff either  $x'_1, x'_2$  are assigned to  $(x_1, \alpha_1), (x_2, \alpha_2)$  with  $(x_1, x_2) \in E(G)$  and with  $\alpha_1 = \alpha_2$ , or if  $x'_1$  is assigned to a pair  $(x_1, \alpha_1)$  where  $x'_2 = x_1$ , see Figure (cf. [10]):



Consider a graph of the Archimedean polytope  $(6,6,5)$  and on each its edge apply the edge-hexagon substitution. Into each face of the obtained graph insert a new vertex and join it with new edges to the vertices of the face boundary. In the resulting graph, every 5-path is of the weight of at least 29. ■

**Proof of Theorem 2.** To prove the upper bound suppose that there exists a counterexample  $G$  in which every 4-path has a weight of at least 32.

The following propositions are easy to prove:

**Proposition 1.** *Each  $k$ -vertex with  $7 \leq k \leq 16$  is adjacent to at most  $\lfloor \frac{k}{2} \rfloor - 5$ -vertices.*

**Proposition 2.** *Each  $k$ -vertex,  $k \geq 17$ , is adjacent to at most  $\lfloor \frac{3k}{4} \rfloor - 5$ -vertices.*

We use again the Discharging method. As before, the initial assignment of charges is  $\mu(x) = \deg_G(x) - 6$  for each vertex  $x \in V(G)$ . The local redistribution of charges is based on the following rules:

**Rule 1.** Each  $k$ -vertex  $x$ ,  $k \geq 6$ , sends the charge  $\frac{k-6}{m(x)}$  to each adjacent  $-5$ -vertex;  $m(x)$  is the number of  $-5$ -vertices adjacent to  $x$ . If  $m(x) = 0$ , no charge is transferred.

The following table shows the minimal charge sent by a  $k$ -vertex  $x$ ,  $k \geq 7$ , to an adjacent  $-5$ -vertex, according to Rule 1 (the corresponding values  $m(x)$  are computed due to Propositions 1 and 2):

$k$	7	8	9	10	11	12	13	14	15	16	17	18	19	20	$\geq 21$
$min.charge$	$\frac{1}{3}$	$\frac{1}{2}$	$\frac{3}{4}$	$\frac{4}{5}$	1	1	$\frac{7}{6}$	$\frac{8}{7}$	$\frac{9}{7}$	$\frac{5}{4}$	$\frac{11}{12}$	$\frac{12}{13}$	$\frac{13}{14}$	$\frac{14}{15}$	$\geq 1$

As seen from the table, the only cases when the minimal charge is less than 1 are those with  $k \in \{7, 8, 9, 10, 17, 18, 19, 20\}$ .

Let  $\bar{\mu}$  denote the charge of a vertex after application of Rule 1. A vertex  $y$  is said to be *overcharged* if  $\bar{\mu}(y) > 0$ , and *undercharged* if  $\bar{\mu}(y) < 0$ .

**Rule 2.** Each overcharged  $-5$ -vertex  $x$  sends the charge  $\frac{\bar{\mu}(x)}{\bar{m}(x)}$  to each adjacent undercharged  $4$ -vertex;  $\bar{m}(x)$  is the number of undercharged  $4$ -vertices adjacent to  $x$ . If  $\bar{m}(x) = 0$ , no charge is transferred.

Let  $\tilde{\mu}$  be the charge of vertices after application of Rule 2. Note that  $\bar{\mu}(y) \geq 0$  implies that  $\tilde{\mu}(y) \geq 0$ . We will show that after redistribution of charges we have  $\tilde{\mu}(x) \geq 0$  for each vertex  $x \in G$ , a contradiction. To this end, several cases have to be considered.

*Case 1.* Let  $x$  be a  $k$ -vertex,  $k \geq 6$ . Then either all its charge is sent to adjacent  $-5$ -vertices ( $\bar{\mu}(x) = 0$ ) or there is no transfer from  $x$  and  $\bar{\mu}(x) = k - 6 \geq 0$ .

*Case 2.* Let  $x$  be a  $5$ -vertex. Then  $x$  is adjacent to at least three  $+9$ -vertices (otherwise it is adjacent to at least three  $-8$ -vertices and we can find a  $4$ -path of weight of at most  $8 \cdot 3 + 5 = 29 < 31$ ); hence  $\bar{\mu}(x) \geq -1 + 3 \cdot \frac{3}{4} = \frac{5}{4} > 0$  (thus every  $5$ -vertex is overcharged).

*Case 3.* Let  $x$  be a  $4$ -vertex. Then  $x$  is adjacent to at least two  $+10$ -vertices (otherwise it is adjacent to at least three  $-9$ -vertices and we can find a  $4$ -path of weight of at most  $9 \cdot 3 + 4 = 31$ ). If  $x$  is adjacent to at least three  $+10$ -vertices then  $\bar{\mu}(x) \geq -2 + 3 \cdot \frac{4}{5} = \frac{2}{5} > 0$ ; so, suppose that  $x$  is adjacent to exactly two  $+10$ -vertices  $u, v$ . If both  $u, v$  are  $+21$ -vertices, or one of them is  $+21$ -vertex and the degree of another one is between 11 and 16, or both their degrees are between 11 and 16, then  $u$  and  $v$  send 1 to  $x$  (see Table) and  $\bar{\mu}(x) \geq -2 + 2 \cdot 1 = 0$ . Hence (without loss of generality) it is enough to consider the following possibilities for degrees of  $u, v$  (denote  $y, z$  the remaining neighbours of  $x$ ):

*Case 3.1.* Both  $u, v$  are 10-vertices. Then both  $y, z$  are +8-vertices (otherwise a 4-path of weight of at most  $4 + 2 \cdot 10 + 7 = 31$  is found) and  $\bar{\mu}(x) \geq -2 + 2 \cdot \frac{4}{5} + 2 \cdot \frac{1}{2} > 0$ .

*Case 3.2.*  $u$  is 10-vertex,  $v$  is +11-vertex. Then the sum of degrees of  $y, z$  is at least 18 (otherwise  $x, y, u, z$  form a 4-path of weight of at most  $10 + 4 + 17 = 31$ ); hence, one of them has to be a +9-vertex. Thus  $\bar{\mu}(x) \geq -2 + \frac{4}{5} + \frac{11}{12} + \frac{3}{4} > 0$ .

*Case 3.3.* The degrees of  $u, v$  are between 17 and 20. If some of  $y, z$  is a +7-vertex, then a simple calculation yields  $\bar{\mu}(x) \geq -2 + 2 \cdot \frac{11}{12} + \frac{1}{3} > 0$ ; if some of them is a 5-vertex, the application of Rule 2 yields  $\tilde{\mu}(x) \geq -2 + 2 \cdot \frac{11}{12} + \frac{5}{2} > 0$ . Now, suppose that  $y, z$  are 6- or 4-vertices; then we have to treat several cases:

*Case 3.3a.*  $y, z$  are 4-vertices forming a triangular face with  $x$ . Then  $u, v$  are 20-vertices. Consider the neighbourhood of the vertices  $u, v, y, z$ ; then the vertices  $u, v$  have at least six +6-neighbours. Thus  $\bar{\mu}(x) \geq -2 + 2 \cdot \frac{20-6}{20-6} = 0$ .

*Case 3.3b.*  $y, z$  are 4-vertices not forming a triangular face with  $x$ . Then all their neighbours, except  $x$ , are +20-vertices and we have  $\bar{\mu}(y) \geq -2 + 3 \cdot \frac{14}{15} = \frac{12}{15}$ ,  $\bar{\mu}(z) \geq -2 + 3 \cdot \frac{14}{15} = \frac{12}{15}$ . Hence  $y, z$  are overcharged and using Rule 2 we have  $\tilde{\mu}(x) \geq -2 + 2 \cdot \frac{14}{15} + 2 \cdot \frac{12}{1} > 0$ .

*Case 3.3c.*  $y, z$  are 6-vertices. Considering that each of their neighbours except  $x$  has to be a +16-vertex, it is easy to see that  $u, v$  have at least six +6-neighbours, thus  $\bar{\mu}(x) \geq -2 + 2 \cdot \frac{17-6}{17-6} = 0$ .

*Case 3.3d.*  $y$  is a 4-vertex,  $z$  is a 6-vertex and they do not form a triangular face with  $x$ . Then each neighbour of  $z$ , except for  $x$ , is a +18-vertex, i.e.,  $u, v$  are +18-vertices and, moreover, they have at least six +6-neighbours. Hence  $\bar{\mu}(x) \geq -2 + 2 \cdot \frac{18-6}{18-6} = 0$ .

*Case 3.3e.*  $y$  is a 4-vertex,  $z$  is a 6-vertex and they form a triangular face with  $x$ . Then  $u, v$  are +18-vertices. Let  $u$  be adjacent to  $y$  and  $v$  to  $z$ . Since every neighbour of  $z$ , except  $x$  and  $y$ , has to be a +18-vertex,  $v$  has at least six +6-neighbours and it sends at least 1 to  $x$ . If  $u$  is a 20-vertex, then it has also at least six +6-neighbours, thus  $\bar{\mu}(x) \geq -2 + 2 \cdot 1 = 0$ . So suppose that  $u$  is 18- or 19-vertex not having at least six +6-neighbours.

If  $u$  is a 19-vertex, then in consequence of Proposition 2 it has exactly five +6-neighbours and sends  $\frac{19-6}{19-5} = \frac{13}{14}$  to  $x$ . Denote  $v'_1, v'_2 \dots v'_l$  the neighbours of  $v$  in the cyclical ordering such that  $v'_1 = z, v'_2 = x, v'_3 = u$ . Due to the neighbourhood of  $u$ ,  $v'_4$  has to be a 5-vertex and  $v'_5$  has to be +17-vertex. From this fact we obtain that  $v$  has at least seven +6-neighbours, so it sends at least  $\frac{18-6}{18-7} = \frac{12}{11}$  to  $x$ . Hence  $\bar{\mu}(x) \geq -2 + \frac{13}{14} + \frac{12}{11} = \frac{3}{154} > 0$ .

If  $u$  is a 18-vertex, then every its neighbour, except  $x$  and  $y$ , has to be a +6-vertex (otherwise a 4-path of weight of at most  $2 \cdot 4 + 18 + 5 = 31$  can be found), so  $u$  even sends at least 2 to  $x$  and clearly  $\bar{\mu}(x) > 0$ .

*Case 3.4.* The degree of  $u$  is between 17 and 20, the degree of  $v$  is either between 11 and 16, or is at least 21. According to the similarity to case 3.3 (note that  $v$  always sends at least 1 to  $x$ ) it is enough to consider the cases when  $y$  or  $z$  are neither +7-vertices nor -5-vertices, that means,  $(\deg_G(y), \deg_G(z)) \in \{(4, 4), (4, 6), (6, 4), (6, 6)\}$ . In these cases, it is routine check to prove that  $u$  has at least 6 +6-neighbours, or we obtain a similar situation as in 3.3e, so  $\bar{\mu}(x) \geq 0$ .

Consider the graph of an icosahedron; into each its triangular face  $[XYZ]$  insert a new triangle  $[ABC]$  and add new edges  $\{A, X\}, \{A, Y\}, \{B, Y\}, \{B, Z\}, \{C, Z\}, \{C, X\}$ . In the resulting graph, every 4-path is of weight of at least  $15 + 3 \cdot 4 = 27$ . ■

### Acknowledgement

A support of Slovak VEGA grant 1/7467/20 is acknowledged.

### References

- [1] K. Ando, S. Iwasaki and A. Kaneko, *Every 3-connected planar graph has a connected subgraph with small degree sum*, Annual Meeting of the Mathematical Society of Japan, 1993 (in Japanese).
- [2] O.V. Borodin, *Solution of problems of Kotzig and Grünbaum concerning the isolation of cycles in planar graphs*, Mat. Zametki **46** (5) (1989) 9–12.
- [3] O.V. Borodin, *Minimal vertex degree sum of a 3-path in plane maps*, Discuss. Math. Graph Theory **17** (1997) 279–284.
- [4] O.V. Borodin and D.R. Woodall, *Short cycles of low weight in normal plane maps with minimum degree 5*, Discuss. Math. Graph Theory **18** (1998) 159–164.
- [5] I. Fabrici and S. Jendrol', *Subgraphs with restricted degrees of their vertices in planar 3-connected graphs*, Graphs and Combinatorics **13** (1997) 245–250.

- [6] I. Fabrici and S. Jendrol', *Subgraphs with restricted degrees of their vertices in planar graphs*, Discrete Math. **191** (1998) 83–90.
- [7] I. Fabrici, E. Hexel, S. Jendrol' and H. Walther, *On vertex-degree restricted paths in polyhedral graphs*, Discrete Math. **212** (2000) 61–73.
- [8] P. Franklin, *The four color problem*, Amer. J. Math. **44** (1922) 225–236.
- [9] S. Jendrol' and T. Madaras, *On light subgraphs in plane graphs of minimum degree five*, Discuss. Math. Graph Theory **16** (1996) 207–217.
- [10] E. Jucovič, *Convex polytopes* (Veda, Bratislava, 1981).
- [11] T. Madaras, *Note on weights of paths in polyhedral graphs*, Discrete Math. **203** (1999) 267–269.
- [12] B. Mohar, *Light paths in 4-connected graphs in the plane and other surfaces*, J. Graph Theory **34** (2000) 170–179.
- [13] P. Wernicke, *Über den kartographischen Vierfarbensatz*, Math. Ann. **58** (1904) 413–426.

Received 14 April 1999

Revised 20 August 2000