

## CONNECTIVITY OF PATH GRAPHS

MARTIN KNOR

*Slovak University of Technology*  
*Faculty of Civil Engineering, Department of Mathematics*  
*Radlinského 11, 813 68 Bratislava, Slovakia*

**e-mail:** knor@vox.svf.stuba.sk

AND

L'UDOVÍT NIEPEL

*Kuwait University, Faculty of Science*  
*Department of Mathematics & Computer Science*  
*P.O. box 5969 Safat 13060, Kuwait*

**e-mail:** NIEPEL@MATH-1.sci.kuniv.edu.kw.

### Abstract

We prove a necessary and sufficient condition under which a connected graph has a connected  $P_3$ -path graph. Moreover, an analogous condition for connectivity of the  $P_k$ -path graph of a connected graph which does not contain a cycle of length smaller than  $k+1$  is derived.

**Keywords:** connectivity, path graph, cycle.

**2000 Mathematics Subject Classification:** 05C40, 05C38.

## 1 Introduction

Let  $G$  be a graph,  $k \geq 1$ , and let  $\mathcal{P}_k$  be the set of all paths of length  $k$  (i.e., with  $k+1$  vertices) in  $G$ . The vertex set of a *path graph*  $P_k(G)$  is the set  $\mathcal{P}_k$ . Two vertices of  $P_k(G)$  are joined by an edge if and only if the edges in the intersection of the corresponding paths form a path of length  $k-1$ , and their union forms either a cycle or a path of length  $k+1$ . It means that the

---

Supported by VEGA grant 1/6293/99.

Supported by Kuwait University grant #SM 172.

vertices are adjacent if and only if one can be obtained from the other by "shifting" the corresponding paths in  $G$ .

Path graphs were investigated by Broersma and Hoede in [2] as a natural generalization of line graphs, since  $P_1(G)$  is the line graph of  $G$ . We have to point out that, in the pioneering paper [2] the number  $k$  in  $P_k(G)$  denotes the number of vertices of the paths and not their length. However, in some applications our notation is more consistent, see e.g., [3]. Traversability of  $P_2$ -path graphs is studied in [9], and a characterization of  $P_2$ -path graphs is given in [2] and [7]. Distance properties of path graphs are studied in [1], [4] and [5], and [6] and [8] are devoted to isomorphisms of path graphs.

Let  $V = V(G)$  be a set of  $n$  distinct symbols. Consider strings of length  $k+1$  of these symbols, in which all  $k+1$  symbols are mutually distinct. Let  $G$  be a graph on vertex set  $V$ , edges of which correspond to pairs of symbols which can be neighbours in our strings. If we do not distinguish between a string and its reverse, then  $P_k(G)$  is connected if and only if every string can be obtained from any other one sequentially, by removing a symbol from one of its ends and adding a symbol to the other end.

Let  $G$  be a connected graph. It is well-known (and trivial to prove) that  $P_1(G)$ , i.e., the line graph of  $G$ , is a connected graph. However, this is not the case for  $P_k$ -path graphs if  $k \geq 2$ . This causes some problems, especially when studying distances in path graphs. For example, in [1] the authors give an upper bound for the diameter of every component of a  $P_k$ -path graph, as the whole graph can be disconnected. By [4, Theorem 1], we have:

**Theorem A.** *Let  $G$  be a connected graph. Then  $P_2(G)$  is disconnected if and only if  $G$  contains two distinct paths  $A$  and  $B$  of length two, such that the degrees of both endvertices of  $A$  are 1 in  $G$ .*


In this paper we generalize Theorem A to  $P_k$ -path graphs when  $G$  does not contain a cycle of length smaller than  $k+1$ . Moreover, we completely solve the case of  $P_3$ -path graphs.

We use standard graph-theoretic notation. Let  $G$  be a graph. The vertex set and the edge set of  $G$ , respectively, are denoted by  $V(G)$  and  $E(G)$ . For two subgraphs,  $H_1$  and  $H_2$  of  $G$ , by  $H_1 \cup H_2$  we denote the union of  $H_1$  and  $H_2$ , and  $H_1 \cap H_2$  denotes their intersection. Let  $u$  and  $v$  be vertices in  $G$ . By  $d_G(u, v)$  we denote the distance from  $u$  to  $v$  in  $G$ , and by  $\deg_G(u)$  the degree of  $u$  is denoted. For the vertex set of a component of  $G$  containing  $u$  we use  $Co(u)$ . A path and a cycle, respectively, of length  $l$  are denoted by  $P_l$  and  $C_l$ .

The outline of the paper is as follows. In Section 2 we give a (necessary and sufficient) condition for a connected graph (under some restrictions) to have a connected  $P_k$ -path graph, and Section 3 is devoted to an analogous condition for  $P_3$ -path graphs of general graphs.

## 2 $P_k$ -Path Graphs

Let  $G$  be a graph,  $k \geq 2$ ,  $0 \leq t \leq k-2$ , and let  $A$  be a path of length  $k$  in  $G$ . By  $P_{k,t}^*$  we denote an induced subgraph of  $G$  which is a tree of diameter  $k+t$  with a diametric path  $(x_t, x_{t-1}, \dots, x_1, v_0, v_1, \dots, v_{k-t}, y_1, y_2, \dots, y_t)$ , such that all endvertices of  $P_{k,t}^*$  have distance  $\leq t$  either to  $v_0$  or to  $v_{k-t}$  and the degrees of  $v_1, v_2, \dots, v_{k-t-1}$  are 2 in  $P_{k,t}^*$ . Moreover, no vertex of  $V(P_{k,t}^*) - \{v_1, v_2, \dots, v_{k-t-1}\}$  is joined by an edge to a vertex in  $V(G) - V(P_{k,t}^*)$ . The path  $(v_0, v_1, \dots, v_{k-t})$  is a *base* of  $P_{k,t}^*$ , and we say that  $A$  *lies in*  $P_{k,t}^*$ ,  $A \in P_{k,t}^*$ , if and only if the base of  $P_{k,t}^*$  is a subpath of  $A$ .



In Figure 1 a  $P_{6,3}^*$  is pictured. Note that this graph contains also two  $P_{6,0}^*$  and one  $P_{6,1}^*$ , but it does not contain  $P_{6,2}^*$ . We remark that by thin halfedges are outlined possible edges joining vertices of  $P_{6,3}^*$  to vertices in  $V(G) - V(P_{6,3}^*)$ .

In this section we prove the following theorem.

**Theorem 1.** *Let  $G$  be a connected graph without cycles of length smaller than  $k+1$ . Then  $P_k(G)$  is disconnected if and only if  $G$  contains  $P_{k,t}^*$ ,  $0 \leq t \leq k-2$ , and a path  $A$  of length  $k$  such that  $A \notin P_{k,t}^*$ .*

For easier handling of paths of length  $k$  in  $G$  (i.e., the vertices of  $P_k(G)$ ) we adopt the following convention. We denote the vertices of  $P_k(G)$  (as well as the vertices of  $G$ ) by small letters  $a, b, \dots$ , while the corresponding paths of length  $k$  in  $G$  will be denoted by capital letters  $A, B, \dots$ . It means that if  $A$  is a path of length  $k$  in  $G$  and  $a$  is a vertex in  $P_k(G)$ , then  $a$  must be the vertex corresponding to the path  $A$ .

**Lemma 2.** *Let  $G$  be a connected graph without cycles of length smaller than  $k+1$ . Moreover, let  $A = (x_0, x_1, \dots, x_k)$  be a path of length  $k$  in  $G$  which is not in  $P_{k,t}^*$ ,  $0 \leq t \leq k-2$ . Then for every  $i$ ,  $0 \leq i \leq k$ , there is an  $a_i \in \text{Co}(a)$  such that  $x_i$  is an endvertex of  $A_i$  and the edge of  $A_i$  incident with  $x_i$  lies in  $A$ .*

**Proof.** Observe that if there is a vertex  $a_i \in \text{Co}(a)$  such that  $x_i$  is an endvertex of  $A_i$ , then choosing  $a_i$  with  $d_{P_k(G)}(a, a_i)$  smallest possible, the endedge of  $A_i$  incident with  $x_i$  is in  $A$ .

Thus, suppose that for some  $i$ ,  $0 < i < k$ , there is no  $a_i \in \text{Co}(a)$  such that  $x_i$  is an endvertex of  $A_i$ . Let  $H$  be a subgraph of  $G$  formed by the vertices and edges of paths  $A'$ , where  $a' \in \text{Co}(a)$ . Clearly,  $(x_{i-1}, x_i, x_{i+1}) \subseteq A'$  for every  $a' \in \text{Co}(a)$ . Let  $R = (v_0, v_1, \dots, v_{k-t})$  be the longest path that share all  $A'$ ,  $a' \in \text{Co}(a)$ . As  $k-t \geq 2$ , we have  $t \leq k-2$ . Further,  $\deg_H(v_1) = \deg_H(v_2) = \dots = \deg_H(v_{k-t-1}) = 2$ , and every endvertex of  $H$  has distance  $\leq t$  either to  $v_0$  or to  $v_{k-t}$ . Since  $H$  does not contain cycles (recall that the length of every cycle in  $G$  is at least  $k+1$ ),  $H$  is  $P_{k,t}^*$ ,  $0 \leq t \leq k-2$ . As  $R \subseteq A$  we have  $A \in P_{k,t}^*$ , a contradiction. ■

Let  $A$  and  $B$  be two paths of length  $k$  in  $G$ . If one endvertex of  $B$ , say  $x$ , lies in  $A$ , but the edge of  $B$  incident with  $x$  is not in  $A$ , then we say that the pair  $(A, B)$  forms  $T$  with a touching vertex  $x$ .

Note that if  $(A, B)$  forms  $T$  in  $G$ , then  $A \cup B$  is not necessarily a tree even if  $G$  does not contain a cycle of length  $\leq k$ .

**Lemma 3.** *Let  $G$  be a graph without cycles of length smaller than  $k+1$ . Moreover, suppose  $G$  does not contain  $P_{k,t}^*$ ,  $0 \leq t \leq k-2$ , and let  $(A, B)$  form  $T$  in  $G$ . Then  $b \in \text{Co}(a)$ .*

**Proof.** Let  $(A, B)$  form  $T$  with a touching vertex  $x$ . By Lemma 2, there is  $a' \in \text{Co}(a)$  such that  $x$  is an endvertex of  $A'$  and the edge of  $A'$  incident with  $x$  lies in  $A$ . As  $G$  does not contain a cycle of length smaller than  $k+1$ , we have  $d_{P_k(G)}(a', b) \leq k$ , and hence  $b \in \text{Co}(a)$ . ■

Now we are able to prove Theorem 1.

**Proof of Theorem 1.** We arrange the proof into three steps.

(i) First suppose that  $G$  contains some  $P_{k,t}^*$ ,  $0 \leq t \leq k-2$ , with a base  $R = (v_0, v_1, \dots, v_{k-t})$ , and a path  $A$  of length  $k$  such that  $A \notin P_{k,t}^*$ . Since the diameter of  $P_{k,t}^*$  is  $k+t$ , there is a path  $B$  of length  $k$  in  $G$  such that  $B \in P_{k,t}^*$ , i.e.,  $R \subseteq B$ . By the structure of  $P_{k,t}^*$ , for every vertex  $b'$  of  $P_k(G)$

which is adjacent to  $b$  we have  $R \subseteq B'$ , too. Hence, for every  $b' \in Co(b)$  it holds  $R \subseteq B'$ . Since  $A$  does not contain  $R$ , we have  $a \notin Co(b)$ , so that  $P_k(G)$  is a disconnected graph.

(ii) Now suppose that  $G$  contains some  $P_{k,t}^*$ ,  $0 \leq t \leq k-2$ , such that for every  $a \in V(P_k(G))$  it holds  $A \in P_{k,t}^*$ . We show that either  $P_k(G)$  is a connected graph, or  $G$  contains  $P_{k,t'}^*$ ,  $0 \leq t' < t$ , and a path  $B$  of length  $k$  such that  $B \notin P_{k,t'}^*$ .

Let  $R = (v_0, v_1, \dots, v_{k-t})$  be the base of  $P_{k,t}^*$ , and let  $b$  be a vertex of  $P_k(G)$  such that  $B \in P_{k,t}^*$  and  $v_0$  is an endvertex of  $B$  (e.g., choose  $B$  as a part of a diametric path of  $P_{k,t}^*$ ). Let  $a$  be a vertex of  $P_k(G)$ ,  $A \in P_{k,t}^*$ . If there is  $a' \in Co(a)$  such that either  $v_0$  or  $v_{k-t}$  is an endvertex of  $A'$ , then either  $d_{P_k(G)}(a', b) \leq 2t$  or  $d_{P_k(G)}(a', b) = t$  (by the structure of  $P_{k,t}^*$  we have  $R \subseteq A'$ ). Hence,  $a \in Co(b)$ .

Thus, suppose that there is a vertex  $a$  in  $P_k(G)$ ,  $A \in P_{k,t}^*$ , such that for every  $a' \in Co(a)$  neither  $v_0$  nor  $v_{k-t}$  is an endvertex of  $A'$ . Let  $H$  be a subgraph of  $G$  formed by the vertices and edges of paths  $A'$ , for which  $a' \in Co(a)$ . Clearly,  $R \subseteq A'$  for every  $a' \in Co(a)$ . Let  $R' = (v'_0, v'_1, \dots, v'_{k-t'})$  be the longest path that share all  $A'$ ,  $a' \in Co(a)$ . Since  $R \subset R'$ , by the choice of  $A$  we have  $v_0 = v'_i$ ,  $v_1 = v'_{i+1}$ ,  $\dots$ ,  $v_{k-t} = v'_{i+k-t}$ , where  $i > 0$  and  $i+k-t < k-t'$ , i.e.,  $t' < t - i$ . Further,  $\deg_H(v'_1) = \deg_H(v'_2) = \dots = \deg_H(v'_{k-t-1}) = 2$ , and every endvertex of  $H$  has distance  $\leq t'$  either to  $v'_0$  or to  $v'_{k-t'}$ . Since  $H$  does not contain cycles,  $H$  is  $P_{k,t'}^*$ ,  $0 \leq t' \leq k-2$ . As  $R' \not\subseteq B$ , we have  $B \notin P_{k,t'}^*$ .

(iii) Finally, suppose that  $G$  does not contain  $P_{k,t}^*$ ,  $0 \leq t \leq k-2$ . We show that  $P_k(G)$  is a connected graph.

Let  $a, b \in V(P_k(G))$ . First suppose that  $A \cap B$  does not contain an edge. Let  $P = (y_0, y_1, \dots, y_l)$  be a shortest path in  $G$  joining a vertex of  $A$  with a vertex of  $B$  (i.e.,  $y_l \in V(B)$ ). By Lemma 2, there is  $b' \in Co(b)$  such that  $y_l$  is an endvertex of  $B'$  and the edge of  $B'$  incident with  $y_l$  lies in  $B$ . Let  $B' = (b'_0, b'_1, \dots, b'_{k-1}, y_l)$ . Then  $P' = (b'_0, b'_1, \dots, b'_{k-1}, y_l, y_{l-1}, \dots, y_0)$  is a walk of length  $k + l$ . Since  $G$  does not contain a cycle of length  $\leq k$ , every subwalk of  $P'$  of length  $k$  is a path. Let  $B''$  be a subpath of length  $k$  of  $P'$  with endvertex  $y_0$ . Then  $d_{P_k(G)}(b', b'') \leq l$ , and hence  $b'' \in Co(b)$ . As  $(A, B'')$  forms  $T$  in  $G$ , we have  $b \in Co(a)$ , by Lemma 3.

Now suppose that  $A \cap B$  contains an edge. Let  $P = (y_0, y_1, \dots, y_l)$  be a longest path that is shared by  $A$  and  $B$ . By Lemma 2, for every  $i$ ,  $0 \leq i \leq l$ , there is  $b_i \in Co(b)$  such that  $y_i$  is an endvertex of  $B_i$ , and the edge of  $B_i$  incident with  $y_i$  lies in  $B$ . If  $B_0$  does not contain the edge  $y_0 y_1$ , then  $(A, B_0)$  forms  $T$  in  $G$ , so that  $b \in Co(a)$ , by Lemma 3. Analogously, if

$B_l$  does not contain  $y_{l-1}y_l$ , then  $b \in \text{Co}(a)$ . Thus, suppose that  $B_0$  contains the edge  $y_0y_1$  and  $B_l$  contains  $y_{l-1}y_l$ . Then there is some  $i$ ,  $0 \leq i < l$ , such that both  $B_i$  and  $B_{i+1}$  contain the edge  $y_iy_{i+1}$ . By Lemma 2, there is  $a' \in \text{Co}(a)$  such that  $y_i$  is an endvertex of  $A'$  and the edge of  $A'$  incident with  $y_i$  lies in  $A$ . If  $A'$  contains the edge  $y_iy_{i+1}$ , then  $d_{P_k(G)}(a', b_{i+1}) \leq k-1$ , and hence  $b \in \text{Co}(a)$ . On the other hand, if  $A'$  does not contain  $y_iy_{i+1}$ , we have  $d_{P_k(G)}(a', b_i) \leq k$ , and hence  $b \in \text{Co}(a)$  as well. ■

### 3 $P_3$ -Path Graphs

Let  $G$  be a graph and let  $A$  be a path of length three in  $G$ . By  $P_3^\circ$  we denote a subgraph of  $G$  induced by vertices of a path of length 3, say  $(v_0, v_1, v_2, v_3)$ , such that neither  $v_0$  nor  $v_3$  has a neighbour in  $V(G) - \{v_1, v_2\}$ . We say that the path  $A$  is in  $P_3^\circ$ ,  $A \in P_3^\circ$ , if  $A = (v_0, v_1, v_2, v_3)$ .

By  $P_4^\circ$  we denote an induced subgraph of  $G$  with a path  $(x, v_0, v_1, v_2, y)$ , in which every neighbour of  $v_0$  (and analogously every neighbour of  $v_2$ ), except  $v_0$ ,  $v_1$  and  $v_2$ , has degree 1, or it has degree 2 and in this case it is adjacent to  $v_1$ . Moreover, no vertex of  $V(P_4^\circ) - \{v_1\}$  is joined by an edge to a vertex of  $V(G) - V(P_4^\circ)$  in  $G$ . The path  $(v_0, v_1, v_2)$  is a *base* of  $P_4^\circ$ , and we say that the *path  $A$  lies in  $P_4^\circ$* ,  $A \in P_4^\circ$ , if the base of  $P_4^\circ$  is a subpath of  $A$ .




Figure 2




Figure 3

On example of a graph  $P_3^\circ$  is pictured in Figure 2 and a graph  $P_4^\circ$  in Figure 3. The edges that must be in  $G$  are painted thick, while edges, that are not necessarily in  $G$ , are painted thin.

Let  $K_4$  be a complete graph on 4 vertices, and let  $S$  be a set (possibly empty) of independent vertices. A graph obtained from  $K_4 \cup S$  by joining all vertices of  $S$  to one special vertex of  $K_4$  is denoted by  $K_4^*$ , see Figure 4. Let  $K_{2,t}$  be a complete bipartite graph,  $t \geq 1$ , and let  $(X, Y)$  be the bipartition of  $K_{2,t}$ ,  $X = \{v_1, v_2\}$ . Join  $t$  sets of independent vertices by edges, each to one vertex of  $Y$ ; further, glue a set of stars (each with at least 3 vertices) by one endvertex, each either to  $v_1$  or to  $v_2$ ; glue a set of triangles by one

vertex, each either to  $v_1$  or to  $v_2$ ; and finally, join  $v_1$  to  $v_2$  by an edge. The resulting graph is denoted by  $K_{2,t}^*$ , see Figure 5.




Figure 4




Figure 5

**Theorem 4.** *Let  $G$  be a connected graph such that  $P_3(G)$  is not empty. Then  $P_3(G)$  is disconnected if and only if one of the following holds:*

- (1)  $G$  contains  $P_t^\circ$ ,  $t \in \{3, 4\}$ , and a path  $A$  of length 3 such that  $A \notin P_t^\circ$ ;
- (2)  $G$  is isomorphic to  $K_4^*$ ;
- (3)  $G$  is isomorphic to  $K_{2,t}^*$ ,  $t \geq 1$ .

If  $A \in P_3^\circ$  in  $G$ , then  $a$  is an isolated vertex in  $P_3(G)$ , and if  $A \in P_4^\circ$ , then  $a$  lies in a complete bipartite graph. Thus, we have the following corollary of Theorem 4.

**Corollary 5.** *Let  $G$  be a connected graph that is not isomorphic to  $K_4^*$  or to  $K_{2,t}^*$ ,  $t \geq 1$ . Then at most one nontrivial component of  $P_3(G)$  is different from a complete bipartite graph.*

In the proof of Theorem 4 we use 6 lemmas.

**Lemma 6.** *Let  $G$  be a connected graph, and let  $a$  and  $b$  be vertices in  $P_3(G)$ . If neither  $A$  nor  $B$  is in some  $P_3^\circ$  or  $P_4^\circ$  in  $G$ , then there are vertices  $c$  and  $d$  in  $P_3(G)$ , such that  $c \in \text{Co}(a)$ ,  $d \in \text{Co}(b)$  and  $C$  and  $D$  share an edge in  $G$ .*

**Proof.** Let  $A \cap B$  do not contain an edge, and let  $P = (y_0, y_1, \dots, y_l)$  be a shortest path in  $G$  joining a vertex of  $A$  with a vertex of  $B$  (i.e.,  $y_l \in V(B)$ ). We show that there is a vertex  $b'$  in  $\text{Co}(b)$ , such that  $y_l$  is an endvertex of  $B'$ .

Suppose that there is no vertex  $b'$  with the required property. Then  $B = (x_0, x_1, y_l, x_3)$ , and since  $B$  is not in  $P_3^\circ$  in  $G$ , there is a vertex  $\bar{b}$  in  $P_3(G)$  such that  $\bar{b}b \in E(P_3(G))$ . By our assumption,  $\bar{B} = (x_1, y_l, x_3, x_4)$  for some  $x_4 \in V(G)$ . Moreover, for every neighbour  $u$  of  $b$  we have  $U = (x_1, y_l, x_3, z)$ ,

where  $z$  has no neighbours in  $V(G) - \{y_l, x_3\}$ ; and for every neighbour  $v$  of  $\bar{b}$  we have  $V = (z, x_1, y_l, x_3)$ , where  $z$  has no neighbours in  $V(G) - \{x_1, y_l\}$ . Hence  $B$  is in some  $P_4^\circ$ , a contradiction.

Thus, there is a vertex  $b' \in Co(b)$ , such that  $y_l$  is an endvertex of  $B'$ . Let  $b''$  be the first vertex on a shortest  $b - b'$  path in  $P_3(G)$ , such that one endvertex of  $B''$  is in  $P$ . Assume that  $B'' = (b_3'', b_2'', b_1'', y_i)$ . Then  $P' = (b_3'', b_2'', b_1'', y_i, y_{i-1}, \dots, y_0)$  is a path of length  $i+3 \geq 3$ . Let  $B^*$  be a subpath of  $P$  of length 3, such that  $y_0$  is an endvertex of  $B^*$ . Then  $d_{P_3(G)}(b'', b^*) = i$ , and hence,  $b^* \in Co(b)$ .

Denote  $B^* = (y_0, b_1^*, b_2^*, b_3^*)$ , and suppose that  $A \cap B^*$  does not contain an edge. Let  $A = (a_0, a_1, a_2, a_3)$ . Distinguish two cases.

- (i)  $y_0 = a_1$ . Then  $b_1^* \neq a_0$  and  $b_1^* \neq a_2$ , so that at least one of  $a_0$  and  $a_2$ , say  $a_0$ , is different from  $b_2^*$ . Since  $a_0$  is not an interior vertex of  $B^*$ ,  $D = (a_0, y_0, b_1^*, b_2^*)$  is a path of length 3 in  $G$ . As  $b^*d \in E(P_3(G))$ , we have  $d \in Co(b)$  and  $A \cap D$  contains an edge.
- (ii)  $y_0 = a_0$ . If  $b_1^* \neq a_2$  then  $C = (b_1^*, y_0, a_1, a_2)$  is a path of length 3 in  $G$ ,  $c \in Co(a)$ ,  $b^* \in Co(b)$ , and  $C \cap B^*$  contains an edge. On the other hand, if  $b_1^* = a_2$  then  $D = (a_1, y_0, a_2, b_2^*)$  is a path of length 3 in  $G$ ,  $d \in Co(b)$ , and  $A \cap D$  contains an edge. ■

**Lemma 7.** *Let  $G$  be a connected graph, and let  $a$  and  $b$  be two vertices in  $P_3(G)$  such that  $b \notin Co(a)$  and  $A \cap B$  contains a path of length two. Moreover, suppose  $G$  does not contain  $P_3^\circ$  or  $P_4^\circ$ . Then  $G$  is isomorphic either to  $K_4^*$  or to  $K_{2,t}^*$  for some  $t \geq 1$ .*

**Proof.** Let  $A = (x_0, x_1, x_2, x_3)$  and  $B = (x_0, x_1, x_2, x_4)$ ,  $x_3 \neq x_4$ . Since  $b \notin Co(a)$ ,  $x_0$  has no neighbour in  $V(G) - \{x_1, x_2\}$ . Thus, both  $x_3$  and  $x_4$  have some neighbours in  $V(G) - \{x_1, x_2\}$ , as  $G$  does not contain  $P_3^\circ$ . Let  $y$  be a vertex of  $G$  such that  $x_1y \in E(G)$  and  $y \notin \{x_0, x_2, x_3, x_4\}$ . Then  $a' \in Co(a)$  and  $b' \in Co(b)$ , where  $A' = (y, x_1, x_2, x_3)$  and  $B' = (y, x_1, x_2, x_4)$ . Since  $b \notin Co(a)$  we have  $b' \notin Co(a')$ , and hence,  $y$  has no neighbour in  $V(G) - \{x_1, x_2\}$ .

Suppose that  $x_3x_4 \in E(G)$  and distinguish three cases.

*Case 1.*  $x_1x_3, x_1x_4 \in E(G)$ , see Figure 6.

Let  $G'$  be a graph obtained from  $G$  by joining  $x_0$  to  $x_2$ . Then  $A, (x_1, x_2, x_3, x_4), (x_2, x_3, x_4, x_1), (x_3, x_4, x_1, x_0), (x_4, x_1, x_0, x_2), (x_1, x_0, x_2, x_4), (x_0, x_2, x_4, x_3), (x_2, x_4, x_3, x_1), (x_1, x_2, x_4, x_3), B$  is a sequence of paths



whose images produce a walk of length 9 from  $a$  to  $b$  in  $P_3(G')$ . (We remark that  $d_{P_3(G')}(a, b) = 9$ .) Thus  $b \in Co(a)$ , a contradiction. Hence  $\deg_G(x_0) = 1$ .

Let  $C_1 = (x_1, x_2, x_3, x_4)$  and  $C_2 = (x_1, x_2, x_4, x_3)$  be two cycles of length 4 in  $G$ . For every subpath  $A'$  of  $C_1$  of length 3 we have  $a' \in Co(a)$ , and for every subpath  $B'$  of  $C_2$  of length 3 we have  $b' \in Co(b)$ . Let  $y$  be a vertex in  $V(G) - \{x_1, \dots, x_4\}$  which is joined to some  $x \in \{x_1, \dots, x_4\}$ . Since  $C_1 \cap C_2$  contains an edge incident with  $x$ , there are paths  $A''$  and  $B''$  of length 3 in  $G$ , both containing the edge  $yx$ , such that  $a'' \in Co(a)$ ,  $b'' \in Co(b)$  and  $A'' \cap B''$  contains  $P_2$ . Thus, analogously as above it can be shown that  $\deg_G(y) = 1$ . Finally, as  $G$  does not contain  $P_3^\circ$  we have  $x = x_1$ , and hence  $G \cong K_4^*$ .




Figure 6




Figure 7




Figure 8

*Case 2.*  $x_1x_3 \in E(G)$  and  $x_1x_4 \notin E(G)$ , see Figure 7 and Figure 8 (by dotted lines edges that are missing in  $G$  are pictured).

Since  $(x_1, x_2, x_3)$  is not a base of  $P_4^\circ$ , either there is a vertex  $y \in V(G) - \{x_0, \dots, x_4\}$  such that  $yx_4 \in E(G)$ , or there is a path of length 2 glued by one endvertex to  $x_3$  (the other vertices of the path are not in  $\{x_0, \dots, x_4\}$ ).

First suppose that there is  $x_5 \in V(G) - \{x_0, \dots, x_4\}$  such that  $x_4x_5 \in E(G)$ , see Figure 7. Let  $G'$  be a graph obtained from  $G$  by joining  $x_0$  to  $x_2$ . Then  $A, (x_1, x_2, x_3, x_4), (x_2, x_3, x_4, x_5), (x_0, x_2, x_3, x_4), (x_1, x_0, x_2, x_3), (x_3, x_1, x_0, x_2), (x_4, x_3, x_1, x_0), (x_2, x_4, x_3, x_1), (x_1, x_2, x_4, x_3), B$  is a sequence of paths whose images produce a walk of length 9 from  $a$  to  $b$  in  $P_3(G')$ . Thus  $b \in Co(a)$ , a contradiction.

Hence  $\deg_G(x_0) = 1$ . Analogously, for every vertex  $x$ , such that  $xx_2, xx_3 \in E(G)$ , every neighbour of  $x$  (different from  $x_2$  and  $x_3$ ) has degree 1 in  $G$ .

Let  $y_1$  and  $y_2$  be vertices in  $V(G) - \{x_0, \dots, x_5\}$ , such that  $x_2y_1, y_1y_2 \in E(G)$ . If  $y_2$  is joined by an edge to a vertex, say  $z$ , of  $V(G) - \{x_2, y_1\}$ , then for  $C = (x_2, y_1, y_2, z)$  we have  $c \in Co(a)$  and  $c \in Co(b)$ . Hence  $b \in Co(a)$ , a contradiction. Since  $G$  contains  $P_3^\circ$  if there is a vertex of degree 1 joined to  $x_2$ , we have  $G \cong K_{2,t}^*$  for some  $t \geq 2$ .

Now suppose that there are  $x_5, x_6 \in V(G) - \{x_0, \dots, x_4\}$  such that  $x_3x_5, x_5x_6 \in E(G)$ , see Figure 8. (Observe that the cases  $x_6 \in \{x_0, x_1, x_4\}$  imply  $b \in Co(a)$ .)

Let  $G'$  be a graph obtained from  $G$  by joining  $x_0$  to  $x_2$ . Then  $A, (x_1, x_2, x_3, x_5), (x_2, x_3, x_5, x_6), (x_0, x_2, x_3, x_5), (x_1, x_0, x_2, x_3), (x_3, x_1, x_0, x_2), (x_4, x_3, x_1, x_0), (x_2, x_4, x_3, x_1), (x_1, x_2, x_4, x_3), B$  is a sequence of paths whose images produce a walk of length 9 from  $a$  to  $b$  in  $P_3(G')$ . Thus  $b \in Co(a)$ , a contradiction.

Hence  $\deg_G(x_0) = 1$ . Analogously, for every vertex  $x$ , such that  $xx_2, xx_3 \in E(G)$ , every neighbour of  $x$  (different from  $x_2$  and  $x_3$ ) has degree 1 in  $G$ . Now analogously as above it can be shown that  $G \cong K_{2,t}^*$  for some  $t \geq 2$ .

*Case 3.*  $x_1x_3, x_1x_4 \notin E(G)$ , see Figure 9.

Since neither  $(x_1, x_2, x_3)$  nor  $(x_1, x_2, x_4)$  is a base of  $P_4^\circ$ , there is a vertex  $x_5 \in V(G) - \{x_0, \dots, x_4\}$  which is adjacent either to  $x_3$  or to  $x_4$ . Assume that  $x_3x_5 \in E(G)$ . As  $b \notin Co(a)$ ,  $x_5$  has no neighbour in  $\{x_0, x_1, x_4\}$ . Since  $(x_1, x_2, x_3)$  is not a base of  $P_4^\circ$ , there is a vertex  $y \in V(G) - \{x_0, \dots, x_5\}$  such that either  $yx_5 \in E(G)$  or  $yx_4 \in E(G)$ .

First suppose that there is a vertex  $x_6 \in V(G) - \{x_0, \dots, x_5\}$  such that  $x_5x_6 \in E(G)$ . Then every neighbour of  $x_4$  (different from  $x_2$  and  $x_3$ ) has degree 1 in  $G$ , otherwise  $b \in Co(a)$ . Analogously, for every vertex  $x$ , such that  $xx_2, xx_3 \in E(G)$ , every neighbour of  $x$  (different from  $x_2$  and  $x_3$ ) has degree 1 in  $G$ . Thus, analogously as above we have  $G \cong K_{2,t}^*$  for some  $t \geq 1$ .

If there is  $x_6 \in V(G) - \{x_0, \dots, x_5\}$  such that  $x_4x_6 \in E(G)$ , then the problem is reduced to the previous case as  $(x_3, x_2, x_4)$  is not a base of  $P_4^\circ$ .




Figure 9




Figure 10

To prove the lemma it remains to consider the case  $x_3x_4 \notin E(G)$ , see Figure 10.

As  $b \notin Co(a)$ , there is no cycle  $(x_3, x_2, x_4, \dots)$  of length at least 4 in  $G$ . Since neither  $A$  nor  $B$  is in  $P_3^\circ$  in  $G$ , there are  $x_5, x_6 \in V(G) - \{x_0, \dots, x_5\}$ ,  $x_5 \neq x_6$ , such that  $x_3x_5, x_4x_6 \in E(G)$ . Moreover, as  $G$  does not contain

$P_4^\circ$  with base  $(x_1, x_2, x_3)$ , there is  $x_7 \in V(G) - \{x_0, \dots, x_6\}$  such that  $x_5x_7 \in E(G)$ , and analogously, there is  $x_8 \in V(G) - \{x_0, \dots, x_7\}$  such that  $x_6x_8 \in E(G)$ . (Observe that  $b \in \text{Co}(a)$  if  $x_7 = x_1$ , and the same holds if  $x_8 = x_1$ .) But now  $d_{P_3(G)}(a, b) \leq 7$ , and hence  $b \in \text{Co}(a)$ , a contradiction. ■

**Lemma 8.** *Let  $G$  be a connected graph, and let  $a$  and  $b$  be two vertices in  $P_3(G)$  such that  $b \notin \text{Co}(a)$  and  $A \cap B$  contains two independent edges. Moreover, suppose  $G$  does not contain  $P_3^\circ$  or  $P_4^\circ$ . Then  $G$  is isomorphic either to  $K_4^*$  or to  $K_{2,t}^*$  for some  $t \geq 1$ , or there are  $a' \in \text{Co}(a)$  and  $b' \in \text{Co}(b)$  such that  $A' \cap B'$  contains a path of length 2.*

**Proof.** Let  $A = (x_0, x_1, x_2, x_3)$ . Since  $b \notin \text{Co}(a)$ ,  $B = (x_0, x_1, x_3, x_2)$ . We may assume that  $x_0$  has no neighbour in  $V(G) - \{x_0, \dots, x_3\}$ , otherwise there are  $a' \in \text{Co}(a)$  and  $b' \in \text{Co}(b)$  such that  $A' \cap B'$  contains  $P_2$ .

Distinguish three cases.

*Case 1.*  $x_0x_2, x_0x_3 \in E(G)$ . Then both  $A$  and  $B$  lie in cycles of length 4. If there is a vertex  $y$  adjacent to a vertex of  $\{x_0, \dots, x_4\}$ , then there are  $a' \in \text{Co}(a)$  and  $b' \in \text{Co}(b)$  such that  $A' \cap B'$  contains  $P_2$ . Thus,  $G \cong K_4$  which is a special  $K_4^*$ .

*Case 2.*  $x_0x_2 \in E(G)$  and  $x_0x_3 \notin E(G)$ , see Figure 11. Since  $A$  is not in  $P_3^\circ$  in  $G$ , there is a vertex  $x_4 \in V(G) - \{x_0, \dots, x_3\}$  such that  $x_3x_4 \in E(G)$ . But then  $a' \in \text{Co}(a)$ ,  $b' \in \text{Co}(b)$  and  $A' \cap B'$  contains  $P_2$ , where  $A' = (x_1, x_2, x_3, x_4)$  and  $B' = (x_0, x_2, x_3, x_4)$ .

*Case 3.*  $x_0x_2, x_0x_3 \notin E(G)$ , see Figure 12. Since neither  $A$  nor  $B$  is in  $P_3^\circ$  in  $G$ , there are vertices  $x_4, x_5 \in V(G) - \{x_0, \dots, x_3\}$  such that  $x_2x_4, x_3x_5 \in E(G)$ . We may assume that the degree of every neighbour of  $x_1$  (except  $x_2$  and  $x_3$ ) is 1 in  $G$ , as the other possibilities we have already solved.

If  $x_4 \neq x_5$ , then there are  $x_6, x_7 \in V(G) - \{x_0, \dots, x_3\}$  such that  $x_4x_6, x_5x_7 \in E(G)$ , as neither  $(x_1, x_3, x_2)$  nor  $(x_1, x_2, x_3)$  is a base of  $P_4^\circ$ . But then  $b \in \text{Co}(a)$ , a contradiction.

Thus, suppose that  $x_4 = x_5$ . By previous subcase, we may assume that  $\deg_G(x_2) = \deg_G(x_3) = 3$ . As  $(x_1, x_2, x_3)$  is not a base of  $P_4^\circ$ , there is  $x_5 \in V(G) - \{x_0, \dots, x_4\}$  such that  $x_4x_5 \in E(G)$ . By our assumptions,  $\deg_G(x_5) = 1$ . Hence,  $\deg_G(x_0) = \deg_G(x_5) = 1$ ,  $\deg_G(x_2) = \deg_G(x_3) = 3$ ,

and all neighbours of  $x_1$  and  $x_4$  (except  $x_2$  and  $x_3$ ) have degree 1 in  $G$ . Thus,  $G \cong K_{2,2}^*$ .  $\blacksquare$




Figure 11




Figure 12

**Lemma 9.** *Let  $G$  be a connected graph, and let  $a$  and  $b$  be two vertices in  $P_3(G)$  such that  $b \notin \text{Co}(a)$  and  $A \cap B$  contains exactly one edge and two vertices outside this edge. Moreover, suppose  $G$  does not contain  $P_3^\circ$  or  $P_4^\circ$ . Then there are  $a' \in \text{Co}(a)$  and  $b' \in \text{Co}(b)$  such that  $A' \cap B'$  contains two independent edges.*

**Proof.** Let  $A = (x_0, x_1, x_2, x_3)$ . Then either  $B = (x_0, x_2, x_1, x_3)$  or  $B = (x_1, x_2, x_0, x_3)$ .

First suppose that  $B = (x_0, x_2, x_1, x_3)$ . Since  $A$  is not in  $P_3^\circ$  in  $G$ , either  $x_0x_3 \in E(G)$  or  $x_3x_4 \in E(G)$  for some  $x_4 \in V(G) - \{x_0, \dots, x_3\}$ . In both these cases there are  $a' \in \text{Co}(a)$  and  $b' \in \text{Co}(b)$  such that  $A' \cap B'$  contains two independent edges.

Now suppose that  $B = (x_1, x_2, x_0, x_3)$ . Then for  $A' = (x_1, x_2, x_3, x_0)$  we have  $a' \in \text{Co}(a)$ , and  $A' \cap B$  contains two independent edges.  $\blacksquare$

**Lemma 10.** *Let  $G$  be a connected graph, and let  $a$  and  $b$  be two vertices in  $P_3(G)$  such that  $b \notin \text{Co}(a)$  and  $A \cap B$  contains exactly one edge and one vertex outside this edge. Moreover, suppose  $G$  does not contain  $P_3^\circ$  or  $P_4^\circ$ . Then there are  $a' \in \text{Co}(a)$  and  $b' \in \text{Co}(b)$  such that  $A' \cap B'$  contains two edges.*

**Proof.** Let  $A = (x_0, x_1, x_2, x_3)$ , and let  $x_4$  be a vertex of  $B$  lying outside  $A$ . Distinguish four cases.

*Case 1.* Suppose that  $x_1x_2$  is the middle edge of  $B$ . Then  $B = (x_3, x_1, x_2, x_4)$ . If  $x_4$  has a neighbour in  $V(G) - \{x_1, x_2\}$ , then for  $B' = (x_0, x_1, x_2, x_4)$  we have  $b' \in \text{Co}(b)$  and  $A \cap B' = P_2$ . Thus, we may assume that both  $x_0$  and  $x_4$  have no neighbour in  $V(G) - \{x_1, x_2\}$ . However, then there is some  $P_3^\circ$  in  $G$ , a contradiction.

*Case 2.* Suppose that  $x_1x_2$  is an endedge of  $B$ .

If  $B = (x_1, x_2, x_0, x_4)$ , then for  $A' = (x_4, x_0, x_1, x_2)$  we have  $a' \in Co(a)$  and  $A' \cap B$  contains two independent edges.

If  $B = (x_1, x_2, x_4, x_0)$  then  $b \in Co(a)$ ; and if  $B = (x_1, x_2, x_4, x_3)$ , then for  $B' = (x_0, x_1, x_2, x_4)$  we have  $b' \in Co(b)$  and  $A \cap B' = P_2$ .




Figure 13




Figure 14

*Case 3.* Suppose that  $x_0x_1$  is an endedge of  $B$  and  $x_1$  is an endvertex of  $B$ . If  $B = (x_1, x_0, x_4, x_2)$ ,  $B = (x_1, x_0, x_3, x_4)$ , or  $B = (x_1, x_0, x_4, x_3)$ , then  $b \in Co(a)$ . Thus, suppose that  $B = (x_1, x_0, x_2, x_4)$ , see Figure 13.

If  $\deg_G(x_1) > 2$ , then for  $B' = (x_1, x_0, x_2, x_3)$  we have  $b' \in Co(b)$  and  $A \cap B'$  contains two independent edges. Thus, suppose that  $\deg_G(x_0) = \deg_G(x_1) = 2$ .

If  $x_3x_4 \in E(G)$ , then analogously as above we have  $\deg_G(x_3) = \deg_G(x_4) = 2$ , and hence, there is  $P_4^\circ$  with base  $(x_0, x_2, x_3)$  in  $G$ , a contradiction. Thus, suppose that  $x_3x_4 \notin E(G)$ .

As  $b \notin Co(a)$ , there is no cycle  $(x_3, x_2, x_4, \dots)$  of length at least 4 in  $G$ . Since neither  $A$  nor  $B$  is in  $P_3^\circ$  in  $G$ , there are  $x_5, x_6 \in V(G) - \{x_0, \dots, x_4\}$ ,  $x_5 \neq x_6$ , such that  $x_3x_5, x_4x_6 \in E(G)$ . Moreover, as  $G$  does not contain  $P_4^\circ$  with base  $(x_0, x_2, x_3)$ , there is a vertex  $x_7 \in V(G) - \{x_0, \dots, x_6\}$  such that  $x_5x_7 \in E(G)$ . Thus, for  $A' = (x_6, x_4, x_2, x_3)$  and  $B' = (x_0, x_2, x_4, x_6)$  we have  $a' \in Co(a)$ ,  $b' \in Co(b)$  and  $A' \cap B' = P_2$ .

*Case 4.* Suppose that  $x_0x_1$  is an endedge of  $B$  and  $x_0$  is an endvertex of  $B$ .

If  $B = (x_0, x_1, x_4, x_3)$ , then  $b \in Co(a)$ . Since the cases  $B = (x_0, x_1, x_4, x_2)$  and  $B = (x_0, x_1, x_3, x_4)$  are equivalent, suppose that  $B = (x_0, x_1, x_4, x_2)$ , see Figure 14.

We have  $x_0x_3 \notin E(G)$ , otherwise  $b \in Co(a)$ . Since  $A$  is not in  $P_3^\circ$  in  $G$ , there is  $y \in V(G) - \{x_0, \dots, x_3\}$  such that either  $x_0y \in E(G)$  or  $x_3y \in E(G)$ . Assume that  $x_0y \in E(G)$ . If  $y \neq x_4$ , then for  $A' = (y, x_0, x_1, x_2)$  and  $B' = (y, x_0, x_1, x_4)$  we have  $a' \in Co(a)$ ,  $b' \in Co(b)$  and  $A' \cap B' = P_2$ . On the other hand, if  $y = x_4$ , then for  $A' = (x_2, x_4, x_0, x_1)$  we have  $a' \in Co(a)$  and

$A' \cap B$  contains two independent edges. ■

**Lemma 11.** *Let  $G$  be a connected graph, and let  $a$  and  $b$  be two vertices in  $P_3(G)$  such that  $b \notin \text{Co}(a)$  and  $A \cap B$  contains exactly one edge and no vertex outside this edge. Moreover, suppose  $G$  does not contain  $P_3^\circ$  or  $P_4^\circ$ . Then there are  $a' \in \text{Co}(a)$  and  $b' \in \text{Co}(b)$  such that  $A' \cap B'$  contains two edges.*

**Proof.** Let  $A = (x_0, x_1, x_2, x_3)$ , and let  $x_4$  and  $x_5$  be vertices of  $B$  lying outside  $A$ . If  $A' \cap B'$  does not contain  $P_2$  for every  $a' \in \text{Co}(a)$  and  $b' \in \text{Co}(b)$ , then either  $B = (x_0, x_1, x_4, x_5)$  or  $B = (x_4, x_1, x_2, x_5)$ .

First suppose that  $B = (x_0, x_1, x_4, x_5)$ , see Figure 15. If there is  $y \in V(G) - \{x_1, x_2\}$  such that  $yx_3 \in E(G)$ , then for  $A' = (x_5, x_4, x_1, x_2)$  we have  $a' \in \text{Co}(a)$  and  $A' \cap B = P_2$ . Hence, we may assume that  $x_3$  has no neighbour in  $V(G) - \{x_1, x_2\}$ . Since  $A$  is not in  $P_3^\circ$  in  $G$ , there is  $y \in V(G) - \{x_1, x_2\}$  such that  $yx_0 \in E(G)$ . If  $y \neq x_4$ , then for  $A' = (y, x_0, x_1, x_2)$  and  $B' = (y, x_0, x_1, x_4)$  we have  $a' \in \text{Co}(a)$ ,  $b' \in \text{Co}(b)$  and  $A' \cap B' = P_2$ . On the other hand, if  $x_0x_4 \in E(G)$ , then for  $A' = (x_5, x_4, x_0, x_1)$  we have  $a' \in \text{Co}(a)$  and  $A' \cap B$  contains two edges.

Thus, suppose that  $B = (x_4, x_1, x_2, x_5)$ . Since  $A$  is not in  $P_3^\circ$  in  $G$ , we may assume that there is  $y \in V(G) - \{x_1, x_2\}$  such that  $x_0y \in E(G)$ . Then for  $A' = (x_0, x_1, x_2, x_5)$  we have  $a' \in \text{Co}(a)$  and  $A' \cap B = P_2$ . ■




Figure 15

Now we prove Theorem 4.

**Proof of Theorem 4.** First suppose that  $G$  contains  $P_3^\circ$  and a path  $A$  of length 3 such that  $A \notin P_3^\circ$ . Then there is a path  $B$  of length 3 in  $G$  such that  $B \in P_3^\circ$ . Since  $b$  is an isolated vertex in  $P_3(G)$ ,  $b \notin \text{Co}(a)$ . Now suppose that  $G$  contains  $P_4^\circ$ , and choose  $B \in P_4^\circ$ . For every vertex  $b' \in \text{Co}(b)$ ,  $B'$  contains the base of  $P_4^\circ$ . Hence,  $P_3(G)$  is disconnected if there is a path  $A$  of length 3 such that  $A \notin P_4^\circ$ .

If  $G$  is isomorphic to  $K_4^*$ , then  $P_3(G)$  has three components, each containing  $C_4$ . Finally, if  $G$  is isomorphic to  $K_{2,t}^*$ ,  $t \geq 1$ , and  $P_3(G)$  is not empty, then some paths of length 3 in  $G$  contain the edge  $v_1v_2$ , while the

other do not, see Figure 5. Let  $a \in V(P_3(G))$  such that  $v_1v_2 \in A$ . Then  $v_1v_2 \in A'$  for every  $a' \in Co(a)$ , so that  $P_3(G)$  is a disconnected graph.

To prove the "only if" part of Theorem 4, first suppose that  $G$  contains  $P_t^\circ$ ,  $t \in \{3, 4\}$ , but no path  $A$  of length 3 such that  $A \not\subseteq P_t^\circ$ . If  $G$  contains  $P_3^\circ$ , then our assumption implies that  $G$  is a path of length 3. On the other hand, if  $G$  contains  $P_4^\circ$  and there is no  $P_3^\circ$  in  $G$ , then  $G$  is a tree of diameter 4 and  $P_3(G)$  is a complete bipartite graph. Thus, in what follows we restrict our considerations to graphs which do not contain  $P_t^\circ$ ,  $t \in \{3, 4\}$ .

Let  $G$  be a graph which does not contain  $P_3^\circ$  or  $P_4^\circ$ , and let  $a$  and  $b$  be vertices of  $P_3(G)$  such that  $b \notin Co(a)$ . By Lemma 6, there are  $a' \in Co(a)$  and  $b' \in Co(b)$  such that  $A' \cap B'$  contains an edge. Hence,  $G$  is either isomorphic to  $K_4^*$  or to  $K_{2,t}^*$ ,  $t \geq 1$ , by Lemmas 7, 8, 9, 10 and 11. ■

### Acknowledgement

The authors sincerely acknowledge the helpful remarks and corrections of the referee.

### References

- [1] A. Belan and P. Jurica, *Diameter in path graphs*, Acta Math. Univ. Comenian. **LXVIII** (1999) 111–126.
- [2] H.J. Broersma and C. Hoede, *Path graphs*, J. Graph Theory **13** (1989) 427–444.
- [3] M. Knor and L. Niepel, *Path, trail and walk graphs*, Acta Math. Univ. Comenian. **LXVIII** (1999) 253–256.
- [4] M. Knor and L. Niepel, *Distances in iterated path graphs*, Discrete Math. (to appear).
- [5] M. Knor and L. Niepel, *Centers in path graphs*, (submitted).
- [6] M. Knor and L. Niepel, *Graphs isomorphic to their path graphs*, (submitted).
- [7] H. Li and Y. Lin, *On the characterization of path graphs*, J. Graph Theory **17** (1993) 463–466.
- [8] X. Li and B. Zhao, *Isomorphisms of  $P_4$ -graphs*, Australasian J. Combin. **15** (1997) 135–143.
- [9] X. Yu, *Trees and unicyclic graphs with Hamiltonian path graphs*, J. Graph Theory **14** (1990) 705–708.

Received 20 July 1999

Revised 20 March 2000