

## A NOTE ON PERIODICITY OF THE 2-DISTANCE OPERATOR

BOHDAN ZELINKA

*Department of Applied Mathematics*  
*Technical University of Liberec*  
*Liberec, Czech Republic*

**To the memory of Ivan Havel**

### Abstract

The paper solves one problem by E. Prisner concerning the 2-distance operator  $T_2$ . This is an operator on the class  $C_f$  of all finite undirected graphs. If  $G$  is a graph from  $C_f$ , then  $T_2(G)$  is the graph with the same vertex set as  $G$  in which two vertices are adjacent if and only if their distance in  $G$  is 2. E. Prisner asks whether the periodicity  $\geq 3$  is possible for  $T_2$ . In this paper an affirmative answer is given. A result concerning the periodicity 2 is added.

**Keywords:** 2-distance operator, complement of a graph.

**2000 Mathematics Subject Classification:** 05C12.

In this paper we consider finite undirected graphs without loops and multiple edges. The vertex set of a graph  $G$  is denoted by  $V(G)$ , its edge set by  $E(G)$ . The symbol  $\overline{G}$  denotes the complement of  $G$ , i.e., the graph with the same vertex set as  $G$  in which two distinct vertices are adjacent if and only if they are not adjacent in  $G$ .

Let  $\phi$  be a graph operator defined on the class  $C_f$  of all finite undirected graphs. For every positive integer  $r$  we define the power  $\phi^r$  so that  $\phi^1 = \phi$  and for  $r \geq 2$  the operator  $\phi^r$  is such that  $\phi^r(G) = \phi(\phi^{r-1}(G))$  for each  $G \in C_f$ . A graph  $G \in C_f$  is called  $\phi$ -periodic, if there exists a positive integer  $r$  such that  $\phi^r(G) \cong G$ . The minimum number  $r$  with this property is the *periodicity* of the graph  $G$  in the operator  $\phi$ .

For an integer  $k \geq 2$  the operator  $T_k$  on  $C_f$  is defined in such a way that for any graph  $G \in C_f$  the graph  $T_k(G)$  has the same vertex set as  $G$

and two distinct vertices are adjacent in  $T_k(G)$  if and only if their distance in  $G$  is  $k$ . The operator  $T_k$  is called the  $k$ -distance operator.

In [2], page 170, E. Prisner asks the following problem:

Is period  $\geq 3$  possible for  $T_2$  ?

An affirmative answer is given by the following theorem.

**Theorem.** *Let  $r$  be an even positive integer. Then there exists a graph  $G_r$  whose periodicity in the operator  $T_2$  is  $r$ .*

**Proof.** Let  $q = 2^r + 1$ . Let  $V_0, V_1, \dots, V_{q-1}$  be pairwise disjoint sets of vertices. Let  $t$  be an integer,  $t \geq 2$  and let  $|V_i| = t^i$  for  $i = 0, 1, \dots, q-1$ . The vertex set of  $G_r$  is  $V(G_r) = \bigcup_{i=0}^{q-1} V_i$ . All sets  $V_0, V_1, \dots, V_{q-1}$  are independent in  $G_r$ . Let  $x \in V_i, y \in V_j$  for some  $i$  and  $j$  from  $\{0, 1, \dots, q-1\}$ . These vertices are adjacent in  $G_r$  if and only if  $j \equiv i+1 \pmod{q}$  or  $j \equiv i-1 \pmod{q}$ . This implies that all sets  $V_0, V_1, \dots, V_{q-1}$  induce complete subgraphs in the graph  $T_2(G_r)$ . If  $x \in V_i, y \in V_j$ , then  $x, y$  are adjacent in  $T_2(G_r)$  if and only if  $j \equiv i+2 \pmod{q}$  or  $j \equiv i-2 \pmod{q}$ . From these facts by induction we obtain that  $T_2^m(G)$  for  $m \geq 2$  has the following structure. If  $m$  is even, then all sets  $V_0, V_1, \dots, V_{q-1}$  are independent; if  $m$  is odd, then they induce complete subgraphs; if  $x \in V_i, y \in V_j$ , then  $x, y$  are adjacent if and only if  $j \equiv i+2^m \pmod{q}$  or  $j \equiv i-2^m \pmod{q}$  in both the cases. This implies that  $T_2^r(G_r) \cong G_r$ . Now it remains to show that  $T_2^m(G)$  is not isomorphic to  $G_r$  for  $1 \leq m < r$ . We do it using the independence number  $\alpha(G)$ . The greatest independent set in  $G_r$  is  $\bigcup_{i=1}^{\frac{1}{2}(q-1)} V_{2i}$  and thus  $\alpha(G_r) = \sum_{i=1}^{\frac{1}{2}(q-1)} t^{2i} = t^2(t^{q-1} - 1)/(t^2 - 1)$ . If  $m$  is odd, then  $\alpha(T_2^m(G)) = \frac{1}{2}(q-1)$ . If  $m$  is even,  $2 \leq m \leq r-2$ , then the set  $V_0 \cup V_{q-2} \cup V_{q-1}$  is independent in  $T_2^m(G_r)$  and thus  $\alpha(T_2^m(G_r)) \geq |V_0 \cup V_{q-2} \cup V_{q-1}| = 1 + t^{q-2} + t^{q-1} > t^2(t^{q-1} - 1)/(t^2 - 1)/(t^2 - 1) = \alpha(G_r)$ ; this inequality may be easily proved. Therefore no graph  $T_2^m(G_r)$  for  $1 \leq m \leq r-1$  is isomorphic to  $G_r$  and thus the periodicity of  $G_r$  in  $T_2$  is  $r$ . ■

We shall remark also the periodicity 2. In [1] F. Harary, C. Hoede and D. Kladdeck have proved that if a graph  $G$  is self-complementary, i.e.,  $\overline{G} \cong G$ , then  $T_2(G) \cong G$  and thus the periodicity of  $G$  in  $T_2$  is 1. A slight generalization of the result is the following proposition. The diameter of  $G$  is denoted by  $\text{diam } G$ .

**Proposition 1.** *Let  $G$  be a graph such that  $\text{diam } G = \text{diam } \overline{G} = 2$  and  $\overline{G}$  is not isomorphic to  $G$ . Then  $G$  is  $T_2$ -periodic with the periodicity 2.*

**Proof.** If two vertices  $x, y$  are adjacent in  $G$ , then their distance in  $G$  is 1 and they are not adjacent in  $T_2(G)$ . If they are not adjacent in  $G$ , then their distance in  $G$  is 2 and realizes  $\text{diam } G$ . Moreover,  $x$  and  $y$  are adjacent in  $T_2(G)$ . Hence  $T_2(G) = \overline{G}$ . As also  $\text{diam } \overline{G} = 2$ , we have  $T_2^2(G) = T_2(T_2(G)) = T_2(\overline{G}) = G$ . ■

We shall create a class of graph which have the property that  $\text{diam } G = \text{diam } \overline{G} = 2$ .

Let  $H_1, H_2, H_3, H_4, H_5$  be pairwise disjoint graphs. The graph  $G(H_1, H_2, H_3, H_4, H_5)$  contains mentioned graphs as subgraphs and has new edges  $xy$  created in the following way. If  $x \in V(H_i)$ ,  $y \in V(H_j)$ , then  $x$  and  $y$  are adjacent in  $G$  if and only if  $j \equiv i + 1 \pmod{5}$  or  $j \equiv i + 4 \pmod{5}$ . The simplest is the graph  $G(K_1, K_1, K_1, K_1, K_1) = C_5$ .

**Proposition 2.** *For any five graphs  $H_1, H_2, H_3, H_4, H_5$  the graph  $G(H_1, H_2, H_3, H_4, H_5)$  has the diameter 2 and so has its complement.*

**Proof.** Let  $x, y$  be two vertices of  $G(H_1, H_2, H_3, H_4, H_5)$ . Let  $i, j$  be such numbers from  $\{1, 2, 3, 4, 5\}$  that  $x \in V(H_i)$ ,  $y \in V(H_j)$ .

If  $i = j$ , then both  $x, y$  are in the graph  $H_i$ . If they are adjacent in  $G$ , then their distance is 1. If they are not adjacent, then there exists a path of length 2 connecting them; its inner vertex is in  $V(H_{j+1}) \cup V(H_{i+4})$ , the subscripts being taken modulo 5. If  $j \equiv i + 1 \pmod{5}$  or  $j \equiv i + 4 \pmod{5}$ , then  $x, y$  are adjacent in  $G$  and their distance is 1.

If  $j \equiv i + 2 \pmod{5}$  or  $j \equiv i + 3 \pmod{5}$  then  $x, y$  are not adjacent, but there exists a path of length 2 connecting them; its inner vertex is in  $V(H_{i+1}) \cup V(H_{i+4})$ . Therefore  $\text{diam } G = 2$ . The complement of  $G(H_1, H_2, H_3, H_4, H_5)$  is isomorphic to  $G(\overline{H}_1, \overline{H}_2, \overline{H}_3, \overline{H}_4, \overline{H}_5)$  and thus also  $\text{diam } \overline{G} = 2$ . ■

## References

- [1] F. Harary, C. Hoede and D. Kladdeck, *Graph-valued functions related to step graphs*, J. Comb. Ing. Syst. Sci. **7** (1982) 231–246.
- [2] E. Prisner, Graph Dynamics (Longman House, Burnt Mill, Harlow, 1995).

Received 18 February 2000

Revised 5 July 2000