Discussiones Mathematicae Graph Theory 21 (2001) 43–62

FULL DOMINATION IN GRAPHS

ROBERT C. BRIGHAM

Department of Mathematics University of Central Florida, Orlando, FL 32816

GARY CHARTRAND

Department of Mathematics and Statistics Western Michigan University, Kalamazoo, MI 49008

RONALD D. DUTTON

Program of Computer Science University of Central Florida, Orlando, FL 32816

AND

Ping Zhang¹

Department of Mathematics and Statistics Western Michigan University, Kalamazoo, MI 49008

Abstract

For each vertex v in a graph G, let there be associated a subgraph H_v of G. The vertex v is said to dominate H_v as well as dominate each vertex and edge of H_v . A set S of vertices of G is called a full dominating set if every vertex of G is dominated by some vertex of S, as is every edge of G. The minimum cardinality of a full dominating set of G is its full domination number $\gamma_{FH}(G)$. A full dominating set of G of cardinality $\gamma_{FH}(G)$ is called a γ_{FH} -set of G. We study three types of full domination in graphs: full star domination, where H_v is the maximum star centered at v, full closed domination, where H_v is the subgraph induced by the closed neighborhood of v, and full open domination, where H_v is the subgraph induced by the subgraph induced by the open neighborhood of v.

Keywords: full domination, full star domination, full closed domination, full open domination.

2000 Mathematics Subject Classification: 05C12.

 $^{^1\}mathrm{Research}$ supported in part by the Western Michigan University Research Development Award Program Fund

1. Introduction

A vertex v in a graph G is said to *dominate* itself and each of its neighbors. A set $S \subseteq V(G)$ is called a *dominating set* for G if every vertex of G is dominated by some vertex of S. The minimum cardinality of a dominating set is the *domination number* $\gamma(G)$ of G. A dominating set of cardinality $\gamma(G)$ is a γ -set for G. There has been increased interest in recent years in the study of domination in graphs. Indeed, the books [2, 3] by Haynes, Hedetniemi, and Slater are devoted exclusively to this subject. In domination, a vertex dominates a set of vertices (according to some rule); while in covering, a vertex covers the edges incident with it. We combine these concepts to describe another variation of domination.

For a graph G, let H be a function that maps each vertex v of G into a subgraph H_v of G. In this context, the vertex v is said to dominate H_v as well as dominate each vertex and edge of H_v . A set S of vertices of G is called a full dominating set if every vertex and every edge of G is dominated by some vertex of S. For each full dominating set S of G and $v \in V(G) - S$, the set $S \cup \{v\}$ is also a full dominating set. If G has no isolated vertices, then we need only be concerned with each edge of G being dominated by some vertex of S. The minimum cardinality of a full dominating set of G is its full domination number (with respect to the function H) and is denoted by $\gamma_{FH}(G)$. A full dominating set of G of cardinality $\gamma_{FH}(G)$ is called a γ_{FH} -set of G. Certainly, $\gamma_{FH}(G)$ is defined for a graph G if and only if V(G)is a full dominating set for G.

In this paper we study three examples of full domination, namely full star domination, where H_v is the maximum star S_v centered at v, full closed domination, where $H_v = \langle N[v] \rangle$, the subgraph induced by the closed neighborhood of v, and full open domination, where $H_v = \langle N(v) \rangle$, the subgraph induced by the open neighborhood of v.

2. Full Star Domination in Graphs

We denote the *full star domination number* of a graph G by $\gamma_{FS}(G)$. Certainly, $\gamma_{FS}(G)$ is defined for every graph G. Indeed, if G is a graph without isolated vertices, then $\gamma_{FS}(G) = \alpha_o(G)$, the vertex covering number of G (the minimum number of vertices that cover all edges of G). If G has I(G)isolated vertices, then $\gamma_{FS}(G) = \alpha_o(G) + I(G)$. Therefore, the full star domination number is not a new parameter; it only provides a new setting for an old one. A well-known theorem of Gallai [1] states that if G is a graph of order n without isolated vertices, then $\alpha_o(G) + \beta_o(G) = n$, where $\beta_o(G)$ is the vertex independence number of G. This gives us the following.

Observation 2.1. For every graph G of order n without isolated vertices,

$$\gamma_{FS}(G) = n - \beta_o(G).$$

Since every full star dominating set of a graph is also a dominating set, it follows that $\gamma(G) \leq \gamma_{FS}(G)$. By Observation 2.1,

$$1 \le \gamma(G) \le \gamma_{FS}(G) \le n - 1$$

for every graph G of order n with at most n-2 isolated vertices. We now consider the realizability of three integers a, b, n as the domination number, full star domination number, and order, respectively, of some graph without isolated vertices. Thus any such triple a, b, n described above must satisfy $1 \le a \le b \le n-1$. By Observation 2.1, however, $\gamma_{FS}(G) = n-1$ if and only if $G = K_n$, which implies that $\gamma(G) = 1$. Hence we may assume that $1 \le a \le b \le n-2$. On the other hand, the independent domination number i(G) satisfies

$$\gamma(G) \le i(G) \le \beta_o(G) = n - \gamma_{FS}(G).$$

This implies that $\gamma(G) + \gamma_{FS}(G) \leq n$, thereby obtaining Ore's [6] well-known inequality $\gamma(G) \leq n/2$ for graphs G of order n without isolated vertices. We now present the desired realization result.

Proposition 2.2. For every triple a, b, n of integers with $n \ge 3$, $1 \le a \le b \le n-2$, and $a+b \le n$, there exists a graph G of order n without isolated vertices such that $\gamma(G) = a$ and $\gamma_{FS}(G) = b$.

Proof. We consider two cases.

Case 1. $a + b \leq n - 1$. Let K_{b+1} be the complete graph with vertex set $\{u_1, u_2, \ldots, u_{b+1}\}$ and let G be the graph obtained from K_{b+1} by adding n-b-1 new vertices $v_1, v_2, \ldots, v_{n-b-1}$, the a-1 edges $u_i v_i$ $(1 \leq i \leq a-1)$, and the n-b-a edges $u_a v_i$ $(a \leq i \leq n-b-1)$. The graph G is shown in Figure 1. Since $\{u_1, u_2, \ldots, u_a\}$ is a γ -set and $\{u_1, u_2, \ldots, u_b\}$ is a γ_{FS} -set for G, it follows that $\gamma(G) = a$ and $\gamma_{FS}(G) = b$.



Figure 1. The graph G in Case 1

Case 2. a + b = n. Let $K_{\ell_1}, K_{\ell_2}, \ldots, K_{\ell_a}$ be complete graphs, where $\ell_i \geq 2$ for all i and $\sum_{i=1}^{a} \ell_i = n$. Also, let v_{i1} and v_{i2} be distinct vertices in K_{ℓ_i} $(1 \leq i \leq a)$. Then the graph G is obtained from the graph $\bigcup_{i=1}^{a} K_{\ell_i}$ by adding the a - 1 edges $v_{i1} v_{i+1,2}$ for $1 \leq i \leq a - 1$. For a = 4, the graph G is shown in Figure 2. Since $\{v_{11}, v_{21}, \ldots, v_{a1}\}$ is a γ -set, $\gamma(G) = a$. On the other hand,

$$\bigcup_{i=1}^{a} V(K_{\ell_i}) - \{v_{12}, v_{22}, \dots, v_{a2}\}$$

is a γ_{FS} -set and so $\gamma_{FS}(G) = n - a = b$.

46



Figure 2. The graph G in Case 2 when a = 4

3. Full Closed Domination in Graphs

Recall that a set S of vertices in a graph G is a full closed dominating set if every vertex and edge of G belongs to $\langle N[v] \rangle$ for some $v \in S$. The minimum cardinality of a full closed dominating set is the full closed domination number $\gamma_{FC}(G)$. A full closed dominating set of cardinality $\gamma_{FC}(G)$ is referred to as a γ_{FC} -set. This parameter was first introduced by Sampathkumar and Neeralagi in [5], where it was called the neighborhood number of a graph, and further studied by Jayaram, Kwong, and Straight in [4]. The following two propositions appeared in [5].

Proposition 3.1. For every graph G, $\gamma(G) \leq \gamma_{FC}(G) \leq \gamma_{FS}(G)$.

Proposition 3.2. If G is a triangle-free graph, then $\gamma_{FC}(G) = \gamma_{FS}(G)$.

The converse of Proposition 3.2 is not true in general unless $\gamma_{FC}(G) = \gamma_{FS}(G) = 1$, in which case G is a star. To see this, we recall that the corona of a graph G with $V(G) = \{v_1, v_2, \ldots, v_n\}$ is that graph of order 2n obtained from G by adding n new vertices u_1, u_2, \ldots, u_n and the n new edges $u_i v_i$ $(1 \leq i \leq n)$. For $n \geq 3$, let G_n denote the corona of K_n . Define G_2 as the graph obtained from G_3 by deleting an end-vertex. Then $\gamma_{FS}(G_n) = \gamma_{FC}(G_n) = n$ for $n \geq 2$ but certainly G_n is not triangle-free.

If $\gamma(G) = 1$, then $\gamma_{FC}(G) = 1$ while $1 \leq \gamma_{FS}(G) \leq n-1$. For each integer k with $1 \leq k \leq n-1$, the graph H obtained by deleting the edges of a complete subgraph of order n-k from K_n has $\gamma(H) = \gamma_{FC}(H) = 1$ and $\gamma_{FS}(H) = k$. For $\gamma(G) \geq 2$, the following realization result appeared in [4].

Theorem 3.3. For every triple a, b, c of integers with $2 \le a \le b \le c$, there exists a graph G with $\gamma(G) = a$, $\gamma_{FC}(G) = b$, and $\gamma_{FS}(G) = c$.

It is often of interest to know how the value of a graphical parameter is affected when a small change is made in a graph. In this connection, we now consider this question in the case of $\gamma_{FC}(G)$ when an edge is deleted from G. We show, in fact, that such an operation produces a graph whose full closed domination number differs from that of the original graph by at most 1.

Proposition 3.4. For each edge e of a graph G,

$$|\gamma_{FC}(G) - \gamma_{FC}(G - e)| \le 1.$$

Proof. Let e = uv be an edge of G and let S be a γ_{FC} -set of G - e. Then $S \cup \{u\}$ is a full closed dominating set of G. So $\gamma_{FC}(G) \leq |S \cup \{u\}| \leq \gamma_{FC}(G-e) + 1$. Next we show that $\gamma_{FC}(G-e) \leq \gamma_{FC}(G) + 1$. We consider two cases.

Case 1. There exists a γ_{FC} -set S' of G such that $u, v \notin S'$. Then S' is a full closed dominating set of G - e as well. Therefore, $\gamma_{FC}(G - e) \leq |S'| = \gamma_{FC}(G) < \gamma_{FC}(G) + 1$.

Case 2. For every γ_{FC} -set S of G, at least one of u and v belongs to S. Since $S \cup \{u, v\}$ is a full closed dominating set of G - e, it follows that

$$\gamma_{FC}(G-e) \le |S \cup \{u, v\}| \le |S| + 1 = \gamma_{FC}(G) + 1.$$

The bounds presented in Proposition 3.4 are sharp. To see this, we consider the graph G of Figure 3, where $\gamma_{FC}(G) = 3$ and the vertices of a γ_{FC} -set in G are indicated by solid circles. Observe that



Figure 3. How the full closed domination number is affected by the removal of an edge

In view of Proposition 3.4, the edge set of a graph G can be partitioned into the following subsets:

$$E_{0}(G) = \{e \in E(G) : \gamma_{FC}(G-e) = \gamma_{FC}(G)\},\$$

$$E_{-}(G) = \{e \in E(G) : \gamma_{FC}(G-e) = \gamma_{FC}(G) - 1\},\$$

$$E_{+}(G) = \{e \in E(G) : \gamma_{FC}(G-e) = \gamma_{FC}(G) + 1\}.\$$

The graph G of Figure 3 shows that it is possible for all three of these subsets to be nonempty for a single graph G. We now present some facts concerning elements in $E_{-}(G)$ and $E_{+}(G)$.

Proposition 3.5. Let G be a graph containing an edge e = uv. Then $e \in E_{-}(G)$ if and only if for every γ_{FC} -set S' of G - e,

(a) neither u, v, nor any common neighbor of u and v belongs to S', and

(b) for each $w \in N_G[u] \cap N_G[v]$, the set $S' \cup \{w\}$ is a γ_{FC} -set of G.

Proof. Suppose that $e = uv \in E_{-}(G)$. Let S' be a γ_{FC} -set of G - e. We first verify (a). Assume, to the contrary, that either u, v, or some common neighbor of u and v belongs to S'. Thus S' is also a full closed dominating set of G. So $\gamma_{FC}(G) \leq |S'|$, a contradiction. To verify (b), let $w \in N_G[u] \cap N_G[v]$. Then $w \notin S'$ by (a). Let $S = S' \cup \{w\}$. Thus $|S| = |S'| + 1 = \gamma_{FC}(G)$. Since S is a full closed dominating set of G, it follows that S is a γ_{FC} -set for G and so (b) holds.

For the converse, let S' be a γ_{FC} -set of G - e, satisfying (a) and (b). It then follows from (a) that $S' \cap (N_G[u] \cap N_G[v]) = \emptyset$. Let $w \in N_G[u] \cap N_G[v]$. By (b) the set $S = S' \cup \{w\}$ is a γ_{FC} -set of G and so $|S| = |S'| + 1 = \gamma_{FC}(G)$. Thus $|S| = \gamma_{FC}(G) - 1$, implying that $e \in E_-(G)$.

Proposition 3.6. Let G be a graph containing an edge e = uv. Then $e \in E_+(G)$ if and only if for every γ_{FC} -set S of G,

- (a) $|S \cap \{u, v\}| = 1$, and
- (b) $S \cup \{u, v\}$ is a γ_{FC} -set of G e.

Proof. Let $e = uv \in E_+(G)$ and let S be a γ_{FC} -set of G. First we verify (a). Assume, to the contrary, that $|S \cap \{u, v\}| \neq 1$. If $S \cap \{u, v\} = \emptyset$, then, since e is dominated by some vertex in S, there is a vertex $w \in S$ adjacent to both u and v. However, then, S is a full closed dominating set for G - e, contradicting the fact that $e \in E_+(G)$. On the other hand, if $\{u, v\} \subseteq S$, then, once again, S is full closed dominating set for G - e, a contradiction. Next we verify (b). Certainly, $S \cup \{u, v\}$ is a full closed dominating set of G - e. By (a), however,

$$|S \cup \{u, v\}| = |S| + 1 = \gamma_{FC}(G) + 1 = \gamma_{FC}(G - e)$$

and so $S \cup \{u, v\}$ is a γ_{FC} -set for G - e.

For the converse, let S be a γ_{FC} -set of G that satisfies (a) and (b). By (a), S contains exactly one of u and v. Let $S' = S \cup \{u, v\}$ and so |S'| = |S| + 1. By (b), S' is a γ_{FC} -set of G - e. Thus $\gamma_{FC}(G - e) = |S'| = |S| + 1 = \gamma_{FC}(G) + 1$, implying that $e \in E_+(G)$. If we were to delete two edges from G, one belonging to $E_{-}(G)$ and the other belonging to $E_{+}(G)$, then the full closed domination number of the resulting graph is the same as $\gamma_{FC}(G)$.

Proposition 3.7. Let G be a graph. If $e_1 \in E_-(G)$ and $e_2 \in E_+(G)$, then

$$\gamma_{FC}(G - e_1 - e_2) = \gamma_{FC}(G)$$

Proof. Removing e_1 first and then e_2 shows that $\gamma_{FC}(G - e_1 - e_2) \leq \gamma_{FC}(G)$; while removing e_2 first and then e_1 produces the inequality $\gamma_{FC}(G - e_2 - e_1) \geq \gamma_{FC}(G)$.

Accordingly, if the edges e_1 and e_2 of graph G of Figure 3 are deleted, then $\gamma_{FC}(G - e_1 - e_2) = 3$ since $\gamma_{FC}(G) = 3$. Observe that the edges e_1 and e_2 of this graph are not adjacent. This is no coincidence as we next show.

Proposition 3.8. Let G be a graph. If $e_1 \in E_-(G)$ and $e_2 \in E_+(G)$, then e_1 and e_2 are not adjacent in G.

Proof. Assume, to the contrary, that there exists a graph G containing adjacent edges e_1 and e_2 with $e_1 \in E_-(G)$ and $e_2 \in E_+(G)$. Let $e_1 = uv$ and $e_2 = vw$. Let S' be a γ_{FC} -set for G-uv. By Proposition 3.5, $v \notin S'$. Let $S'' = S' \cup \{v\}$ and consider the graph G - vw. The edge uv is dominated by $v \in S''$. Since vw is dominated by some vertex of S', it follows that either vw is dominated by $w \in S'$ or dominated by some $x \in S'$, where x is adjacent to both v and w. In either case, w is dominated in G - vw by some vertex of S''. Hence, S'' is a full closed dominating set for G - vw.

$$|S''| = \gamma_{FC}(G - uv) + 1 = \gamma_{FC}(G) < \gamma_{FC}(G - vw),$$

which is impossible.

By Proposition 3.8 then, for the graph G of Figure 3, it follows that $e_3 \in E_0(G)$. Indeed, if G is a connected graph in which $E_+(G) \neq \emptyset$ and $E_-(G) \neq \emptyset$, then $E_0(G) \neq \emptyset$. There are numerous graphs G in which every edge of G belongs to $E_0(G)$, such as even cycles and K_n $(n \geq 3)$. There is, however, no graph G in which every edge belongs to $E_+(G)$.

Proposition 3.9. No graph G exists every edge of which belongs to $E_+(G)$.

Proof. Assume, to the contrary, that there exists a graph G such that $E(G) = E_+(G)$. Let S be a γ_{FC} -set of G. Then, by Proposition 3.6, for every edge uv in G, one of u and v belongs to S and the other to V(G) - S. This implies that G is a bipartite graph with partite sets S and V(G) - S. Thus G is triangle-free. By Observation 2.1, $\gamma_{FC}(G) = \gamma_{FS}(G) = \alpha_o(G)$. However,

$$\gamma_{FC}(G) + 1 = \gamma_{FC}(G - e) = \alpha_o(G - e) \le \alpha_o(G) = \gamma_{FC}(G)$$

for every edge e in G, which contradicts the fact that $e \in E_+(G)$.

There are graphs G, though, every edge of which belongs to $E_{-}(G)$. For example, odd cycles of order at least 5 have this property.

For a set S of vertices of a graph G and a vertex v of G, the *distance* between v and S is defined as

$$d(v, S) = \min\{d(v, u) : u \in S\}.$$

The *diameter* of S is defined as

$$\operatorname{diam} S = \max\{d(u, v) : u, v \in S\}.$$

Thus $\operatorname{diam} V(G) = \operatorname{diam} G$.

For a nonempty set S of vertices in a connected graph G, a Steiner Stree is a tree of minimum size in G containing S. Certainly, every end-vertex of a Steiner S-tree belongs to S. An edge e = uv in a Steiner S-tree T is called S-free if both $u \notin S$ and $v \notin S$.

Lemma 3.10. For every γ_{FC} -set S of a connected graph G, there exists a Steiner S-tree containing no S-free edges.

Proof. Assume, to the contrary, that there is a connected graph G and a γ_{FC} -set S of G such that every Steiner S-tree in G contains S-free edges. Among all Steiner S-trees, let T be a Steiner S-tree containing a minimum number of S-free edges. Then T contains an S-free edge e = uv and a vertex $x \in S$ such that x, u, v is a path in T. Since S is a γ_{FC} -set of G, it follows that e is dominated by some vertex in S. If e is dominated by x, then necessarily x is adjacent to both u and v. Hence (T-uv) + xv is a Steiner S-tree in G containing fewer S-free edges than T, which is impossible. Thus e is dominated by some vertex $w \in S$, where $w \neq x$. Let T_u and T_v be the two components of T-uv, where T_u contains u and T_v contains v. Necessarily, w belongs to exactly one of T_u and T_v , say T_u . Then (T-uv) + wv is a Steiner S-tree in G containing fewer S-free edges than T, again an impossibility.

Lemma 3.11. If S is a γ_{FC} -set of a connected graph G, then the order of any Steiner S-tree is at most $2\gamma_{FC}(G) - 1$.

Proof. Let T be a Steiner S-tree containing no S-free edges. Then $V(T) = S \cup W$, where $S \cap W = \emptyset$. Let |W| = a. Thus the order of T is $a + \gamma_{FC}(G)$ and the size of T is $a + \gamma_{FC}(G) - 1$. Assume that $a \ge \gamma_{FC}(G)$. Since T is a Steiner S-tree, every end-vertex in T belongs to S. Thus every vertex in W has degree at least 2 in T. Also, since T has no S-free edge, every vertex in W is adjacent only to vertices of S in T.

Therefore, the size of T is at least

$$\sum_{w \in W} \deg_T w \ge 2a \ge a + \gamma_{FC},$$

producing a contradiction.

Corollary 3.12. If S is a γ_{FC} -set in a connected graph G, then

diam
$$S \leq 2\gamma_{FC}(G) - 2.$$

Proof. Let T be a Steiner S-tree of G and suppose that that order of T is k. By Lemma 3.11, $k \leq 2\gamma_{FC}(G) - 1$. Among all trees of order k, the path P_k has the greatest diameter, namely k - 1, and $k - 1 \leq 2\gamma_{FC}(G) - 2$.

Theorem 3.13. If G is a graph of diameter d, then

$$\gamma_{FC}(G) \ge d/2.$$

Proof. Let x and y be vertices of G such that d(x, y) = d and let S be a γ_{FC} -set in G. Then x is dominated by some vertex $u \in S$ and y is dominated by some $v \in S$. Either u = x or $ux \in E(G)$. Similarly, either v = y or $vy \in E(G)$. Thus, using Corollary 3.12, we have

$$d = \operatorname{diam} G = d(x, y) \le d(u, v) + 2 \le \operatorname{diam} S + 2 \le 2\gamma_{FC}(G),$$

producing the desired result.

To show that the bound presented in Theorem 3.13 is sharp, let $G = P_{2k+1}$ be the path of order 2k + 1. Then diamG = 2k and $\gamma_{FC}(G) = k$, as desired.

A set S of vertices in a graph G is an *open dominating set* (or *total dominating set*) if every vertex of G is adjacent to at least one vertex of S. An open dominating set of minimum cardinality is a *minimum open dominating*

set and its cardinality is the open domination number $\gamma_t(G)$, also called the total domination number. The open domination number is also referred to as the total domination number. No graph containing isolated vertices has an open dominating set.

In order to obtain a relationship between the open domination number and the full closed domination number, we present the following lemma.

Lemma 3.14. For every γ_{FC} -set S in a connected graph and each vertex $v \in S$,

$$d\left(v, S - \{v\}\right) \le 2.$$

Proof. Assume, to the contrary, that there is a γ_{FC} -set S in a connected graph G and a vertex $v \in S$ such that $d(v, S - \{v\}) = k \ge 3$. Let $w \in S$ such that $d(v, w) = d(v, S - \{v\})$ and let $P : v = u_0, u_1, \ldots, u_k = w$ be a v - w geodesic in G. Thus, neither u_1 nor u_2 is in S; for otherwise, $d(v, S - \{v\}) \le 2 < k$. Hence the edge $e = u_1u_2$ is dominated by some vertex $y \in S$ (that is necessarily adjacent to both u_1 and u_2). Hence $d(v, y) \le 2 < k$, a contradiction.

We have already seen (in Corollary 3.12) that if S is a γ_{FC} -set in a connected graph G, then diam $S \leq 2\gamma_{FC}(G) - 2$. We now show that $\gamma_t(G)$ has a similar upper bound.

Theorem 3.15. For every connected graph G,

$$\gamma_t(G) \le 2\gamma_{FC}(G) - 1.$$

Proof. Let S be a γ_{FC} -set. Since S is also a dominating set for G, every vertex in V(G) - S is dominated by and therefore adjacent to some vertex in S. Consequently, S openly dominates all vertices in V(G) - S. By Lemma 3.14, for every vertex $u \in S$, there is a vertex $v \neq u$ in S such that $d(u, v) \leq 2$. If every vertex in S is adjacent to some vertex in S, then S is also an open dominating set and so $\gamma_t(G) \leq \gamma_{FC}(G)$. On the other hand, suppose that there is a vertex $x \in S$ that is adjacent to no vertex in S. Then there is a vertex $y \in S$ such that d(x, y) = 2. Let w be a vertex of G adjacent to x and y. Hence $w \notin S$. For each vertex $u \in S - \{x, y\}$, let u' be a vertex of G that is adjacent to u. So u' may or may not be in S. Let $S' = \{u' \mid u \in S - \{x, y\}\}$. Then $S \cup S' \cup \{w\}$ is an open dominating set, $|S \cup S' \cup \{w\}| \leq 2\gamma_{FC}(G) - 1$, and so $\gamma_t(G) \leq 2\gamma_{FC}(G) - 1$.

To see that the upper bound in Theorem 3.15 is sharp, we show that for each integer $k \geq 5$, there exists a connected graph G_k such that $\gamma_{FC}(G_k) = k$

and $\gamma_t(G) = 2k - 1$. Let G_k be the graph obtained from a cycle C_k : $v_0, v_1, v_2, \ldots, v_{k-1}, v_0$ by (i) adding a vertex a_0 and the edges a_0v_0 and v_0v_i for $2 \le i \le k - 2$ and (ii) adding the vertices a_i and b_i $(1 \le i \le k - 1)$ and the edges $a_i b_i$ and $b_i v_i$ for $1 \le i \le k - 1$. The graph G_5 is shown in Figure 4.



Figure 4. The graph G_5

4. Full Open Domination in Graphs

A vertex v in a graph G openly dominates the subgraph $\langle N(v) \rangle$ induced by the (open) neighborhood N(v) of v, but v does not openly dominate itself or any edge incident with it. A set S of vertices in G is a full open dominating set if every vertex and every edge of G belongs to $\langle N(v) \rangle$ for some $v \in S$. The minimum cardinality of a full open dominating set is the full open domination number $\gamma_{FO}(G)$. A full open dominating set of cardinality $\gamma_{FO}(G)$ is referred to as a γ_{FO} -set. Note that a graph G has a full open dominating set if and only if G contains no isolated vertices and every edge of G lies on a triangle in G. Consequently, we have the following.

Observation 4.1. Let S be a full open dominating set in a graph G. Every vertex of S (and consequently every edge joining two vertices of S) belongs to a triangle every vertex of which belongs to S.

To illustrate these concepts, consider the graphs $G_1 = P_5 + K_1$ and $G_2 = K_{2,2,2}$ shown in Figure 5. In G_1 , since each edge $v_i v_{i+1}$ $(1 \le i \le 4)$ is openly dominated only by u, the edge uv_1 is openly dominated only by v_2 , and the edge uv_5 is openly dominated only by v_4 , it follows that u, v_2, v_4

belong to every γ_{FO} -set of G_1 . However, the set $\{u, v_2, v_4\}$ is not a full open dominating set of G_1 as the edges uv_2 and uv_4 are not openly dominated by any vertex in $\{u, v_2, v_4\}$. Since $S_1 = \{u, v_2, v_3, v_4\}$ is a full open dominating set, S_1 is a γ_{FO} -set of G_1 and so $\gamma_{FO}(G_1) = 4$. In G_2 , the set $S_2 =$ $\{x_1, x_2, x_3\}$ is a full open dominating set. Moreover, there is no 2-element full open dominating set. Thus, S_2 is a γ_{FO} -set of G_2 and $\gamma_{FO}(G_2) = 3$.



Figure 5. Graphs $G_1 = P_5 + K_1$ and $G_2 = K_{2,2,2}$

By Observation 4.1, every full open dominating set of a graph G must contain at least three vertices and so $\gamma_{FO}(G) \geq 3$. Certainly, every full open dominating set of a graph G is also a full closed dominating set and so $\gamma_{FO}(G) \geq \gamma_{FC}(G)$. This observation yields the following lower bound for $\gamma_{FO}(G)$.

Corollary 4.2. For a graph G without isolated vertices and in which every edge belongs to a triangle,

 $\gamma_{FO}(G) \ge \max\{3, \gamma_{FC}(G)\}.$

Certainly, if G is a nontrivial connected graph of order n, then $1 \leq \gamma(G) \leq \gamma_{FC}(G) \leq \gamma_{FS}(G) \leq n-1$. Hence no nontrivial connected graph G of order n has $\gamma_{FC}(G) = n$ or $\gamma_{FS}(G) = n$. However, this is not true for $\gamma_{FO}(G)$, as we show next. For a graph G consisting of k ($k \geq 1$) disjoint copies of a graph H, we write G = kH. In particular, G = H for k = 1.

Theorem 4.3. For $n \ge 3$, there exists a connected graph G of order n such that $\gamma_{FO}(G) = n$ if and only if $n \notin \{4, 6\}$.

Proof. We first show that for n = 4 or n = 6, there is no connected graph G of order n with $\gamma_{FO}(G) = n$. If n = 4, then $K_4 - e$ and K_4 are the

56

only graphs of order 4 in which every edge belongs to a triangle. However, $\gamma_{FO}(K_4 - e) = \gamma_{FO}(K_4) = 3$. Next we show that there is no connected graph of order 6 with full open domination number 6. Assume, to the contrary, that there is a connected graph G of order 6 such that $\gamma_{FO}(G) = 6$. Let V(G) = $\{v_1, v_2, \ldots, v_6\}$. Since every edge of G belongs to a triangle in G, there exist at least two triangles in G that have a common edge. Thus G contains $K_4 - e$ as a subgraph. Assume, without loss of generality, that $v_2v_4 \in E(G)$, $v_i v_{i+1} \in E(G)$ for i = 1, 2, 3, and $v_1 v_4 \in E(G)$. Since $\{v_1, v_2, v_3, v_4\}$ openly dominates the induced subgraph $\langle \{v_1, v_2, v_3, v_4\} \rangle$, which is $K_4 - e$ or K_4 , the vertex v_5 must openly dominate v_6 or some edge incident with v_6 . If v_5 openly dominates v_6 , then $v_5v_6 \in E(G)$. However, then, the edge v_5v_6 must be openly dominated by some vertex v_i for $i \in \{1, 2, 3, 4\}$. If v_5 openly dominates some edge e that is incident with v_6 , then $e = v_i v_6$ for some $i \in \{1, 2, 3, 4\}$ and v_5 is adjacent to both v_i and v_6 . In either case, G contains a triangle v_i, v_5, v_6, v_i for some $i \in \{1, 2, 3, 4\}$. Thus, we may assume that G contains at least one of the two graphs G_1 or G_2 of Figure 6 as a subgraph. Let $S = V(G) - \{v_1\}$. Since $\langle S \rangle$ contains $2K_2 + K_1$ as a subgraph, every edge of $\langle S \rangle$ belongs to a triangle. Thus S is a full open dominating set of $\langle S \rangle$. Moreover, the vertex v_1 and all edges incident with v_1 are openly dominated by S. This implies that S is also a full open dominating set of G. Therefore, $\gamma_{FO}(G) \leq |S| = 5$, which is a contradiction.



Figure 6. Subgraphs G_1 and G_2

For the converse, assume that $n \geq 3$ but $n \neq 4, 6$. We construct a graph G of order n with $\gamma_{FO}(G) = n$. If n = 2k + 1 for some integer $k \geq 1$, let $G = kK_2 + K_1$ for some positive integer k. Then the order of G is 2k + 1. Since V(G) is the only full open dominating set, $\gamma_{FO}(G) = 2k + 1$. Now let n = 2k some integer $k \geq 4$. For k = 4, let F be the graph of Figure 7. Note that for every vertex w in G, there is an edge that is only openly dominated by w. Hence V(G) is the only full open dominating set and so $\gamma_{FO}(G) = 8$.



Figure 7. A graph F of order 8 with $\gamma_{FO}(F) = 8$

For $k \geq 5$, let G be obtained from the graph F of Figure 7 and the graph $(k-4)K_2$ by joining every vertex of $(k-4)K_2$ to the vertex v in F. Then the order of G is 2k and $\gamma_{FO}(F) = 2k$, as desired.

Since there is no graph G of order 4 and $\gamma_{FO}(G) = 4$ while the disconnected graph $G = 2K_3$ has order 6 and $\gamma_{FO}(G) = 6$, we have the following corollary.

Corollary 4.4. For $n \ge 3$, there exists a graph G of order n such that $\gamma_{FO}(G) = n$ if and only if $n \ne 4$.

We have seen that if G is a graph in which every edge belongs to a triangle and $\gamma_{FC}(G) = a$ and $\gamma_{FO}(G) = b$, then $1 \le a \le b$ and $b \ge 3$. Next we show that the converse of this fact is also true.

Theorem 4.5. For each pair a, b of integers with $1 \le a \le b$ and $b \ge 3$, there exists a connected graph G in which every edge belongs to a triangle with $\gamma_{FC}(G) = a$ and $\gamma_{FO}(G) = b$.

Proof. We consider three cases, according to whether a = 1, a = 2, or $a \ge 3$.

Case 1. a = 1. Suppose, first, that b is odd. Then b = 2k + 1 for some integer $k \ge 1$. Let $G = kK_2 + K_1$, where $\deg_G u = 2k$. Since $\{u\}$ is a full closed dominating set, $\gamma_{FC}(G) = 1$. Moreover, V(G) is the only full open dominating set, so $\gamma_{FO}(G) = |V(G)| = 2k + 1 = b$. Next suppose that b is even. Then b = 2k for some integer $k \ge 2$. Here we let G = $(P_5 \cup (k-2)K_2) + K_1$, where $P_5 : v_1, v_2, v_3, v_4, v_5$ and $\deg_G u = 2k + 1$. Then $G = P_5 + K_1$ (shown in Figure 5) for k = 2. Again, $\{u\}$ is a full closed dominating set and so $\gamma_{FC}(G) = 1$. On the other hand, the set $\{u, v_2, v_3, v_4\} \cup V((k-2)K_2)$ is a γ_{FO} -set, implying that $\gamma_{FO}(G) = 2k$.

Case 2. a = 2. For b = 3, let $G = K_{2,2,2}$ (shown in Figure 5). Then $\gamma_{FC}(G) = 2$ and $\gamma_{FO}(G) = 3$. For b = 2k + 1, where $k \ge 2$, let G be

obtained from $K_{2,2,2}$ and $F = (k-1)K_2 + K_1$, where $\deg_F u = 2k-2$, by identifying some vertex in $K_{2,2,2}$ and the vertex u in F. Then $\gamma_{FC}(G) = 2$ and $\gamma_{FO}(G) = 3 + 2(k-1) = 2k + 1 = b$.

Now suppose that b is even. Then b = 2k for some integer $k \ge 2$. Let $G_1 = P_5 + \overline{K}_2$, where $P_5 : v_1, v_2, v_3, v_4, v_5$ and the remaining two vertices of G_1 are u and v; and for $k \ge 3$, let $G_2 = (k-2)K_2 + K_1$, where $\deg_{G_2} x = 2k - 4$. For k = 2, let $G = G_1$ and for $k \ge 3$, let G be the graph obtained from G_1 and G_2 by identifying u and x. Then $\gamma_{FC}(G) = 2$, while $\gamma_{FO}(G) = 4 + 2(k-2) = 2k = b$.

Case 3. $a \ge 3$. We consider three subcases here, depending on whether b = a, b = a + 1, or $b \ge a + 2$.

Subcase 3.1. b = a. Let F_0 be a copy of the complete graph K_a with $V(F_0) = \{u_1, u_2, \ldots, u_a\}$. For each i with $1 \le i \le a$, let $F_i : x_i, y_i$ be a copy of K_2 . Then let G be the graph obtained from the graphs F_i $(0 \le i \le a)$ by adding the 4a new edges $u_{i-1}x_i, u_ix_i, u_iy_i$, and $u_{i+1}y_i$ for all $1 \le i \le a$, where each subscript is one of the integers $1, 2, \ldots, a$ modulo a. The graph G is shown in Figure 8 for a = 3, 4. We show that $\gamma_{FO}(G) = \gamma_{FC}(G) = a$.



Figure 8. Graphs G with $\gamma_{FO}(G) = \gamma_{FC}(G) = a$ for a = 3, 4

Let $S = \{u_1, u_2, \ldots, u_a\}$. Since S is a full closed and full open dominating set of G, it follows that $\gamma_{FC}(G) \leq a$ and $\gamma_{FO}(G) \leq a$. On the other hand, each edge $x_i y_i$ $(1 \leq i \leq k)$ in G is dominated only by x_i , y_i , or u_i . Hence every full closed dominating set of G must contain at least one vertex from each set $\{u_i, x_i, y_i\}$ for all $1 \leq i \leq a$. Thus $\gamma_{FC}(G) \geq a$. Since $\gamma_{FO}(G) \geq \gamma_{FC}(G)$, it follows that $\gamma_{FO}(G) \geq a$. Therefore, $\gamma_{FO}(G) = \gamma_{FC}(G) = a$. Subcase 3.2. b = a+1. Let G be obtained from the graph G constructed in Subcase 3.1 by first subdividing the edge x_1y_1 into x_1z and zy_1 and then adding the edge zu_1 . The graph G is shown in Figure 9 for a = 3. Since $S = \{u_1, u_2, \ldots, u_a\}$ is a γ_{FC} -set of G, it follows that $\gamma_{FC}(G) = a$. On the other hand, $S \cup \{x_1\}$ is a γ_{FC} -set and so $\gamma_{FO}(G) = a + 1$.



Figure 9. A graph G with $\gamma_{FO}(G) = 4$ and $\gamma_{FC}(G) = 3$

Subcase 3.3. $b \ge a + 2$. Suppose first that b = a + 2k, where $k \ge 1$. For each integer *i* with $1 \le i \le k$, let $H_i : v_i, w_i$ be a copy of K_2 . Now let *H* be the graph obtained from the graph *G* in Subcase 3.1 and the graphs H_i $(1 \le i \le k)$ by adding the 2k edges u_1v_i, u_1w_i for all *i* with $1 \le i \le k$. For a = 3 and k = 1 (so b = 5), the graph *H* is shown in Figure 10(a).

Next, suppose that b = a + 2k + 1, where $k \ge 1$. Let H be the graph obtained from the graph G in Subcase 3.2 and the graphs H_i $(1 \le i \le k)$ by adding the 2k edges u_1v_i , u_1w_i for all $1 \le i \le k$. For a = 3 and k = 1 (so b = 6), the graph H is shown in Figure 10(b).

Since $S = \{u_1, u_2, \ldots, u_a\}$ is a γ_{FC} -set in H for all $b \ge a + 2$, it follows that $\gamma_{FC}(H) = a$. Next we show that $\gamma_{FO}(H) = b$. Let $V = \{v_1, v_2, \ldots, v_k\}$ and $W = \{w_1, w_2, \ldots, w_k\}$. For b = a + 2k, the set $S_1 = S \cup V \cup W$ is a γ_{FO} -set of H and so $\gamma_{FO}(H) = |S_1| = a + 2k = b$. For b = a + 2k + 1, the set $S_2 = S \cup \{x_1\} \cup V \cup W$ is γ_{FO} -set of H and so $\gamma_{FO}(H) = |S_2| = a + 1 + 2k = b$.

Certainly, every full open dominating set of a graph G is also an open dominating set. Thus if G is a graph without isolated vertices in which every edge is in a triangle, then $\gamma_{FO}(G) \geq \gamma_t(G)$. Next we show that there is no graph G with $\gamma_{FO}(G) = \gamma_t(G)$. 60



Figure 10. Graphs in Subcase 3.3 for a = 3 and b = 5, 6

Proposition 4.6. If G is a graph without isolated vertices in which every edge is in a triangle, then $\gamma_{FO}(G) > \gamma_t(G)$.

Proof. Assume, to the contrary, that there exists a graph G with $\gamma_t(G) = \gamma_{FO}(G)$. Let S be a γ_{FO} -set of G. Since every full open dominating set is also an open dominating set and $\gamma_t(G) = \gamma_{FO}(G)$, it follows that S is also a γ_t -set in G. Let $u \in S$. We consider two cases.

Case 1. There exists a vertex x that is openly dominated by vertex $u \in S$ but not by any vertex in $S - \{u\}$. This implies that x is adjacent to u, but x is not adjacent to any vertex in $S - \{u\}$. On the other hand, since S is a γ_{FO} -set of G, the edge ux is openly dominated by some vertex in $v \in S - \{u\}$. Hence ux belongs to $\langle N(v) \rangle$, implying that x is adjacent to $v \in S - \{u\}$, a contradiction.

Case 2. Each vertex in G that is openly dominated by u is also openly dominated by some vertex in $S - \{u\}$. Since S is a γ_t -set in G, there is a vertex v adjacent to u such that v is not adjacent to any other vertex in $S - \{u\}$. However, then, the edge uv is not openly dominated by any vertex in S, a contradiction.

Next we show that every pair a, b of integers with $2 \le a < b$ is realizable as the open domination number and the full open domination number, respectively, of some graph.

Theorem 4.7. For every pair a, b of integers with $2 \le a < b$, there exists a graph G with $\gamma_t(G) = a$ and $\gamma_{FO}(G) = b$.

Proof. We consider two cases.

Case 1. b = a + 1 or b = a + 2. Let G_a be the graph obtained from K_{a+1} with $V(K_{a+1}) = \{v_1, v_2, \ldots, v_{a+1}\}$ by adding, for each edge $e_{ij} = v_i v_j$, where $1 \le i < j \le a+1$, a new vertex w_{ij} and joining it to v_i and v_j . The graphs G_2 and G_3 are shown in Figure 11. Let $H_a = G_{a+1} - w_{12}$. The graph H_2 is shown in Figure 11. Since $\{v_1, v_2, \ldots, v_a\}$ is a γ_t -set of G_a and $\{v_1, v_2, \ldots, v_{a+1}\}$ is a γ_{FO} -set of G_a , it follows that $\gamma_t(G_a) = a$ and $\gamma_{FO}(G_a) = a + 1$. Moreover, since $\{v_3, v_4, \ldots, v_{a+2}\}$ is a γ_t -set of H_a and $\{v_1, v_2, \ldots, v_{a+2}\}$ is a γ_{FO} -set of H_a , we have $\gamma_t(H_a) = a$ and $\gamma_{FO}(H_a) = a + 2$.



Figure 11. The graphs G_2 , G_3 , and H_2

Case 2. $b \ge a+3$. Suppose, first, that b = a + 2k + 1 $(k \ge 1)$. Let G be the graph obtained from the graph G_a (from Case 1) and kK_2 by joining v_1 to each of the vertices of kK_2 . If b = a + 2k $(k \ge 2)$, then let G be the graph obtained from the graph H_a (of Case 1) and $(k-1)K_2$ by joining v_3 to each vertex of $(k-1)K_2$. It is routine to verify that $\gamma_t(G) = a$ and $\gamma_{FO}(G) = b$.

We conclude this paper with a problem.

Problem 4.8. Determine all triples a, b, c of integers with $\max\{a, b\} \leq c$, $a \geq 2, c \geq 3$, and a < c for which there exists a graph G with $\gamma_t(G) = a$, $\gamma_{FC}(G) = b$, and $\gamma_{FO}(G) = c$.

References

- T. Gallai, Über extreme Punkt- und Kantenmengen, Ann. Univ. Sci. Budapest, Eötvös Sect. Math. 2 (1959) 133–138.
- [2] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, Fundamentals of Domination in Graphs (Marcel Dekker, New York, 1998).
- [3] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, Domination in Graphs: Advanced Topics (Marcel Dekker, New York, 1998).
- [4] S.R. Jayaram, Y.H.H. Kwong and H.J. Straight, Neighborhood sets in graphs, Indian J. Pure Appl. Math. 22 (1991) 259–268.
- [5] E. Sampathkumar and P.S. Neeralagi, *The neighborhood number of a graph*, Indian J. Pure Appl. Math. **16** (1985) 126–136.
- [6] O. Ore, Theory of Graphs, Amer. Math. Soc. Colloq. Publ. 38 (Amer. Math. Soc. Providence, RI, 1962).

Received 5 July 2000 Revised 17 October 2000