Discussiones Mathematicae Graph Theory 22 (2002) 111–112

A PROOF OF MENGER'S THEOREM BY CONTRACTION

FRANK GÖRING

Department of Mathematics Technical University of Ilmenau D-98684 Ilmenau Germany

Abstract

A short proof of the classical theorem of Menger concerning the number of disjoint AB-paths of a finite graph for two subsets A and B of its vertex set is given. The main idea of the proof is to contract an edge of the graph.

Keywords: connectivity, disjoint paths, digraph, Menger. 2000 Mathematics Subject Classifications: 05C40.

Proofs of Menger's Theorem are given in [7, 6, 4, 8, 2]. A short proof is given by T. Böhme, F. Göring and J. Harant in [1]; another short proof based on edge deletion is given by the author in [5]. The new idea here is to get a short proof by contracting an arbitrary edge of the original graph.

For terminology and notation not defined here we refer to [3]. A graph with no edges is denoted by its vertex set. Let G be a finite graph (loops and multiple edges being allowed). For an edge e of G let G - e and G/e denote the graphs obtained from G by removing e and contracting e to one vertex v_e , respectively. For (possibly empty) sets of vertices A and B of G let an AB-separator be a set of vertices of G such that the graph obtained from Gby deleting these vertices contains no path from A to B. Note that a single vertex of $A \cap B$ is considered as a path from A to B, too. An AB-connector is a subgraph of G such that each of its components is a path from A to Bhaving only one vertex in common with A and B, respectively. In particular the empty graph is also an AB-connector. If we contract an egde incident with a vertex of A or B then the resulting vertex is considered to be a vertex of A or B, respectively. **Theorem** (Menger, 1927). Let G be a finite graph, A and B sets of vertices of G, and s the minimum number of vertices forming an AB-separator. Then there is an AB-connector C with $|C \cap A| = s$.

Proof. If G is edgeless then set $C = A \cap B$. Suppose, G is a counterexample with |E(G)| minimal. Then G contains an edge e from x to y and G/e has an AB-separator S with |S| < s, otherwise we are done. Obviously, $v_e \in S$. Then $P = (S \setminus \{v_e\}) \cup \{x, y\}$ is an AB-separator of G with |P| = |S| + 1 = s. An AP-separator (as well as an PB-separator) of G - e is an AB-separator of G. Consequently, G - e has an AP-connector X and a PB-connector Y containing P. Since $X \cap Y = P$, the set $C = (X \cup Y)$ is an AB-connector of G with $|C \cap A| = s$, a contradiction.

References

- T. Böhme, F. Göring and J. Harant, *Menger's Theorem*, J. Graph Theory 37 (2001) 35–36.
- [2] W. McCuaig, A simple proof of Menger's theorem, J. Graph Theory 8 (1984) 427–429.
- [3] R. Diestel, Graph Theory (2nd edition), (Springer-Verlag, New York, 2000).
- [4] G.A. Dirac, Short proof of Menger's graph theorem, Mathematika 13 (1966) 42-44.
- [5] F. Goering, Short Proof of Menger's Theorem, to appear in Discrete Math.
- [6] T. Grünwald (later Gallai), Ein neuer Beweis eines Mengerschen Satzes, J. London Math. Soc. 13 (1938) 188–192.
- [7] K. Menger, Zur allgemeinen Kurventheorie, Fund. Math. 10 (1927) 96–115.
- [8] J.S. Pym, A proof of Menger's theorem, Monatshefte Math. 73 (1969) 81-88.

Received 8 June 2000 Revised 21 May 2001