

## GENERALIZED EDGE-CHROMATIC NUMBERS AND ADDITIVE HEREDITARY PROPERTIES OF GRAPHS

MICHAEL J. DORFLING

*Department of Mathematics*  
*Faculty of Science*  
*Rand Afrikaans University*  
*P.O. Box 524, Auckland Park, 2006 South Africa*

AND

SAMANTHA DORFLING<sup>1</sup>

*Department of Mathematics and Applied Mathematics*  
*Faculty of Science*  
*University of the Free State*  
*P.O. Box 339, Bloemfontein, 9300 South Africa*  
**e-mail:** DorfliS@sci.uovs.ac.za

### Abstract

An additive hereditary property of graphs is a class of simple graphs which is closed under unions, subgraphs and isomorphisms. Let  $\mathcal{P}$  and  $\mathcal{Q}$  be hereditary properties of graphs. The generalized edge-chromatic number  $\rho'_{\mathcal{Q}}(\mathcal{P})$  is defined as the least integer  $n$  such that  $\mathcal{P} \subseteq n\mathcal{Q}$ . We investigate the generalized edge-chromatic numbers of the properties  $\rightarrow H$ ,  $\mathcal{I}_k$ ,  $\mathcal{O}_k$ ,  $\mathcal{W}_k^*$ ,  $\mathcal{S}_k$  and  $\mathcal{D}_k$ .

**Keywords:** property of graphs, additive, hereditary, generalized edge-chromatic number.

**2000 Mathematics Subject Classification:** 05C15.

---

<sup>1</sup>This research forms part of the author's PhD studies at the Rand Afrikaans University.

## 1. Introduction

Following [1] we denote the class of all finite simple graphs by  $\mathcal{I}$ .

A *property* of graphs is a non-empty isomorphism-closed subclass of  $\mathcal{I}$ . We say that a graph  $G$  *has the property*  $\mathcal{P}$  if  $G \in \mathcal{P}$ . A property  $\mathcal{P}$  is called *hereditary* if  $G \in \mathcal{P}$  and  $H \subseteq G$  implies  $H \in \mathcal{P}$ .  $\mathcal{P}$  is called *additive* if  $G \cup H \in \mathcal{P}$  whenever  $G \in \mathcal{P}$  and  $H \in \mathcal{P}$ . A *homomorphism* of a graph  $G$  to a graph  $H$  is a mapping of the vertex set  $V(G)$  into  $V(H)$  such that if  $e = \{u, v\} \in E(G)$ , then  $f(e) = \{f(u), f(v)\} \in E(H)$ . Given a graph  $G$  and a positive integer  $k$  we define  $G[k]$  to be the graph with  $V(G[k]) = V(G) \times \{1, 2, \dots, k\}$  and  $E(G[k]) = \{(u, l_1)(v, l_2) : uv \in E(G)\}$ ;  $G[k]$  is called a *multiplication* of  $G$ . The *clique number*  $\omega(G)$  of a graph  $G$  is the maximum order of a complete subgraph of  $G$ . A *trail* in a graph is a sequence  $u_1u_2, u_2u_3, \dots, u_{k-1}u_k$  of edges, with no edge repeating. If  $u_1 \neq u_k$  then the trail is *open*. Since we will only be interested in the length of a trail, we associate a trail  $T$  with the set of edges in  $T$ .

**Example 1.** For a positive integer  $k$  and a given graph  $H$  we define the following well-known properties:

$$\mathcal{O} = \{G \in \mathcal{I} : E(G) = \emptyset\},$$

$$\mathcal{I}_k = \{G \in \mathcal{I} : G \text{ does not contain } K_{k+2}\},$$

$$\mathcal{O}_k = \{G \in \mathcal{I} : \text{each component of } G \text{ has at most } k+1 \text{ vertices}\},$$

$$\mathcal{W}_k = \{G \in \mathcal{I} : \text{each path in } G \text{ has at most } k \text{ edges}\},$$

$$\mathcal{W}_k^* = \{G \in \mathcal{I} : \text{each open trail in } G \text{ has at most } k \text{ edges}\},$$

$$\mathcal{S}_k = \{G \in \mathcal{I} : \text{the maximum degree of } G \text{ is at most } k\},$$

$$\mathcal{D}_k = \{G \in \mathcal{I} : G \text{ is } k\text{-degenerate, i.e., every subgraph of } G \text{ has a vertex of degree at most } k\},$$

$$\rightarrow H = \{G \in \mathcal{I} : \text{there is a homomorphism from } G \text{ to } H\},$$

$$\mathcal{O}^k = \{G \in \mathcal{I} : G \text{ is } k\text{-colourable}\} \Rightarrow K_k.$$

Note that for a graph  $G$  we have that  $G \in \rightarrow H$  iff  $G$  is a subgraph of a multiplication of  $H$ . A property of the form  $\rightarrow H$  is called a *hom-property*.

Every hereditary property  $\mathcal{P}$  is determined by the set of *minimal forbidden subgraphs*  $\mathbf{F}(\mathcal{P}) = \{G \in \overline{\mathcal{P}} : \text{every proper subgraph of } G \text{ is in } \mathcal{P}\}$ .

If  $G = (V, E)$  is a graph and  $E' \subseteq E$  then the *subgraph of  $G$  induced by  $E'$*  is the graph  $(V, E')$  and is denoted by  $G[E']$ .

Let  $\mathcal{Q}_1, \mathcal{Q}_2, \dots, \mathcal{Q}_n$  be arbitrary hereditary properties of graphs. An *edge  $(\mathcal{Q}_1, \mathcal{Q}_2, \dots, \mathcal{Q}_n)$ -decomposition* of a graph  $G$  is a decomposition

$\{E_1, E_2, \dots, E_n\}$  of  $E(G)$  such that for each  $i = 1, 2, \dots, n$  the induced subgraph  $G[E_i]$  has the property  $\mathcal{Q}_i$ . The property  $\mathcal{R} = \mathcal{Q}_1 \oplus \mathcal{Q}_2 \oplus \dots \oplus \mathcal{Q}_n$  is defined as the set of all graphs having an edge  $(\mathcal{Q}_1, \mathcal{Q}_2, \dots, \mathcal{Q}_n)$ -decomposition. It is easy to see that if  $\mathcal{Q}_1, \mathcal{Q}_2, \dots, \mathcal{Q}_n$  are additive and hereditary, then  $\mathcal{R} = \mathcal{Q}_1 \oplus \mathcal{Q}_2 \oplus \dots \oplus \mathcal{Q}_n$  is additive and hereditary too. If  $\mathcal{Q}_1 = \mathcal{Q}_2 = \dots = \mathcal{Q}_n = \mathcal{Q}$ , then we write  $n\mathcal{Q} = \mathcal{Q}_1 \oplus \mathcal{Q}_2 \oplus \dots \oplus \mathcal{Q}_n$ .

The generalized edge-chromatic number  $\rho'_{\mathcal{Q}}(G)$  of a graph  $G$  is defined as the least integer  $n$  such that  $G \in n\mathcal{Q}$ . For a property  $\mathcal{P}$ ,  $\rho'_{\mathcal{Q}}(\mathcal{P})$  is then the least  $n$  such that  $\mathcal{P} \subseteq n\mathcal{Q}$ .

As an example of the non-existence of  $\rho'_{\mathcal{Q}}(\mathcal{P})$  we have  $\rho'_{\mathcal{S}_1}(\mathcal{D}_1)$  since there exist graphs in  $\mathcal{D}_1$  of arbitrary maximum degree. Theorem 1.1 by J. Nešetřil and V. Rödl (see [6]) implies that for some properties  $\mathcal{P}$ ,  $\rho'_{\mathcal{Q}}(\mathcal{P})$  exists iff  $\rho'_{\mathcal{Q}}(\mathcal{P}) = 1$ . Here a graph  $G$  is called *3-chromatic connected* if there is no  $S \subseteq V(G)$  such that  $G - S$  is disconnected and  $G[S]$  is bipartite.

**Theorem 1.1** [6]. *Let  $\mathbf{F}(\mathcal{P})$  be a set of 3-chromatic connected graphs. Then for every positive integer  $k$  and graph  $G \in \mathcal{P}$  there exists a graph  $H \in \mathcal{P}$  such that for any decomposition  $\{E_1, E_2, \dots, E_k\}$  of  $E(H)$  there is an  $i \in \{1, 2, \dots, k\}$ , for which  $G \subseteq H[E_i]$ .* ■

**Corollary 1.2.** *If  $\mathbf{F}(\mathcal{P})$  is a set of 3-chromatic connected graphs, then for any hereditary property  $\mathcal{Q}$ ,  $\rho'_{\mathcal{Q}}(\mathcal{P})$  exists if and only if  $\mathcal{P} \subseteq \mathcal{Q}$ .* ■

**Proof.** Suppose that  $\mathcal{P} \not\subseteq \mathcal{Q}$  but  $\mathcal{P} \in n\mathcal{Q}$  for some  $n$ . Let  $G \in \mathcal{P}$  and  $G \notin \mathcal{Q}$ . By Theorem 1.1 there is an  $H \in \mathcal{P}$  such that for every decomposition  $\{E_1, E_2, \dots, E_n\}$  of  $E(H)$  there is an  $i \in \{1, 2, \dots, n\}$  for which  $G \subseteq H[E_i]$ . Let  $\{E_1, E_2, \dots, E_n\}$  be an  $n\mathcal{Q}$ -decomposition of  $E(H)$ . Then  $G \subseteq H[E_i] \in \mathcal{Q}$  for some  $i$ , a contradiction. The converse is trivial. ■

In particular, for every  $k$  and any hereditary property  $\mathcal{Q}$  we have that  $\rho'_{\mathcal{Q}}(\mathcal{I}_k)$  exists iff  $\mathcal{I}_k \subseteq \mathcal{Q}$ .

**Lemma 1.3.** *Let  $\mathcal{P}_1, \mathcal{P}_2$  and  $\mathcal{Q}$  be any properties. If  $\mathcal{P}_1 \subseteq \mathcal{P}_2$ , then  $\rho'_{\mathcal{Q}}(\mathcal{P}_1) \leq \rho'_{\mathcal{Q}}(\mathcal{P}_2)$ .* ■

**Lemma 1.4.** *Let  $\mathcal{Q}_1, \mathcal{Q}_2$  and  $\mathcal{P}$  be any properties. If  $\mathcal{Q}_1 \subseteq \mathcal{Q}_2$ , then  $\rho'_{\mathcal{Q}_2}(\mathcal{P}) \leq \rho'_{\mathcal{Q}_1}(\mathcal{P})$ .* ■

The lattice of (additive) hereditary properties is discussed in [1] — we use the supremum and infimum of properties in our next result without further discussion. A similar result is proved in [5].

**Theorem 1.5.** *Let  $\mathcal{P}_1$  and  $\mathcal{P}_2$  be hereditary properties and  $\mathcal{Q}$  an additive hereditary property such that  $\rho'_{\mathcal{Q}}(\mathcal{P}_1)$  and  $\rho'_{\mathcal{Q}}(\mathcal{P}_2)$  are finite. The following hold:*

- (i)  $\rho'_{\mathcal{Q}}(\mathcal{P}_1 \cup \mathcal{P}_2) = \rho'_{\mathcal{Q}}(\mathcal{P}_1 \vee \mathcal{P}_2) = \max\{\rho'_{\mathcal{Q}}(\mathcal{P}_1), \rho'_{\mathcal{Q}}(\mathcal{P}_2)\}.$
- (ii)  $\rho'_{\mathcal{Q}}(\mathcal{P}_1 \cap \mathcal{P}_2) \leq \min\{\rho'_{\mathcal{Q}}(\mathcal{P}_1), \rho'_{\mathcal{Q}}(\mathcal{P}_2)\}.$
- (iii)  $\max\{\rho'_{\mathcal{Q}}(\mathcal{P}_1), \rho'_{\mathcal{Q}}(\mathcal{P}_2)\} \leq \rho'_{\mathcal{Q}}(\mathcal{P}_1 \oplus \mathcal{P}_2) \leq \rho'_{\mathcal{Q}}(\mathcal{P}_1) + \rho'_{\mathcal{Q}}(\mathcal{P}_2).$  ■

In the rest of this paper we aim to study the generalized edge-chromatic number  $\rho'_{\mathcal{Q}}(\mathcal{P})$  with  $\mathcal{Q}$  and  $\mathcal{P}$  amongst the properties listed in Example 1.

## 2. Some Values of $\rho'_{\mathcal{Q}}(\mathcal{P})$

The well-known results of Vizing and Petersen on edge-colourings of graphs imply the following result — see [3] for details.

**Theorem 2.1.** *Let  $p$  and  $q$  be any positive integers. Then*

- 1.  $\mathcal{S}_p \oplus \mathcal{S}_q \subseteq \mathcal{S}_{p+q}.$
- 2.  $\mathcal{S}_p \subseteq (p+1)\mathcal{S}_1.$
- 3. *If  $p$  and  $q$  are even then  $\mathcal{S}_{p+q} = \mathcal{S}_p \oplus \mathcal{S}_q.$*
- 4. *If  $q$  is odd then  $\mathcal{S}_{p+q} \not\subseteq \mathcal{S}_p \oplus \mathcal{S}_q.$*  ■

**Corollary 2.2.** *For all positive integers  $k$  and  $n$ ,*

$$\rho'_{\mathcal{S}_n}(\mathcal{S}_k) = \begin{cases} \left\lceil \frac{k}{n} \right\rceil, & n \text{ even or } k \leq n, \\ \left\lceil \frac{k+1}{n} \right\rceil, & \text{otherwise.} \end{cases}$$

**Proof.** The result is clearly true if  $k \leq n$ . If  $n$  is even then it follows from Part 3 of Theorem 2.1 that  $\mathcal{S}_k \subseteq \left\lceil \frac{k}{n} \right\rceil \mathcal{S}_n$  while the lower bound follows by observing that  $k > n \left( \left\lceil \frac{k}{n} \right\rceil - 1 \right)$  so that  $\mathcal{S}_k \not\subseteq \mathcal{S}_{n(\left\lceil \frac{k}{n} \right\rceil - 1)} = \left( \left\lceil \frac{k}{n} \right\rceil - 1 \right) \mathcal{S}_n$ .

Now let  $n$  be odd and  $k > n$ . By Theorem 2.1 we have that  $\mathcal{S}_k \subseteq (k+1)\mathcal{S}_1 \subseteq n \left\lceil \frac{k+1}{n} \right\rceil \mathcal{S}_1 \subseteq \left\lceil \frac{k+1}{n} \right\rceil \mathcal{S}_n$ . Let  $c = \left\lceil \frac{k+1}{n} \right\rceil - 1$ . Since  $\left\lceil \frac{k+1}{n} \right\rceil \leq \frac{k+1}{n} + \frac{n-1}{n}$  it follows that  $k \geq nc$ . If  $c = 1$  then, since  $k > n$ ,  $\rho'_{\mathcal{S}_n}(\mathcal{S}_k) \geq 2 = c+1 = \left\lceil \frac{k+1}{n} \right\rceil$ , so assume that  $c \geq 2$ . Now  $\mathcal{S}_k \supseteq \mathcal{S}_{cn} = \mathcal{S}_{(c-1)n+n} \not\subseteq \mathcal{S}_{(c-1)n} \oplus \mathcal{S}_n \supseteq (c-1)\mathcal{S}_n \oplus \mathcal{S}_n \supseteq c\mathcal{S}_n$  so that  $\rho'_{\mathcal{S}_n}(\mathcal{S}_k) \geq c+1$ . ■

Our next result states that, in some cases, the determination of the generalized edge-chromatic number  $\rho'_Q(\rightarrow H)$  reduces to the determination of  $\rho'_Q(H)$ .

**Theorem 2.3.** *For any additive hereditary property  $\mathcal{Q}$  which is closed under multiplications and any graph  $H$ ,  $\rho'_Q(\rightarrow H) = \rho'_Q(H)$ .*

**Proof.** Since  $H \in \rightarrow H$  we have  $\rho'_Q(\rightarrow H) \geq \rho'_Q(H)$ . Now suppose that  $H \in m\mathcal{Q}$  and let  $(E_1, E_2, \dots, E_m)$  be an  $m\mathcal{Q}$ -decomposition of  $E(H)$ . If  $G \in \rightarrow H$  then  $G$  is a subgraph of a multiplication of  $H$ . Let, for every  $i \in \{1, 2, \dots, m\}$ ,  $E'_i = \{(u, l_1)(v, l_2) : uv \in E_i\}$ . Then  $G[E'_i]$  is a subgraph of a multiplication of  $H[E_i]$  for every  $i$  and, since  $\mathcal{Q}$  is closed under multiplications and hereditary,  $G[E'_i] \in \mathcal{Q}$ . Therefore  $(E'_1, E'_2, \dots, E'_m)$  is an  $m\mathcal{Q}$ -decomposition of  $E(G)$ , hence  $\rho'_Q(\rightarrow H) \leq \rho'_Q(H)$ . ■

**Theorem 2.4.** *For all positive integers  $n \geq 2$  and  $k$ , if  $\mathcal{P}$  satisfies  $\mathcal{O}_{k-1} \subseteq \mathcal{P} \subseteq \mathcal{O}^k$ , then  $\rho'_{\mathcal{O}^n}(\mathcal{P}) = \lceil \log_n k \rceil$ .*

**Proof.** It is well known that  $\mathcal{O}^{ab} = \mathcal{O}^a \oplus \mathcal{O}^b$  (see e.g. [3]). This implies that  $\mathcal{O}^k \subseteq \mathcal{O}^{n^{\lceil \log_n k \rceil}} = \lceil \log_n k \rceil \mathcal{O}^n$  hence  $\rho'_{\mathcal{O}^n}(\mathcal{O}^k) \leq \lceil \log_n k \rceil$ .

Since  $n^{\lceil \log_n k \rceil - 1} < n^{\log_n k} = k$  it follows that  $K_k \notin \mathcal{O}^{n^{\lceil \log_n k \rceil - 1}} = (\lceil \log_n k \rceil - 1)\mathcal{O}^n$ . Therefore  $\mathcal{O}_{k-1} \not\subseteq (\lceil \log_n k \rceil - 1)\mathcal{O}^n$  and thus  $\rho'_{\mathcal{O}^n}(\mathcal{O}_{k-1}) \geq \lceil \log_n k \rceil$ . Therefore, by Lemma 1.3 it follows that  $\rho'_{\mathcal{O}^n}(\mathcal{P}) = \lceil \log_n k \rceil$ . ■

For our next result we define  $\rho_\chi(\mathcal{P})$  to be the least  $k$  such that  $\mathcal{P} \subseteq \mathcal{O}^k$  and  $\chi^*(\mathcal{P})$  to be the greatest  $k$  such that  $\mathcal{O}^k \subseteq \mathcal{P}$ .

**Corollary 2.5.** *For any additive hereditary properties  $\mathcal{Q}$ ,  $\mathcal{P} \neq \mathcal{I}$  for which  $\rho_\chi(\mathcal{P})$  and  $\rho_\chi(\mathcal{Q})$  exist,  $\lceil \log_{\rho_\chi(\mathcal{Q})} \chi^*(\mathcal{P}) \rceil \leq \rho'_Q(\mathcal{P}) \leq \lceil \log_{\chi^*(\mathcal{Q})} \rho_\chi(\mathcal{P}) \rceil$ .*

**Proof.** Since  $\mathcal{O}^{\chi^*(\mathcal{Q})} \subseteq \mathcal{Q}$  and  $\mathcal{P} \subseteq \mathcal{O}^{\rho_\chi(\mathcal{P})}$  we have by Lemma 1.3, Lemma 1.4 and Theorem 2.4 that  $\lceil \log_{\chi^*(\mathcal{Q})} \rho_\chi(\mathcal{P}) \rceil \geq \rho'_Q(\mathcal{P})$ . Similarly, since  $\mathcal{Q} \subseteq \mathcal{O}^{\rho_\chi(\mathcal{Q})}$  and  $\mathcal{O}^{\chi^*(\mathcal{P})} \subseteq \mathcal{P}$  we have that  $\rho'_Q(\mathcal{P}) \geq \lceil \log_{\rho_\chi(\mathcal{Q})} \chi^*(\mathcal{P}) \rceil$ . ■

Since, for any graph  $H$ ,  $\rho_\chi(\rightarrow H) = \chi(H)$  and  $\chi^*(\rightarrow H) = \omega(H)$  we have the following corollary.

**Corollary 2.6.** *For all graphs  $G$  and  $H$ ,*

$$\lceil \log_{\chi(G)} \omega(H) \rceil \leq \rho'_{\rightarrow G}(\rightarrow H) \leq \lceil \log_{\omega(G)} \chi(H) \rceil. \quad \blacksquare$$

### 3. Some Results on $\mathcal{D}_k$

The next result is stated in [2].

**Theorem 3.1.** *For all positive integers  $a$  and  $b$ , we have  $\mathcal{D}_{a+b} \subseteq \mathcal{D}_a \oplus \mathcal{D}_b$ .* ■

From this theorem it follows that, for all positive integers  $c$  and  $n$ ,  $\mathcal{D}_{cn} \subseteq c\mathcal{D}_n$ . We now show that this cannot be improved, even if we restrict the graphs to be bipartite.

**Theorem 3.2.** *For all positive integers  $c$  and  $n$ ,  $\mathcal{D}_{cn+1} \cap \mathcal{O}^2 \not\subseteq c\mathcal{D}_n$ .*

**Proof.** Let  $t = (n+1)c^{cn+1}$ . Clearly,  $G = K_{cn+1,t} \in \mathcal{D}_{cn+1} \cap \mathcal{O}^2$ . We show that  $G \notin c\mathcal{D}_n$ : Suppose, to the contrary, that  $\{E_1, E_2, \dots, E_c\}$  is a  $c\mathcal{D}_n$ -decomposition of  $E(G)$ . Let  $V_1 = \{v_1, v_2, \dots, v_{cn+1}\}$  be the partite set of order  $cn+1$  and  $V_2$  the partite set of order  $t$ . Consider the edges incident with  $v_1$ . At least  $t/c$  of them must be in the same colour class, hence there is a subset  $U_1$  of  $V_2$  with  $|U_1| = t/c$  such that all edges in  $G[U_1 \cup V_1]$  incident with  $v_1$  have the same colour. Similarly, there is a subset  $U_2$  of  $U_1$  with  $|U_2| = t/c^2$  such that all edges in  $G[U_2 \cup V_1]$  incident with  $v_2$  have the same colour (not necessarily the same as for  $v_1$ ). Continuing in this way we obtain a subset  $U$  of  $V_2$  with  $|U| = n+1$  such that, for every  $v \in V_1$ , all edges of  $G[U \cup V_1]$  incident with  $v$  have the same colour.

Since there are  $c$  colours it follows that for some  $i \in \{1, 2, \dots, c\}$  we have that  $K_{n+1,n+1} \subseteq G[E_i]$ . This is a contradiction, since  $K_{n+1,n+1} \notin \mathcal{D}_n$ . Thus  $K_{cn+1,t} \notin c\mathcal{D}_n$ . ■

**Theorem 3.3.** *For all positive integers  $k$  and  $n$ , we have that*

$$\rho'_{\mathcal{D}_n}(\mathcal{D}_k) = \left\lceil \frac{k}{n} \right\rceil.$$

**Proof.** It follows from Theorem 3.1, by induction on  $c$ , that  $\mathcal{D}_{cn} \subseteq c\mathcal{D}_n$  for all  $c$  and  $n$ . Now let  $k$  and  $n$  be positive integers and let  $c = \left\lceil \frac{k}{n} \right\rceil$ . Then  $k \leq cn$  hence  $\mathcal{D}_k \subseteq \mathcal{D}_{cn} \subseteq c\mathcal{D}_n$  and the upper bound follows.

For the lower bound, since  $k \geq (c-1)n+1$  we have that  $\mathcal{D}_k \supseteq \mathcal{D}_{(c-1)n+1} \not\subseteq (c-1)\mathcal{D}_n$  by Theorem 3.2. ■

We know that if  $pq > a+b$ , then  $\mathcal{D}_{a+b} \subseteq \mathcal{O}^{a+b+1} \subseteq \mathcal{O}^{pq} = \mathcal{O}^p \oplus \mathcal{O}^q$  and  $\mathcal{D}_{a+b} \subseteq \mathcal{D}_a \oplus \mathcal{D}_b$ . Our next result shows that for graphs in  $\mathcal{D}_{a+b}$  we can find simultaneous  $(\mathcal{O}^p, \mathcal{O}^q)$ - and  $(\mathcal{D}_a, \mathcal{D}_b)$ -partitions. First a set-theoretic lemma.

**Lemma 3.4.** *Let  $a, b, p$  and  $q$  be positive integers such that  $a \geq b$ ,  $2 \leq q \leq b+1$  and  $pq > a+b$ . If  $X$  is a set with  $a+b$  elements and  $\{U_1, U_2, \dots, U_p\}$  and  $\{V_1, V_2, \dots, V_q\}$  are partitions of  $X$  then there exists a partition  $\{A, B\}$  of  $X$  and  $i$  and  $j$  such that  $|A| = a$ ,  $A \cap U_i = \emptyset$  and  $B \cap V_j = \emptyset$ .*

**Proof.** It is sufficient (and necessary) to find  $i$  and  $j$  such that  $U_i \cap V_j = \emptyset$ ,  $|U_i| \leq b$  and  $|V_j| \leq a$ . Let  $k$  be the number of  $U_i$ 's such that  $|U_i| > b$  and  $m$  the number of  $V_j$ 's such that  $|V_j| > a$ . We will show that  $(p-k)(q-m) > c = |X \setminus (\cup\{U_i : |U_i| > b\} \cup \cup\{V_j : |V_j| > a\})|$ . It then follows that among the sets of the required size there is a disjoint pair (there are  $(p-k)(q-m)$  ways to choose a pair  $(U_i, V_j)$  of sets of the required size. Since the  $U_i$ 's are pairwise disjoint and the  $V_j$ 's are pairwise disjoint it would follow that  $c \geq (p-k)(q-m)$  if all such pairs have nonempty intersection). Note that  $m \leq 1$  since  $a \geq b$  and that  $c \leq \min\{a+b-k(b+1), a+b-m(a+1)\}$ . Also,  $k < p$ , for otherwise we get  $a+b = |X| \geq p(b+1) \geq pq$ . We have three cases to consider.

(1)  $m = 0$ : In this case we have  $(p-k)q = pq - kq \geq a+b+1-k(b+1) > c$ .

(2)  $m = 1$  and  $k \leq \frac{a+1}{b+1}$ : We want to show that  $(p-k)(q-1) > b-1$  since  $c \leq b-1$ . If  $q = b+1$  this is clearly true, hence we assume that  $q \leq b$ . We have

$$\begin{aligned} \frac{b-1}{q-1} + kq - a &\leq \frac{a+1}{b+1}q - a + \frac{b-1}{q-1} \\ &= a\left(\frac{q}{b+1} - 1\right) + \frac{b-1}{q-1} + \frac{q}{b+1} \\ &\leq b\left(\frac{q}{b+1} - 1\right) + \frac{b-1}{q-1} + \frac{q}{b+1} \quad \text{since } a \geq b \text{ and } q \leq b \\ &= b\left(\frac{1}{q-1} - 1\right) + q - \frac{1}{q-1} \\ &\leq q\left(\frac{1}{q-1} - 1\right) + q - \frac{1}{q-1} \\ &= 1 \end{aligned}$$

Suppose now that  $(p-k)(q-1) \leq b-1$ . Then we have  $pq \leq \frac{b-1}{q-1}q + kq = b-1 + \frac{b-1}{q-1} + kq - a + a \leq a+b$ , a contradiction.

(3)  $m = 1$  and  $k > \frac{a+1}{b+1}$ : Again we may assume that  $q \leq b$ . We show that  $(p-k)(q-1) > a+b-k(b+1) \geq c$ . We have

$$\begin{aligned}
& -k(b+1) + \frac{a+b-k(b+1)}{q-1} + kq \\
&= \frac{a+b}{q-1} + k\left(q - (b+1) - \frac{b+1}{q-1}\right) \\
&\leq \frac{a+b}{q-1} + \frac{a+1}{b+1}\left(q - (b+1) - \frac{b+1}{q-1}\right) \quad \text{since } q \leq b \\
&= a\left(\frac{q}{b+1} - 1\right) + \frac{q}{b+1} + \frac{b-q}{q-1} \\
&\leq b\left(\frac{q}{b+1} - 1\right) + \frac{q}{b+1} + \frac{b-q}{q-1} \\
&= (q-b)\left(1 - \frac{1}{q-1}\right) \\
&\leq 0
\end{aligned}$$

Suppose now that  $(p-k)(q-1) \leq a+b-k(b+1)$ . Then we have  $pq \leq q\frac{a+b-k(b+1)}{q-1} + kq = a+b-k(b+1) + \frac{a+b-k(b+1)}{q-1} + kq \leq a+b$ . ■

**Theorem 3.5.** *Let  $a$ ,  $b$ ,  $p$  and  $q$  be positive integers such that  $a \geq b$ ,  $2 \leq q \leq b+1$  and  $pq > a+b$ . Then  $\mathcal{D}_{a+b} \subseteq (\mathcal{D}_a \cap \mathcal{O}^p) \oplus (\mathcal{D}_b \cap \mathcal{O}^q)$ .*

**Proof.** Let  $G$  be a counterexample of minimum order and let  $v$  be a vertex of  $G$  of degree at most  $a+b$ . Then  $G-v$  has a  $(\mathcal{D}_a \cap \mathcal{O}^p, \mathcal{D}_b \cap \mathcal{O}^q)$ -decomposition and Lemma 3.4 is exactly what we need to extend this decomposition to  $G$  for a contradiction. ■

These results now put us in a position to refine Theorem 3.3.

**Theorem 3.6.** *For all positive integers  $k$ ,  $n$  and  $p \geq 2$  we have that:*

$$\begin{aligned}
\rho'_{\mathcal{D}_n \cap \mathcal{O}^p}(\mathcal{D}_k) &= \left\lceil \log_p(k+1) \right\rceil, \text{ if } k \leq n, \\
&= \left\lceil \frac{k}{n} \right\rceil, \text{ if } k > n \text{ and } p^2 > 2n, \\
&\leq \left\lceil \log_p(n+1) \right\rceil + \left\lceil \frac{k}{n} \right\rceil - 1, \text{ otherwise.}
\end{aligned}$$

**Proof.** Firstly we note that from Theorem 3.5 it follows that  $\mathcal{D}_{cn} \subseteq \mathcal{D}_{(c-1)n} \oplus (\mathcal{D}_n \cap \mathcal{O}^2) \subseteq \mathcal{D}_{(c-1)n} \oplus (\mathcal{D}_n \cap \mathcal{O}^p)$  for all  $c \geq 2$  and therefore  $\mathcal{D}_{cn} \subseteq \mathcal{D}_{2n} \oplus (c-2)(\mathcal{D}_n \cap \mathcal{O}^p)$ .



Suppose that  $k \leq n$ . Then  $\rho'_{\mathcal{D}_n \cap \mathcal{O}^p}(\mathcal{D}_k) = \rho'_{\mathcal{O}^p}(\mathcal{D}_k) = \lceil \log_p(k+1) \rceil$  by Theorem 2.4.

Now suppose that  $k > n$  and  $p^2 > 2n$ . Then  $\mathcal{D}_{cn} \subseteq \mathcal{D}_{2n} \oplus (c-2)(\mathcal{D}_n \cap \mathcal{O}^p) \subseteq c(\mathcal{D}_n \cap \mathcal{O}^p)$ , using Theorem 3.5 and the fact that  $p^2 > 2n$ . Now  $\mathcal{D}_k \subseteq \mathcal{D}_{\lceil \frac{k}{n} \rceil n} \subseteq \lceil \frac{k}{n} \rceil (\mathcal{D}_n \cap \mathcal{O}^p)$  giving the upper bound. The lower bound follows from Theorem 3.3 and Lemma 1.4.

Suppose that  $p^2 \leq 2n$ . From  $\mathcal{D}_{cn} \subseteq \mathcal{D}_{2n} \oplus (c-2)(\mathcal{D}_n \cap \mathcal{O}^p)$  we get that  $\mathcal{D}_{cn} \subseteq \mathcal{D}_n \oplus (c-1)(\mathcal{D}_n \cap \mathcal{O}^p)$ . Moreover, by Theorem 2.4 we have that  $\mathcal{D}_n \subseteq \mathcal{O}^{n+1} \subseteq \lceil \log_p(n+1) \rceil (\mathcal{D}_n \cap \mathcal{O}^p)$ . Therefore  $\mathcal{D}_k \subseteq \mathcal{D}_{\lceil \frac{k}{n} \rceil n} \subseteq \mathcal{D}_n \oplus \left( \lceil \frac{k}{n} \rceil - 1 \right) (\mathcal{D}_n \cap \mathcal{O}^p) \subseteq \left( \lceil \log_p(n+1) \rceil + \lceil \frac{k}{n} \rceil - 1 \right) (\mathcal{D}_n \cap \mathcal{O}^p)$  giving the desired bound. ■

#### 4. Results on $\mathcal{W}_k^*$ and $\mathcal{W}_k$

It has been conjectured (see e.g. [4]) that the generalized vertex-chromatic number  $\rho_{\mathcal{W}_n}(\mathcal{W}_k)$  equals  $\lceil \frac{k+1}{n+1} \rceil$ . We now consider the similar problems of determining  $\rho'_{\mathcal{W}_n^*}(\mathcal{W}_k^*)$  and  $\rho'_{\mathcal{W}_n}(\mathcal{W}_k)$ .

We will say that two trails in a graph *intersect* if they have a common edge.

**Theorem 4.1.** *For  $a \geq 9$  and  $b \geq 1$  we have  $\mathcal{W}_{\lceil \frac{2a-6}{3} \rceil + b}^* \subseteq \mathcal{W}_a^* \oplus \mathcal{W}_b^*$ .*

**Proof.** Consider any graph  $G$  in  $\mathcal{W}_{\lceil \frac{2a-6}{3} \rceil + b}^*$ . Take  $E_1$  to be a maximal subset of  $E(G)$  such that  $G[E_1]$  is in  $\mathcal{W}_a^*$ . Let  $E_2 = E(G) - E_1$ . Suppose that there is an open trail  $T$  in  $G[E_2]$  of length  $b+1$  and let  $e_1$  and  $e_2$  denote the end-edges of  $T$ . Since  $E_1$  is maximal in  $\mathcal{W}_a^*$  it follows that there is an open trail  $T_1$  of length  $a+1$  in  $G[E_1 \cup \{e_1\}]$  and an open trail  $T_2$  of length  $a+1$  in  $G[E_1 \cup \{e_2\}]$ . Let  $T_{11}$  and  $T_{12}$  denote the trails on either side of  $e_1$  such that  $T_{11} \cup \{e_1\} \cup T_{12} = T_1$ . Similarly, let  $T_{21} \cup \{e_2\} \cup T_{22} = T_2$ . Now suppose, without loss of generality, that  $x = |E(T_{11})| \leq y = |E(T_{12})|$ , so that  $x + y = a$ .

It is easily seen that if  $y \geq \lceil \frac{2a}{3} \rceil + 1$ , then by taking the trail  $T_{12} \cup T$  or  $T_{12} \cup (T - e_1)$ , as the case may be, we get a trail of length at least  $\lceil \frac{2a}{3} \rceil + 1 + b$  and therefore an open trail of length at least  $\lceil \frac{2a}{3} \rceil + 1 + b - 1 \geq \frac{2a-2}{3} + b > \frac{2a-4}{3} + b \geq \lceil \frac{2a-6}{3} \rceil + b$  in  $G$ , a contradiction. Therefore

$\lceil \frac{a}{2} \rceil \leq y \leq \lfloor \frac{2a}{3} \rfloor$ . Moreover, each  $T_{ij}$ ,  $i, j \in \{1, 2\}$  has length at least  $\lfloor \frac{a}{3} \rfloor$ , since  $x = a - y \geq a - \lfloor \frac{2a}{3} \rfloor \geq a - \frac{2a}{3} = \frac{a}{3} \geq \lfloor \frac{a}{3} \rfloor$ .

Note that  $T_{11}$  and  $T_{12}$  are necessarily edge disjoint as are  $T_{21}$  and  $T_{22}$ .  $T_{12}$  must intersect  $T_{21}$  and  $T_{22}$ , otherwise we get an open trail of length at least  $\lceil \frac{a}{2} \rceil + b - 2 + \lfloor \frac{a}{3} \rfloor \geq \frac{a}{2} + \frac{a-2}{3} + b - 2 = \frac{5a-16}{6} + b > \lceil \frac{2a-6}{3} \rceil + b$  in  $G$ ; containing  $T_{12}$ ,  $T - e_1 - e_2$  and  $T_{21}$  or  $T_{22}$ .

In the following, when we say that  $T_{21}$  intersects  $T_{12}$  *first* we mean that there is a trail starting from an end-vertex of  $e_2$ , following  $T_{21}$  and ending with an edge of  $T_{12}$ , containing no edge of  $T_{11}$ . Similarly for  $T_{22}$  intersecting  $T_{12}$  first or  $T_{2i}$  intersecting  $T_{11}$  first. Note that since  $T_{11}$  and  $T_{12}$  are disjoint and  $T_{12}$  intersects  $T_{21}$  and  $T_{22}$ , we must have that  $T_{2i}$ ,  $i \in \{1, 2\}$  intersects one of  $T_{11}$  and  $T_{12}$  first.

Suppose that both  $T_{21}$  and  $T_{22}$  intersect  $T_{12}$  first. Then we obtain an open trail of length at least  $x + b - 1 + \lceil \frac{y}{2} \rceil \geq a - y + \frac{y}{2} + b - 1 \geq a - \frac{1}{2}y - 1 + b \geq a - \frac{1}{2} \lfloor \frac{2a}{3} \rfloor - 1 + b \geq a - \frac{1}{2}(\frac{2a}{3}) - 1 + b = \frac{2a-3}{3} + b > \lceil \frac{2a-6}{3} \rceil + b$  in  $G$ ; containing  $T_{11}$ ,  $T - e_1$  and at least a half of  $T_{12}$ .

Now, suppose that  $T_{21}$  or  $T_{22}$  intersects  $T_{11}$  first, say  $T_{21}$ . Then we obtain an open trail of length at least  $y + \lceil \frac{x}{2} \rceil + b - 2 = y + \lceil \frac{1}{2}(a - y) \rceil + b - 2 \geq y + \frac{a-y}{2} + b - 2 \geq \frac{a}{2} + \frac{1}{2} \lceil \frac{a}{2} \rceil + b - 2 \geq \frac{3a}{4} + b - 2 > \lceil \frac{2a-6}{3} \rceil + b$  in  $G$ ; containing  $T_{12}$ ,  $T - e_1 - e_2$  and at least a half of  $T_{11}$ . ■

We remark that a similar result has been proved for vertex partitions and  $\mathcal{W}_k$  in [5].

**Theorem 4.2.** *For all positive integers  $k$  and  $n \geq 9$ ,  $\rho'_{\mathcal{W}_n^*}(\mathcal{W}_k^*) \leq \lceil \frac{3k}{2n-6} \rceil$ .*

**Proof.** From Theorem 4.1 it follows by induction on  $c$  that  $\mathcal{W}_{c \lceil \frac{2n-6}{3} \rceil}^* \subseteq c\mathcal{W}_n^*$  for all positive integers  $c$  and  $n$ . Now, with  $c = \lceil \frac{3k}{2n-6} \rceil$  we have that  $\mathcal{W}_k^* \subseteq \mathcal{W}_{c \lceil \frac{2n-6}{3} \rceil}^* \subseteq c\mathcal{W}_n^*$ . ■

**Theorem 4.3.** *For all positive integers  $k$  and  $n \geq 2$ ,  $\lfloor \frac{k-2}{n-1} \rfloor + 1 \leq \rho'_{\mathcal{W}_n}(\mathcal{W}_k) \leq 2k$ .*

**Proof.** We first show that  $\mathcal{W}_{2ac+2} \not\subseteq c\mathcal{W}_{2a+1}$  for every positive integer  $c$ : Clearly,  $G = K_{ac+1,t} \in \mathcal{W}_{2ac+2}$  for every  $t$ . Let  $t$  be large and suppose that  $G \in c\mathcal{W}_{2a+1}$ . Let  $\{E_1, E_2, \dots, E_c\}$  be a corresponding decomposition

of  $E(G)$ . As in the proof of Theorem 3.2 we get, if  $t$  is large enough, for some  $i \in \{1, 2, \dots, c\}$  that  $K_{a+1, a+2} \subseteq G[E_i]$ , a contradiction.

Now let  $a = \frac{n-1}{2}$  and  $c = \left\lfloor \frac{k-2}{n-1} \right\rfloor$ . Since  $k \geq 2ac + 2$  we have  $\mathcal{W}_k \supseteq \mathcal{W}_{2ac+2} \not\subseteq c\mathcal{W}_n$ . Therefore  $\rho'_{\mathcal{W}_n}(\mathcal{W}_k) \geq c + 1$ .

For the upper bound we have  $\mathcal{W}_k \subseteq \mathcal{D}_k \subseteq k\mathcal{D}_1 \subseteq 2k\mathcal{W}_2 \subseteq 2k\mathcal{W}_n$  from Theorem 3.3 and the well-known fact that every tree has a  $2(\mathcal{W}_2 \cap \mathcal{D}_1)$  edge decomposition. ■

### Acknowledgement

The authors wish to thank their supervisor, Prof. I. Broere, for his criticism and assistance in the final preparation of this paper.

### References

- [1] M. Borowiecki, I. Broere, M. Frick, P. Mihók and G. Semanišin, *A survey of hereditary properties of graphs*, Discuss. Math. Graph Theory **17** (1997) 5–50.
- [2] M. Borowiecki and M. Hałuszczak, *Decompositions of some classes of graphs*, Report No. IM-3-99 (Institute of Mathematics, Technical University of Zielona Góra, 1999).
- [3] I. Broere and M. J. Dorfling, *The decomposability of additive hereditary properties of graphs*, Discuss. Math. Graph Theory **20** (2000) 281–291.
- [4] I. Broere, M.J. Dorfling, J.E. Dunbar and M. Frick, *A path(ological) partition problem*, Discuss. Math. Graph Theory **18** (1998) 113–125.
- [5] I. Broere, S. Dorfling and E. Jonck, *Generalized chromatic numbers and additive hereditary properties of graphs*, Discuss. Math. Graph Theory **22** (2002) 259–270.
- [6] J. Nešetřil and V. Rödl, *Simple proof of the existence of restricted Ramsey graphs by means of a partite construction*, Combinatorica **1** (2) (1981) 199–202.

Received 13 June 2001

Revised 5 April 2002