

## TREE-LIKE ISOMETRIC SUBGRAPHS OF HYPERCUBES

BOŠTJAN BREŠAR\*

*University of Maribor*  
*FERI, Smetanova 17, 2000 Maribor, Slovenia*  
**e-mail:** bostjan.bresar@uni-mb.si

WILFRIED IMRICH

*Montanuniversität Leoben*  
*A-8700 Leoben, Austria*  
**e-mail:** imrich@unileoben.ac.at

AND

SANDI KLAVŽAR\*

*Department of Mathematics, PEF, University of Maribor*  
*Koroška cesta 160, 2000 Maribor, Slovenia*  
**e-mail:** sandi.klavzar@uni-mb.si

### Abstract

Tree-like isometric subgraphs of hypercubes, or tree-like partial cubes as we shall call them, are a generalization of median graphs. Just as median graphs they capture numerous properties of trees, but may contain larger classes of graphs that may be easier to recognize than the class of median graphs. We investigate the structure of tree-like partial cubes, characterize them, and provide examples of similarities with trees and median graphs. For instance, we show that the cube graph of a tree-like partial cube is dismantlable. This in particular implies that every tree-like partial cube  $G$  contains a cube that is invariant under every automorphism of  $G$ . We also show that weak retractions preserve tree-like partial cubes, which in turn implies that

---

\*Supported by the Ministry of Education, Science and Sport of Slovenia under the grants Z1-3073, and 0101-P-504, respectively.

every contraction of a tree-like partial cube fixes a cube. The paper ends with several Frucht-type results and a list of open problems.

**Keywords:** isometric embeddings, partial cubes, expansion procedures, trees, median graphs, graph automorphisms, automorphism groups, dismantlable graphs.

**2000 Mathematics Subject Classification:** 05C75, 05C12, 05C05, 05C25.

## 1. Introduction

By a *partial cube* we mean an isometric subgraph of a hypercube. One of the most important subclasses of the class of partial cubes is the class of median graphs. They possess rich structure and have many interesting properties, cf. [18]. There is also an injection of the class of triangle-free graphs into the class of median graphs of diameter 4, see [16].

Unfortunately, just a small portion of the properties of median graphs extends to partial cubes. For instance, every median graph contains a cube that is invariant under all automorphisms of  $G$  [4]. Clearly, this result is not generally true in the class of partial cubes, the simplest example being  $C_6$ . Also, regular median graphs are precisely hypercubes [20], whereas the class of regular partial cubes seems to be rather rich, cf. [17]. Thus one can pose the following question: Is there a natural class of graphs between median graphs and partial cubes that captures “most” of the properties of median graphs?

This question is also interesting from the algorithmic point of view, that is, from the point of view of the complexity of recognizing classes of graphs. The fastest known algorithms that decide whether a given graph  $G$  is a partial cube have time complexity  $O(mn)$ , where  $m$  is the number of edges and  $n$  the number of vertices of  $G$ , see [1, 15]. In the case of median graphs the recognition complexity is that of recognizing triangle-free graphs, which is currently only slightly better than  $O(m\sqrt{n})$ , cf. [13, 16, 15]. It is believed that this recognition complexity cannot be significantly improved.

On the other hand, it may be possible to find a nontrivial class of partial cubes that contains all median graphs but can be recognized faster. Tree-like partial cubes that we introduce in this paper may be a step in that direction. For a class of partial cubes that are not median graphs but can be recognized faster see [6].

In the next section we introduce the main concept of the paper—tree-like partial cubes. In Section 3 we characterize them as the graphs containing no so-called gated periphery-free subgraph. This in particular implies that one can use *any* sequence of peripheral contractions to obtain  $K_1$  from a tree-like partial cube. We follow with a section in that we list several properties that are shared by median graphs and tree-like partial cubes. In particular we show that hypercubes are the only regular tree-like partial cubes and that the cube graph of a tree-like partial cube is dismantlable. Then, in Section 5, we first deduce from the latter result that every tree-like partial cube  $G$  contains a cube that is invariant under every automorphism of  $G$ . We continue by proving that weak retractions preserve tree-like partial cubes and deduce from this result that every nonexpansive map of a tree-like partial cube fixes a cube. In the last section we list some more properties of tree-like partial cubes and give a list of open problems. Along the way we obtain new and shorter proofs of several results on median graphs.

## 2. Notation and Preliminaries

The *Cartesian product*  $G \square H$  of graphs  $G$  and  $H$  is the graph with vertex set  $V(G) \times V(H)$  in which the vertex  $(a, x)$  is adjacent to the vertex  $(b, y)$  whenever  $ab \in E(G)$  and  $x = y$ , or  $a = b$  and  $xy \in E(H)$ . The Cartesian product of  $k$  copies of  $K_2$  is a *hypercube* or *k-cube*  $Q_k$ ; for short also called *cube*. A graph  $G$  is called *prime* (with respect to the Cartesian product) if it cannot be represented as the product of two nontrivial graphs, that is, if  $G = G_1 \square G_2$  implies that  $G_1$  or  $G_2$  is the one-vertex graph  $K_1$ .

A subgraph  $H$  of  $G$  is called *isometric* if  $d_H(u, v) = d_G(u, v)$  for all  $u, v \in V(H)$ , where  $d_G(u, v)$  denotes the length of a shortest  $u, v$ -path. A subgraph  $H$  of a graph  $G$  is *convex* if for any two vertices  $u, v$  of  $H$  all shortest paths between  $u$  and  $v$  in  $G$  are already in  $H$ . A subgraph  $H$  of a graph  $G$  is called *gated* in  $G$  if for every  $x \in V(G)$  there exists a vertex  $u$  in  $H$  such that  $u$  lies on a shortest  $x, v$ -path for all  $v \in V(H)$ . Clearly gated subgraphs are convex and convex subgraphs isometric. Isometric subgraphs of hypercubes are called *partial cubes*. By the above, convex and gated subgraphs of partial cubes are also partial cubes.

A graph  $G$  is a *median graph* if there exists a unique vertex  $x$  to every triple of vertices  $u, v$ , and  $w$  such that  $x$  lies simultaneously on a shortest  $u, v$ -path, a shortest  $u, w$ -path, and a shortest  $w, v$ -path. Median graphs are partial cubes, cf. [15, 21].

Two edges  $e = xy$  and  $f = uv$  of a graph  $G$  are in the Djoković-Winkler [11, 29] relation  $\Theta$  if  $d_G(x, u) + d_G(y, v) \neq d_G(x, v) + d_G(y, u)$ . Winkler [29] proved that a bipartite graph is a partial cube if and only if  $\Theta$  is transitive.

Let  $G = (V, E)$  be a connected, bipartite graph and  $ab$  an edge of  $G$ . Then the following sets are of relevance for partial cubes:

$$\begin{aligned} W_{ab} &= \{w \in V \mid d_G(a, w) < d_G(b, w)\}, \\ U_{ab} &= \{w \in W_{ab} \mid w \text{ has a neighbor in } W_{ba}\}, \\ F_{ab} &= \{e \in E \mid e \text{ is an edge between } W_{ab} \text{ and } W_{ba}\}. \end{aligned}$$

Clearly,  $W_{ab}$  and  $W_{ba}$  are disjoint and  $V = W_{ab} \cup W_{ba}$  because  $G$  is bipartite.

Let  $G = (V, E)$  be a graph,  $V_1$  and  $V_2$  subsets of  $V$  with nonempty intersection, and  $V = V_1 \cup V_2$ . Suppose that  $\langle V_1 \rangle$  and  $\langle V_2 \rangle$  are isometric in  $G$  and that no vertex of  $V_1 \setminus V_2$  is adjacent to a vertex of  $V_2 \setminus V_1$ . ( $\langle W \rangle$  stands for the subgraph induced by  $W$ .) Then the *expansion* of a graph  $G$  with respect to  $V_1$  and  $V_2$  is the graph obtained from  $G$  by the following procedure:

- (i) Replacement of each vertex  $v \in V_1 \cap V_2$  by vertices  $v_1, v_2$  and insertion of the edge  $v_1v_2$ .
- (ii) Insertion of edges between  $v_1$  and the neighbors of  $v$  in  $V_1 \setminus V_2$  as well as between  $v_2$  and the neighbors of  $v$  in  $V_2 \setminus V_1$ .
- (iii) Insertion of the edges  $v_1u_1$  and  $v_2u_2$  whenever  $v, u \in V_1 \cap V_2$  are adjacent in  $G$ .

If  $V_1 \cap V_2$  is convex in  $G$ , we speak of a *convex expansion*, if  $V_1 \cap V_2$  is isometric in  $G$ , then the expansion is called *isometric*. *Contraction* is the operation inverse to the expansion. Partial cubes were characterized as graphs that can be obtained from  $K_1$  by a sequence of expansions [9].

An expansion is called *peripheral* if at least one of the covering sets  $V_1$  or  $V_2$  is equal to  $V(G)$ . In this case the other set equals the intersection, which is thus necessarily isometric in  $G$ . Now, a graph is called a *tree-like partial cube* if it can be obtained by a sequence of peripheral expansions from  $K_1$ . Finally, a subset  $U_{ab}$  is called a *periphery* if  $U_{ab} = W_{ab}$ . The corresponding  $\Theta$ -class  $F_{ab}$  will be called *periphery inducing*.

### 3. Characterization of Tree-Like Partial Cubes

Median graphs can be characterized as graphs that can be obtained from  $K_1$  by a sequence of convex expansions [19, 21]. Moreover, by [22, Lemma 9],

these expansions can be assumed to be peripheral. Hence, by definition, tree-like partial cubes extend this important property of median graphs.

It is obvious that every tree-like partial cube has a periphery. This is not true for partial cubes in general. For instance, the even cycles  $C_{2n}$ ,  $n \geq 3$ , or the graph of Figure 1 are partial cubes without a periphery. We call such graphs *periphery-free* partial cubes. Another example of periphery-free partial cubes are graphs obtained from complete graphs by subdivision of every edge, see [17].




Figure 1: A periphery-free partial cube

We will characterize tree-like partial cubes via periphery-free subgraphs. It is easy to prove the following proposition.

**Proposition 3.1.** *Let  $G$  be a periphery-free partial cube, and  $H$  be obtained from  $G$  by an expansion. Then  $H$  is periphery-free if and only if the expansion is not peripheral.*

We can then characterize the new class of graphs as follows:

**Theorem 3.2.** *A partial cube is tree-like if and only if it contains no gated periphery-free subgraph.*

**Proof.** Let  $G$  be a smallest tree-like partial cube that contains a gated periphery-free subgraph. Let  $H$  be a periphery-free graph isomorphic to a gated subgraph  $G_1$  of  $G$ , and  $G$  be obtained by a peripheral expansion from a tree-like partial cube  $G'$ . By minimality,  $G'$  contains no gated periphery-free subgraph.

Suppose that  $F_{ab}$  is a  $\Theta$ -class of  $G$  that intersects the edge-set of  $G_1$ . We claim that  $F_{ab}$  is not periphery inducing. Let  $xy \in F_{ab} \cap E(G_1)$ . Since  $H$  is periphery-free there exists a vertex  $u \in V(H)$  that is in  $W_{xy}$  but not in  $U_{xy}$  with respect to  $H$ . By the convexity of  $G_1$  in  $G$ , the vertex corresponding to  $u$  is also not in  $U_{xy}$  with respect to  $G$ , hence  $F_{xy}$  is not periphery inducing in  $G$ .

Therefore  $G_1$  is contained either in a subgraph of  $G$  corresponding to  $G'$  or in  $G \setminus G'$ . In the first case we infer from the isometry of  $G'$  that  $G_1$  is also gated in  $G'$ , in contradiction to the minimality assumption. In the second case we consider the subgraph  $G_2$  of  $G'$  isomorphic to  $G_1$ , induced by vertices that are matched via the periphery inducing  $\Theta$ -class. We claim that  $G_2$  is gated in  $G'$ . Indeed, the distance from any vertex  $x$  of  $G'$  to a vertex of  $G_1$  is exactly 1 plus the distance from  $x$  to the corresponding vertex of  $G_2$ . Hence the gatedness of  $G_2$  clearly follows from the gatedness of  $G_1$ . Thus  $G_2$  is a periphery-free subgraph of  $G'$ , again in contradiction to the minimality assumption.

For the converse assume that  $G$  is a partial cube that contains no gated periphery-free subgraphs. Then  $G$  is not periphery-free and one can obtain  $G$  by a peripheral expansion from a graph  $G'$ . If  $G'$  would contain a gated periphery-free subgraph  $G_1$  then  $G_1$  would be gated also in  $G$  (we use the same arguments as above by considering the distances of vertices from  $G \setminus G'$  to vertices of  $G'$ ). By induction on the number of vertices we infer that  $G'$  is a tree-like partial cube, and thus also  $G$ . ■

The theorem directly implies that convex subgraphs of tree-like partial cubes are tree-like partial cubes. In particular we infer the following.

**Corollary 3.3.** *For any periphery  $U$  of a tree-like partial cube  $G$ ,  $G \setminus U$  is a tree-like partial cube.*

This means that one can use any sequence of peripheral contractions to obtain  $K_1$  from a tree-like partial cube. Note that this is a generalization of the elimination procedure in trees where pendant vertices in trees are contracted (or removed). This justifies the name “tree-like” partial cubes.

**Corollary 3.4.**  *$G \square H$  is a tree-like partial cube if and only if  $G$  and  $H$  are tree-like partial cubes.*

**Proof.**  $G$  and  $H$  are isometric (in fact, even convex) subgraphs of  $G \square H$ . Hence  $G \square H$  is a partial cube if and only if  $G$  and  $H$  are partial cubes.

Moreover, the  $\Theta$ -classes of  $G \square H$  naturally correspond to the  $\Theta$ -classes of  $G$  and  $H$ , cf. [15, Lemma 4.3]. Therefore, if  $G$  and  $H$  are tree-like, then so is  $G \square H$ . For the converse we use induction on  $|V(G)| + |V(H)|$  combined with Corollary 3.3. ■

## 4. More Properties of Tree-Like Partial Cubes

By definition tree-like partial cubes can be obtained from  $K_1$  by a sequence of peripheral expansions, just as median graphs can be obtained from  $K_1$  by a sequence of peripheral (convex) expansions [22]. In this section we list several additional properties that can be extended from median graphs to tree-like partial cubes. We begin with the characterization of regular tree-like partial cubes.

**Theorem 4.1.** *Regular tree-like partial cubes are hypercubes.*

**Proof.** Let  $G$  be a regular tree-like partial cube and suppose that it is not a hypercube. If  $G$  is isomorphic to  $K_2 \square U$  for some peripheral subgraph  $U$ , then by Corollary 3.4,  $U$  is also a regular tree-like partial cube that is not a hypercube, and induction completes the proof. On the other hand, if for some peripheral subgraph  $U$ ,  $G$  is not isomorphic to  $K_2 \square U$ , then  $K_2 \square U$  is a proper induced subgraph of  $G$ . But then  $G$  clearly cannot be regular. ■

Since Cartesian products of regular partial cubes are regular partial cubes, the problem of characterizing regular partial cubes reduces to partial cubes that are prime with respect to the Cartesian product. By the same idea as above we can easily prove the following corollary:

**Corollary 4.2.** *Let  $G$  be a regular partial cube on at least three vertices that is prime with respect to the Cartesian product. Then  $G$  is periphery-free.*

Intersection graphs of maximal hypercubes will be briefly called *cube graphs*. Thus  $H$  is the cube graph of a graph  $G$ , in symbols  $H = Q(G)$ , when the vertices of  $H$  are the maximal hypercubes of  $G$ , two vertices in  $H$  being adjacent whenever the corresponding hypercubes in  $G$  intersect. It was observed by Bandelt and van de Vel [5] that the cube graph of a median graph is always *Helly*, that is a graph in that balls have the Helly property. We cannot extend this property to tree-like partial cubes, as the example in Figure 2 shows.




Figure 2: A tree-like partial cube and its cube graph

On the other hand, Helly graphs belong to the class of dismantlable graphs that are defined by an elimination procedure, that is a generalization of the elimination of simplicial vertices in chordal graphs. We say that a vertex  $u$  in a graph  $G$  is *dominated* by its neighbor  $v$  if all neighbors of  $u$  except  $v$  are also neighbors of  $v$ . If  $G$  can be reduced to the one-vertex graph by successive removal of dominated vertices then  $G$  is called a *dismantlable graph*. Dismantlable graphs were studied in [24] under the name *cop-win graphs*, see also [10]. Below we show that the cube graph of a tree-like partial cube is dismantlable.

Let  $G^\Delta$  denote the graph obtained from a graph  $G$  that has the same vertex set as  $G$  and in that two vertices are adjacent whenever they are in the same hypercube of  $G$  (this operation is from [5]).

**Theorem 4.3.** *For any tree-like partial cube  $G$ , the graphs  $G^\Delta$  and  $Q(G)$  are dismantlable.*

**Proof.** To see that  $G^\Delta$  is dismantlable, let  $U$  be the subgraph of  $G$  obtained in the last expansion step. Then each vertex  $u$  in  $U$  has a unique neighbor  $u'$  in  $G \setminus U$ . It is clear that every maximal hypercube that includes  $u$  includes also  $u'$ , thus  $u$  is a dominated vertex in  $G^\Delta$ . By removing all dominated vertices of  $U$  one by one we obtain the graph  $G \setminus U$  that is a smaller tree-like partial cube than  $G$ , and we can conclude the argument by induction (of course,  $K_1^\Delta = K_1$ , which is dismantlable by definition).

The cube graph of a partial cube  $G$  coincides with the clique graph of  $G^\Delta$  (where the *clique graph* of a graph is the intersection graph of maximal



complete subgraphs of a graph). Bandelt and Prisner [3] showed that dismantlable graphs are invariant under the clique graph transformation, that is, the clique graph of a dismantlable graph is again dismantlable. Combining the above two observations we conclude that the cube graph of a tree-like partial cube is dismantlable. ■

## 5. Mappings of Tree-Like Partial Cubes

Recall that any tree  $T$  contains either a vertex or an edge that is invariant under every automorphism of  $T$ , cf. [23]. This property extends to median graphs in the sense that automorphisms of a median graph always fix a cube [4].

Similarly, automorphisms of dismantlable graphs always fix a complete subgraph. Indeed, just observe that the set of all dominated vertices is invariant under automorphisms of a dismantlable graph, and by simultaneously removing all these vertices the resulting graph is again dismantlable, so the argument follows by induction.

Observe that each automorphism of a tree-like partial cube  $G$  induces an automorphism of  $Q(G)$ . By Theorem 4.3,  $Q(G)$  is dismantlable, so it contains a complete subgraph  $K$  invariant under all automorphisms of  $Q(G)$ . Vertices of  $K$  are pairwise intersecting hypercubes of  $G$ , and their intersection is a hypercube that is invariant under all automorphisms of  $G$ .

**Corollary 5.1.** *Let  $G$  be a tree-like partial cube. Then  $G$  contains a hypercube that is invariant under every automorphism of  $G$ .*

Recall that a *retraction* of  $G$  is a homomorphism  $r$  from  $G$  to a subgraph  $H$  of  $G$  such that  $r|_H$  is the identity. The subgraph  $H$  is called a *retract* of  $G$ . It is always an isometric subgraph of  $G$ . One of the most important characterizations of median graphs says that they are precisely retracts of hypercubes [2]. This result holds also for *weak retractions* where we allow that  $r$  collapses edges to vertices. Alternatively, a weak retraction is an idempotent homomorphism if  $G$  is considered as a reflexive graph, that is a graph in which every vertex carries a loop. As hypercubes are median graphs, one direction of these results is that (weak) retracts of median graphs are median graphs. We can show this for tree-like partial cubes as well.

**Theorem 5.2.** *Every weak retract of a tree-like partial cube is a tree-like partial cube.*

**Proof.** The proof is by induction on the number of vertices of a tree-like partial cube. Let  $r : G \rightarrow H$  be a weak retraction of a tree-like partial cube  $G$  onto  $H$ , and  $U$  a periphery of  $G$ . If  $H \subseteq U$  then  $H = r(G) = r(U)$ . Applying Theorem 3.2 we infer that  $U$  is also a tree-like partial cube, and it has fewer vertices than  $G$  thus  $H$  is a tree-like partial cube by induction. In the rest of the proof we assume that  $H \cap (G \setminus U) \neq \emptyset$ .

We consider two cases. For the first case suppose that for any  $u \in U$ , and the unique neighbor  $v$  of  $u$  in  $G \setminus U$ , we have  $u \in H$  implies  $v \in H$ . We claim that then  $H \cap (G \setminus U)$  is a weak retract of  $G \setminus U$ . We define a mapping  $r' : G \setminus U \rightarrow H \cap (G \setminus U)$  as follows. If  $r(x) \in G \setminus U$ , set  $r'(x) = r(x)$ , and if  $r(x) \in U$ , set  $r'(x) = y$  where  $y$  is the unique neighbor of  $r(x)$  in  $G \setminus U$ . Clearly,  $r'$  is a weak retraction because  $r$  is such a mapping, thus  $H \cap (G \setminus U)$  is a tree-like partial cube by induction. Since weak retracts are isometric subgraphs,  $H$  is a partial cube, and it is tree-like because  $U \cap H$  is a periphery in  $H$ .

In the remaining case there exists  $u \in U \cap H$  of which the unique neighbor  $v$  in  $G \setminus U$  is not in  $H$ . Since  $H \cap (G \setminus U) \neq \emptyset$  we may assume that  $u$  is adjacent to  $x \in U \cap H$  of which the unique neighbor  $y \in G \setminus U$  is also in  $H$ . Because  $H$  is isometric in  $G$ , we deduce that  $(W_{vu} \cap W_{vy}) \cap H = \emptyset$  (using that the sets  $W$ , and hence also their intersections, are convex). Therefore  $H = r(G) = r(G \setminus (W_{vu} \cap W_{vy}))$ , and applying Theorem 3.2 we deduce that  $G \setminus (W_{vu} \cap W_{vy})$  is a tree-like partial cube, hence  $H$  is also tree-like a partial cube. ■

This allows an adaption of Corollary 5.1 to nonexpansive maps. First a definition. A mapping  $\varphi : V(G) \rightarrow V(H)$  for which  $d_H(\varphi(u), \varphi(v)) \leq d_G(u, v)$  for any pair of vertices  $u, v \in G$  is called *nonexpansive*.

**Corollary 5.3.** *Every nonexpansive map of a tree-like partial cube fixes a cube.*

**Proof.** Let  $\varphi$  be a nonexpansive map of a tree-like partial cube. Clearly, there is an iterate  $\varphi^r$  such that  $\varphi^{r+1}(G) = \varphi^r(G)$ . Let  $H = \varphi^r(G)$ . Then  $\varphi|_H$  is an automorphism of  $H$  and there is a power  $\varphi^s$  such that  $\varphi^s|_H$  is the identity. Hence any  $\varphi^{is}$ , where  $is > r$ , is a weak retraction of  $G$  with weak retract  $H$ . By Theorem 5.2 and Corollary 5.1  $H$  contains a hypercube invariant under  $\varphi$ . ■

## 6. Concluding Remarks

In this final section we briefly mention some other properties of tree-like partial cubes and pose several questions.

Yet another interesting feature of median graphs are tree-like equalities. Perhaps the most interesting one is the following: Let  $\alpha_i$  be the number of induced  $i$ -cubes in a median graph, then

$$\sum_{i \geq 0} (-1)^i \alpha_i = 1.$$

This result was discovered by Soltan and Chepoi [27] and independently by Škrekovski [26], see also [8] for further generalizations. It is easy to prove that the above equality extends to tree-like partial cubes. We leave the proof to the reader. (Note that for the graph of Figure 1 the corresponding sum is 2.)

Another property of tree-like partial cubes is the following:

**Proposition 6.1.** *Every finite (abstract) group is isomorphic to the automorphism group of some tree-like partial cube.*

**Proof.** Let  $A$  be a finite group. Then, by Frucht's theorem [12],  $A$  is isomorphic to the automorphism group  $\text{Aut}(G)$  of some graph  $G$ . If  $G$  is a tree, we are done. Otherwise, let  $f(G)$  be the graph constructed as follows. Subdivide each edge of  $G$ , add a new vertex  $z$  and connect all original vertices of  $G$  to  $z$ . In the graph  $f(G)$ , let the vertices at level 1 and 2, be the vertices of distance 1, resp. 2, from  $z$ . Thus, the vertices at level 1 correspond to vertices of  $G$ , and vertices at level 2 have degree 2.

We now label the vertices of  $f(G)$  as follows. First label the vertex  $z$  with  $n = |V(G)|$  zeros. Then label its neighbors with strings consisting of  $n - 1$  zeros and with one 1, each in a different position. Finally, vertices at level 2 receive  $n - 2$  zeros and two 1's in the same positions as their two neighbors. It is clear that  $f(G)$  isometrically embeds into  $Q_n$ . Hence  $f(G)$  is a partial cube. In addition, every  $\Theta$ -class of  $f(G)$  is periphery inducing. (Each periphery is isomorphic to  $K_{1,r}$ , where  $r$  is the degree of the corresponding vertex in  $G$ ). We conclude that  $f(G)$  is a tree-like partial cube.

Finally note (cf. [16]) that  $\text{Aut}(G)$  is isomorphic to  $\text{Aut}(f(G))$  and so  $A$  is isomorphic to  $\text{Aut}(f(G))$ . ■

In [16] and [15] it has also been shown that  $f(G)$  is a median graph if and only if  $G$  is triangle free. Since  $f(G)$  is triangle-free, we infer that  $f(f(G))$  is a median graph. We therefore have the following corollary:

**Corollary 6.2.** *Every finite (abstract) group is isomorphic to the automorphism group of a median graph.*

We wish to add that Sabidussi [25] showed that one can impose additional conditions in Frucht's theorem, in particular arbitrary chromatic number. Since two-chromatic graphs are bipartite and since bipartite graphs are triangle-free, there exists a triangle-free graph  $H$  to every group  $A$  such that  $A$  is isomorphic to  $\text{Aut}(H)$ . Our graph  $f(G)$  is a special case of Sabidussi's construction.

We continue with a remark about the hierarchy of partial cubes as introduced in [14] and refined in [7]. According to that classification, tree-like partial cubes belong to the class of graphs obtainable by an isometric expansion procedure and are thus semi-median graphs. However, they are not almost-median in general. To see this, consider the graph obtained by peripherally expanding the (isometric) 6-cycle of the vertex-deleted 3-cube. For an example of an almost-median periphery-free partial cube, see the graph of Figure 1.

We conclude the paper with the following questions.

1. Cube graphs of all standard examples of partial cubes that are not tree-like are not dismantlable. Therefore we ask the following question: *Is a partial cube tree-like precisely when its cube graph is dismantlable?*

2. Automorphisms of periphery-free partial cubes need not preserve a cube. One can easily find such automorphisms of the graph of Figure 1, of even cycles of length at least six, and of subdivisions of complete graphs (the corresponding automorphisms for the latter graphs are derived from the derangement induced automorphisms of complete graphs). Moreover, any vertex transitive partial cube (different from a hypercube) is such an example; for instance the middle level graphs and the permutohedrons.

A natural question arises: *Let  $G$  be a periphery-free partial cube that is minimal with respect to the expansion sequence. Does there exist an automorphism that does not preserve any cube of  $G$ ?*

3. *Is there a (nice) class of tree-like partial cubes, which would play the role of hypercubes in median graphs, in the sense that every tree-like partial cube is a (weak) retract of some graph of this class?*

### Acknowledgment

We wish to thank one of the referees for several intriguing questions and suggestions. In particular the present proof of Corollary 5.1 as well as Theorem 5.2 and Corollary 5.3 were suggested by this referee.

### References

- [1] F. Aurenhammer and J. Hagauer, *Recognizing binary Hamming graphs in  $O(n^2 \log n)$  time*, Math. Systems Theory **28** (1995) 387–396.
- [2] H.-J. Bandelt, *Retracts of hypercubes*, J. Graph Theory **8** (1984) 501–510.
- [3] H.-J. Bandelt and E. Prisner, *Clique graphs and Helly graphs*, J. Combin. Theory (B) **51** (1991) 34–45.
- [4] H.-J. Bandelt, M. van de Vel, *A fixed cube theorem for median graphs*, Discrete Math. **62** (1987) 129–137.
- [5] H.-J. Bandelt and M. van de Vel, *Superextensions and the depth of median graphs*, J. Combin. Theory (A) **57** (1991) 187–202.
- [6] B. Brešar, W. Imrich and S. Klavžar, *Fast recognition algorithms for classes of partial cubes*, Discrete Appl. Math., in press.
- [7] B. Brešar, W. Imrich, S. Klavžar, H.M. Mulder and R. Škrekovski, *Tiled partial cubes*, J. Graph Theory **40** (2002) 91–103.
- [8] B. Brešar, S. Klavžar, R. Škrekovski, *Cubes polynomial and its derivatives*, Electron. J. Combin. **10** (2003) #R3, pp. 11.
- [9] V. Chepoi,  *$d$ -Convexity and isometric subgraphs of Hamming graphs*, Cybernetics **1** (1988) 6–10.
- [10] V. Chepoi, *Bridged graphs are cop-win graphs: an algorithmic proof*, J. Combin. Theory (B) **69** (1997) 97–100.
- [11] D. Djoković, *Distance preserving subgraphs of hypercubes*, J. Combin. Theory (B) **14** (1973) 263–267.
- [12] R. Frucht, *Herstellung von Graphen mit vorgegebener abstrakter Gruppe*, Compositio Math. **6** (1938) 239–250.
- [13] J. Hagauer, W. Imrich and S. Klavžar, *Recognizing median graphs in sub-quadratic time*, Theoret. Comput. Sci. **215** (1999) 123–136.
- [14] W. Imrich and S. Klavžar, *A convexity lemma and expansion procedures for bipartite graphs*, European J. Combin. **19** (1998) 677–685.

- [15] W. Imrich and S. Klavžar, *Product Graphs: Structure and Recognition* (John Wiley & Sons, New York, 2000).
- [16] W. Imrich, S. Klavžar, and H.M. Mulder, *Median graphs and triangle-free graphs*, SIAM J. Discrete Math. **12** (1999) 111–118.
- [17] S. Klavžar and A. Lipovec, *Partial cubes as subdivision graphs and as generalized Petersen graphs*, Discrete Math. **263** (2003) 157–165.
- [18] S. Klavžar and H.M. Mulder, *Median graphs: characterizations, location theory and related structures*, J. Combin. Math. Combin. Comp. **30** (1999) 103–127.
- [19] H.M. Mulder, *The structure of median graphs*, Discrete Math. **24** (1978) 197–204.
- [20] H.M. Mulder, *n-Cubes and median graphs*, J. Graph Theory **4** (1980) 107–110.
- [21] H.M. Mulder, *The Interval Function of a Graph*, (Mathematical Centre Tracts 132, Mathematisch Centrum, Amsterdam, 1980).
- [22] H.M. Mulder, *The expansion procedure for graphs*, in: R. Bodendiek (ed.), *Contemporary Methods in Graph Theory* (Wissenschaftsverlag, Mannheim, 1990) 459–477.
- [23] R. Nowakowski and I. Rival, *Fixed-edge theorem for graphs with loops*, J. Graph Theory **3** (1979) 339–350.
- [24] R. Nowakowski and P. Winkler, *Vertex-to-vertex pursuit in a graph*, Discrete Math. **43** (1983) 235–239.
- [25] G. Sabidussi, *Graphs with given group and given graph-theoretical properties*, Canad. J. Math. **9** (1957) 515–525.
- [26] R. Škrekovski, *Two relations for median graphs*, Discrete Math. **226** (2001) 351–353.
- [27] P.S. Soltan and V. Chepoi, *Solution of the Weber problem for discrete median metric spaces*, (Russian) Trudy Tbiliss. Mat. Inst. Razmadze Akad. Nauk Gruzin. SSR **85** (1987) 52–76.
- [28] C. Tardif, *Prefibers and the Cartesian product of metric spaces*, Discrete Math. **109** (1992) 283–288.
- [29] P. Winkler, *Isometric embeddings in products of complete graphs*, Discrete Appl. Math. **7** (1984) 221–225.

Received 9 January 2002

Revised 11 June 2002