Discussiones Mathematicae Graph Theory 24 (2004) 115–123

# A NOTE ON MINIMALLY 3-CONNECTED GRAPHS \*

VÍCTOR NEUMANN-LARA

Instituto de Matemáticas Universidad Nacional Autónoma de México Ciudad Universitaria, México D.F. 04510

Eduardo Rivera-Campo

Departamento de Matemáticas Universidad Autónoma Metropolitana-Iztapalapa Av. San Rafael Atlixco 186, México D.F. 09340

AND

JORGE URRUTIA

Instituto de Matemáticas Universidad Nacional Autónoma de México Ciudad Universitaria, México D.F. 04510

## Abstract

If G is a minimally 3-connected graph and C is a double cover of the set of edges of G by irreducible walks, then  $|E(G)| \ge 2|C| - 2$ . **Keywords:** minimally 3-connected, walk double cover. **2000 Mathematics Subject Classification:** 05C70, 05C38.

<sup>\*</sup>Partially supported by Conacyt, México.

### 1. INTRODUCTION

A walk  $\alpha$  in a simple graph G is a sequence  $w_0, w_1, \ldots, w_s$  of vertices of G, not necessarily different, such that  $w_{i-1}w_i$  is an edge of G for  $i = 1, 2, \ldots, s$ . An edge e of G is said to be *traversed* in a walk  $\alpha$  if its vertices are consecutive in  $\alpha$ ; an edge may be traversed more than once in a given walk.

A walk  $\alpha$  in a graph G is *irreducible* if  $a \neq b$  for every pair a, b of edges which are traversed consecutively in  $\alpha$ . A set C of irreducible closed walks in a graph G is a *walk double cover* of G if each edge of G is traversed exactly two times, either once in two different walks in C or twice in the same walk in C.

For any simple graph G and any edge e = uv of G we denote by G - ethe graph obtained from G by deleting the edge e, and by  $G \cdot e$  the simple graph obtained from G by identifying the vertices u and v and deleting loops and multiple edges. A *minimally* 3-connected graph is a 3-connected graph G such that, for every edge e of G, the graph G - e is no longer 3-connected.

Whenever possible we follow the terms and notation given in [1]. A wheel  $W_t$  is a graph with t + 1 vertices, obtained from a cycle  $C_t$  with t vertices by adding a new vertex w adjacent to each vertex in  $C_t$ . The cycle  $C_t$  and the vertex w are called the rim and the hub of  $W_t$ , respectively. In this note we prove the following result.

**Theorem 1.1.** Let G be a minimally 3-connected graph with m edges. If C is a walk double cover of G with k walks, then  $m \ge 2k - 2$ . Moreover if  $m \le 2k - 1$ , then G is a planar graph and C is the set of planar faces of G; in particular if m = 2k - 2, then G is a wheel.

### 2. Proof of Theorem 1.1

The following result due to R. Halin [2] will be used in the proof of Theorem 1.1.

**Theorem 2.1.** If e = uv is an edge of a minimally 3-connected graph G with  $\min\{d(u), d(v)\} \ge 4$ , then e lies in no cycle of G of length 3 and  $G \cdot e$  is also minimally 3-connected.

For any graph G and any walk double cover C of G, we denote by m(G) and by k(C) the number of edges of G and the number of walks in C, respectively. **Remark 1.** Let G be a 3-connected graph and C be a walk double cover of G. If two edges uw and wv are consecutive edges in two walks in C, then the degree of w is at least 4.

**Proof of Theorem 1.1.** The smallest 3-connected graph is the wheel  $W_3$  which is planar and has 6 edges. Since each irreducible walk has at least 3 edges, no walk double cover of  $W_3$  has more than 4 walks. Moreover, the only walk double cover of  $W_3$  with 4 walks consists of the planar faces of  $W_3$ .

We proceed by induction assuming  $m \ge 7$  and that the result holds for every minimally 3-connected graph with less than m edges.

If G has an edge e = uv with  $\min\{d(u), d(v)\} \ge 4$ , then by Halin's theorem,  $G \cdot e$  is also minimally 3-connected. Let  $C \cdot e$  denote the set of k walks of  $G \cdot e$  obtained from the walks in C by contracting the edge e.

Also by Halin's theorem, the edge e lies in no cycle of G of length 3; this implies that all walks in  $C \cdot e$  are irreducible. Because C is a walk double cover of G and e is not an edge of  $G \cdot e$ ,  $C \cdot e$  is a walk double cover of  $G \cdot e$ . By induction,  $m(G \cdot e) \geq 2k(C \cdot e) - 2$ ; therefore  $m \geq 2k - 1$ , since  $m(G \cdot e) = m - 1$  and  $k(C \cdot e) = k$ .

If m = 2k - 1, then  $m(G \cdot e) = 2k(C \cdot e) - 2$ ; by induction  $G \cdot e$  is a wheel  $W_t$  and  $C \cdot e$  is the set of planar faces of  $W_t$ . Let x be the vertex of  $W_t$  obtained by identifying u and v. Since u and v have degree at least 4 in G, the vertex x must be the hub of  $W_t$ ; let  $w_0, w_1, \ldots, w_{t-1}$  be the rim of  $W_t$ .

Since e is in no cycle of G of length 3, G is a graph consisting of the cycle  $w_0, w_1, \ldots, w_{t-1}$ , the two adjacent vertices u and v, and one edge joining each vertex  $w_i$  to either u or v.

Suppose there are distinct integers a, b and c such that  $w_a, w_{b+1}$  and  $w_c$  are adjacent to u in G and  $w_{a+1}, w_b$  and  $w_{c+1}$  are adjacent to v in G. The walks  $w_a, x, w_{a+1}, w_b, x, w_{b+1}$  and  $w_c, x, w_{c+1}$  lie in C, since they are faces of  $G \cdot e$ . This implies that  $w_a, u, v, w_{a+1}, w_b, v, u, w_{b+1}$  and  $w_c, u, v, w_{c+1}$  are walks in C which is not possible, since the edge e = uv cannot lie in three walks in C.

Therefore there are integers i and j such that  $w_i, w_{i+1}, \ldots, w_{j-1}$  are adjacent to u in G and  $w_j, w_{j+1}, \ldots, w_{i-1}$  are adjacent to v in G. This shows that G is a planar graph.

Since  $C \cdot e$  is the set of faces of  $G \cdot e = W_t$  and each walk in  $C \cdot e$  is either a walk in C or is obtained from a walk in C by contracting the edge e, the set C must be the set of faces of G. We can now assume that each edge of G has at least one end with degree 3. If C contains no cycle of length 3, then  $2m \ge 4k$  and  $m \ge 2k$ . Therefore we can also assume that C contains at least one cycle of length 3. Let  $C_3$  be the set of cycles in C of length 3; two cases are considered.

Case 1. There is a cycle  $\alpha$  in  $C_3$  such that no pair of edges of  $\alpha$  are traversed consecutively in any other walk in C.

Let u, v and w be the vertices of  $\alpha$ . Since each edge of G has an end with degree 3, without loss of generality, we can assume  $d_G(u) = d_G(v) = 3$ . Let  $u_1$  and  $v_1$  denote the third vertex of G adjacent to u and the third vertex of G adjacent to v, respectively; notice that  $u_1 \neq v_1$ , since G is 3-connected and has at least 5 vertices.

Subcase 1.1. If  $d_G(w) = 3$ , let  $w_1$  denote the third vertex of G adjacent to w; as above  $u_1 \neq w_1 \neq v_1$ . Let G' be the graph obtained from G by contracting the cycle  $\alpha$  to a single point x. We claim that G' can also be obtained from G by a *delta* to *wye* transformation (see Figure 1), and therefore it is also a 3-connected graph.



Figure 1

Since  $d_{G'}(x) = 3$  and  $d_{G'}(z) = d_G(z)$  for each vertex  $z \neq x$  of G', every edge of G' has an end with degree 3; therefore G' is minimally 3-connected.

Let C' be the set of k-1 walks of G' obtained from the walks in  $C \setminus \{\alpha\}$ by contracting the edges uv, vw and wu. Since no pair of edges of  $\alpha$  are consecutive edges in any walk in  $C \setminus \{\alpha\}$ , all walks in C' are irreducible. Moreover, C' is a walk double cover of G', since C is a walk double cover of G and uv, vw and wu are not edges of G'.

By induction  $m(G') \ge 2k(C')-2$ ; hence  $m \ge 2k-1$ , since m(G') = m-3and k(C') = k-1. If m = 2k-1, then m(G') = 2k(C')-2. Again by induction  $G \cdot e$  is a wheel  $W_t$  and C' is the set of planar faces of  $W_t$ . Since x has degree 3 in G', we can assume without loss of generality that x lies in the rim of  $G' = W_t$  and that  $w_1$  is the hub; this implies that G is a graph as in Figure 2 and therefore it is a planar graph in which  $\alpha$  is a face.



Figure 2

Since C' is the set of faces of G' and every walk in C' is either a walk in  $C \setminus \{\alpha\}$  or is obtained from a walk in  $C \setminus \{\alpha\}$  by contracting some of the edges uv, vw and wu, the set C must be the set of planar faces of G.

Subcase 1.2. If  $d_G(w) \ge 4$ , we consider the graph  $G \cdot uv$ . We claim that u and v cannot be contained in a 3-vertex cut of G and, therefore,  $G \cdot uv$  is 3-connected.

Since  $d_{G \cdot uv}(x) = 3$  and  $d_{G \cdot uv}(z) \leq d_G(z)$  for each vertex  $z \neq x$  of  $G \cdot uv$ , every edge of  $G \cdot uv$  has an end with degree 3; therefore  $G \cdot uv$  is minimally 3-connected.

Let  $C \cdot uv$  be the set of k - 1 walks of  $G \cdot uv$  obtained from the walks in  $C \setminus \{\alpha\}$  by contracting the edge uv to a vertex x and substituting each of the edges uw and vw by the edge xw. Each walk in  $C \cdot uv$  is irreducible, because no pair of edges of  $\alpha$  are traversed consecutively in any other walk in C. Since C is a walk double cover of G and uv is not an edge of  $G \cdot uv$ , the set  $C \cdot uv$  is a walk double cover of  $G \cdot uv$ .

By induction  $m(G \cdot uv) \geq 2k(C \cdot uv) - 2$ ; hence  $m \geq 2k - 2$ , since  $m(G \cdot uv) = m - 2$  and  $k(C \cdot uv) = k - 1$ . If  $m \leq 2k - 1$ , then  $m(G \cdot uv) \leq 2k(C \cdot uv) - 1$ ; again by induction,  $G \cdot uv$  is a planar graph and  $C \cdot uv$  is the set of planar faces of  $G \cdot uv$ .

Since  $G \cdot uv$  is 3-connected, there is a planar drawing  $\overline{G \cdot uv}$  of  $G \cdot uv$  in which x is an interior vertex. Let R be the region formed by the three faces of  $\overline{G \cdot uv}$  in which x is a vertex. Since  $w, u_1$  and  $v_1$  lie in the boundary of R

and x is in the interior of R, a planar drawing  $\overline{G}$  of G can be obtained from  $\overline{G \cdot uv}$  by replacing (within the interior of R) the vertex x with two adjacent vertices u and v, and the edges wx,  $u_1x$  and  $v_1x$  with the edges wu, wv,  $u_1u$  and  $v_1v$  as in Figure 3.



Figure 3

Therefore G is a planar graph and  $\alpha$  is a face of G. Furthermore, C is the set of faces of G, since  $C \cdot uv$  is the set of planar faces of  $G \cdot uv$  and each walk in  $C \cdot uv$  is either a walk in  $C \setminus \{\alpha\}$  or is obtained from a walk in  $C \setminus \{\alpha\}$  by contracting the edge uv to the vertex x and substituting each of the edges uw and vw by the edge xw.

If m = 2k - 2, then  $m(G \cdot uv) = 2k(C \cdot uv) - 2$ ; again by induction,  $G \cdot uv$  is a wheel  $W_t$ . Since  $d_{G \cdot uv}(x) = 3$ , we can assume that x lies in the rim of  $G \cdot uv$ .

If w is the hub of  $G \cdot uv$ , then G is the wheel  $W_{t+1}$ , also with hub w. If  $u_1$  is the hub of  $G \cdot uv$ , then G is a graph as in Figure 4. Notice that if t > 3, then  $G - u_1w$  is 3-connected which is not possible since G is minimally 3-connected. Therefore t = 3 and G is the wheel  $W_4$  with hub w. Analogously, if  $v_1$  is the hub of  $G \cdot uv$ , then G is the wheel  $W_4$ .



Figure 4

#### A NOTE ON MINIMALLY 3-CONNECTED GRAPHS

Case 2. For every cycle  $\alpha \in C_3$  there is walk  $\sigma_{\alpha} \neq \alpha$  in C such that two edges of  $\alpha$  are traversed consecutively in  $\sigma_{\alpha}$ .

For this case, we shall prove that the average length of the walks in C is at least 4 and therefore  $2m \ge 4k$  and  $m \ge 2k$ .

For each  $\alpha \in C_3$  let  $u_{\alpha}$ ,  $w_{\alpha}$  and  $v_{\alpha}$  denote the vertices of  $\alpha$ . Without loss of generality we assume that  $u_{\alpha}w_{\alpha}$  and  $w_{\alpha}v_{\alpha}$  are traversed consecutively in  $\sigma_{\alpha}$ . Notice that the walk  $\sigma_{\alpha}$  is uniquely determined since C is a walk double cover of G.

By Remark 1,  $d_G(w_\alpha) \ge 4$ ; therefore  $d_G(u_\alpha) = d_G(v_\alpha) = 3$ , since every edge of G has an end with degree 3. Let  $u'_\alpha$  and  $v'_\alpha$  denote the third vertex of G adjacent to  $u_\alpha$  and the third vertex of G adjacent to  $v_\alpha$ , respectively.

Again by Remark 1, the edges  $w_{\alpha}u_{\alpha}$  and  $u_{\alpha}v_{\alpha}$  are not traversed consecutively in  $\sigma_{\alpha}$ ; therefore  $\sigma_{\alpha}$  must traverse the edge  $u_{\alpha}u'_{\alpha}$ ; analogously  $\sigma_{\alpha}$ traverses the edge  $v_{\alpha}v'_{\alpha}$ . If  $u'_{\alpha} = v'_{\alpha}$ , then  $u_{\alpha}$  and  $v_{\alpha}$  are adjacent only to  $u'_{\alpha} = v'_{\alpha}$ , to  $w_{\alpha}$  and to each other which is not possible since G is a 3connected graph with at least 5 vertices; therefore  $\sigma_{\alpha}$  has length at least 5 for each  $\alpha \in C_3$ . For each  $\tau \in C$  let  $l(\tau)$  denote the length of  $\tau$ .

Consider the equivalence relation in  $C_3$  given by  $\beta \sim \gamma$  if and only if  $\sigma_\beta = \sigma_\gamma$ . For  $\alpha \in C_3$  let  $[\alpha]$  denote the equivalence class of  $\alpha$ .

Let  $\beta$  and  $\gamma$  be two distinct cycles in  $[\alpha]$  and assume, without loss of generality, that the edges  $u_{\beta}w_{\beta}$ ,  $w_{\beta}v_{\beta}$ ,  $u_{\gamma}w_{\gamma}$  and  $w_{\gamma}v_{\gamma}$  are traversed in  $\sigma_{\alpha} = \sigma_{\beta} = \sigma_{\gamma}$  in that relative order. The edges  $u_{\beta}w_{\beta}$  and  $w_{\beta}v_{\beta}$  are not edges of  $\gamma$  since they are traversed in  $\beta$  and in  $\sigma_{\beta} \neq \beta$ ; analogously  $u_{\gamma}w_{\gamma}$ and  $w_{\gamma}v_{\gamma}$  are not edges of  $\beta$ .

Suppose that  $w_{\beta}v_{\beta}$  and  $u_{\gamma}w_{\gamma}$  are traversed consecutively in  $\sigma_{\alpha}$ . Then  $v_{\beta} = u_{\gamma}$  and  $w_{\beta} \neq w_{\gamma}$ , since  $\sigma_{\alpha}$  is an irreducible walk. Moreover,  $u_{\beta} = v_{\gamma}$  since  $d_G(v_{\beta} = u_{\gamma}) = 3$  and  $w_{\beta}$ ,  $w_{\gamma}$ ,  $u_{\beta}$  and  $v_{\gamma}$  are all adjacent to  $v_{\beta} = u_{\gamma}$ . This implies that the vertices  $v_{\beta} = u_{\gamma}$  and  $u_{\beta} = v_{\gamma}$  are adjacent in G only to  $w_{\beta}$ , to  $w_{\gamma}$  and to each other which is not possible since G is 3-connected and has at least 5 vertices.

Therefore, no edges of two distinct cycles in  $[\alpha]$  are traversed consecutively in  $\sigma_{\alpha}$ . This implies that  $\sigma_{\alpha}$  has at least  $3|[\alpha]|$  edges.

By the above arguments

$$\frac{l\left(\sigma_{\alpha}\right)+l\left(\alpha\right)}{2} \ge \frac{5+3}{2} = 4$$

for each  $\alpha \in C_3$  with  $|[\alpha]| = 1$ , and

$$\frac{l(\sigma_{\alpha}) + \sum_{\beta \in [\alpha]} l(\beta)}{|[\alpha]| + 1} \ge \frac{3|[\alpha]| + 3|[\alpha]|}{|[\alpha]| + 1} = \frac{6|[\alpha]|}{|[\alpha]| + 1} \ge 4$$

for each  $\alpha \in C_3$  with  $|[\alpha]| \geq 2$ .

Since all walks in C which are not in  $C_3$  have length at least 4, the average length in C must also be at least 4.

**Corollary 2.2.** Let G be a minimally 3-connected graph with n vertices. If C is a walk double cover of G with k walks, then  $k \leq \frac{3n-4}{2}$ .

**Proof.** Let *m* denote the number of edges in *G*. W. Mader proved in [3] that  $m \leq 3n - 6$ ; by Theorem 1.1,  $k \leq \frac{m+2}{2} \leq \frac{(3n-6)+2}{2} = \frac{3n-4}{2}$ .

**Corollary 2.3.** If G is a minimally 3-connected planar graph with n vertices, then G has at most n faces. Moreover if G has exactly n faces, then G is a wheel.

**Proof.** Since G is 3-connected, its set of faces is a walk double cover. By Theorem 1.1,  $m \ge 2r - 2$ , where m and r are the number of edges and faces of G, respectively. Since n - m + r = 2, it follows  $r \le n$ .

Also by Theorem 1.1, if G is not a wheel, then  $m \ge 2r - 1$ , in which case  $r \le n - 1$ .

**Corollary 2.4.** If G is a minimally 3-connected graph with n vertices embedded in a closed surface S with Euler characteristic  $\chi \neq 2$ , then G has at most  $n - \chi$  faces.

**Proof.** As in Corollary 2.3, the set of faces of G is a walk double cover of G. Since S is not the sphere, C is not the set of planar faces of G. By Theorem 1.1,  $m \ge 2r$ , where m and r are the number of edges and faces of G, respectively. Since  $\chi = n - m + r$ , it follows  $r \le n - \chi$ .

#### References

 J.A. Bondy and U.S.R. Murty, Graph Theory with Applications (North Holland, 1981).

# A NOTE ON MINIMALLY 3-CONNECTED GRAPHS

- R. Halin, Untersuchungen über minimale n-fach zusammenhängende Graphen, Math. Ann. 182 (1969) 175–188.
- W. Mader, Minimale n-fach zusammenhängende Graphen mit maximaler Kantenzahl, J. Reine Angew. Math. 249 (1971) 201–207.

Received 26 February 2002 Revised 13 November 2002