Discussiones Mathematicae Graph Theory 24 (2004) 431–441

# EVEN [a, b]-FACTORS IN GRAPHS

Mekkia Kouider

Laboratoire de Recherche en Informatique UMR 8623 Bât. 490, Université Paris Sud 91405 Orsay, France e-mail: km@lri.fr

AND

PREBEN DAHL VESTERGAARD

Department of Mathematics Aalborg University, Fredrik Bajers Vej 7G DK-9220 Aalborg Øst, Denmark

e-mail: pdv@math.aau.dk

#### Abstract

Let a and b be integers  $4 \le a \le b$ . We give simple, sufficient conditions for graphs to contain an even [a, b]-factor. The conditions are on the order and on the minimum degree, or on the edge-connectivity of the graph.

Keywords: even factor, eulerian, spanning subgraph.2000 Mathematics Subject Classification: 05C70.

## 1. Introduction

We denote by G a graph of order n = |V(G)|. For a vertex x in V(G) let  $d_G(x)$  denote its degree. By  $\delta = \delta(G) = \min\{d_G(x)|x \in V(G)\}$  we denote the minimum degree in G. Let X, Y be an ordered pair of disjoint subsets of V(G), and f, g be mappings from V(G) into N. By e(X, Y) we denote the number of edges with one endvertex in X and the other in Y. By h(X, Y), we denote the number of odd components in  $G - (X \cup Y)$ . A component C

of  $G - (X \cup Y)$  is called odd if  $e(C, Y) + \sum_{c \in V(C)} f(c)$  is an odd number. An even factor of G is a spanning subgraph all of whose degrees are even. If  $g(x) \leq f(x)$  for all x in V(G), by a [g, f]-factor we understand a spanning subgraph F of G satisfying  $g(x) \leq d_F(x) \leq f(x)$ , for all  $x \in V(G)$ .

**Theorem 1** (Lovász' parity [g, f]-factor theorem [13], [3]). Let G be a graph and let g, f be maps from V(G) into the nonnegative integers such that for each  $v \in V(G)$ ,  $g(v) \leq f(v)$  and  $g(v) \equiv f(v) \pmod{2}$ . Then G contains a [g, f]-factor F such that  $d_F(v) \equiv f(v) \pmod{2}$ , for each  $v \in V(G)$ , if and only if, for every ordered pair X, Y of disjoint subsets of V(G)

$$(*) h(X,Y) - \sum_{x \in X} f(x) + \sum_{y \in Y} g(y) - \sum_{y \in Y} d_G(y) + e(X,Y) \le 0.$$

Tutte's f-factor theorem is surveyed in [1]. Let us recall other results on [a, b]-factors. In [7], Kano and Saito proved that, for any nonnegative integers k, r, s, t satisfying  $k \leq r, 1 \leq r, ks \leq rt$ , every graph with degrees in the interval [r, r+s] has a [k, k+t]-factor. Berge, Las Vergnas, and independently Amahashi and Kano, proved for any integer  $b \geq 2$ , that a graph has a [1, b]-factor if and only if b|N(X)| > |X| for all independent vertex sets X of the graph. Kano proved a sufficient condition for a graph to have an [a, b]-factor giving a condition on the sizes |N(X)| for subsets X of V(G) [8]. Cui and Kano generalized Tutte's 1-factor theorem. They consider a map  $f: V(G) \to \{1, 3, 5, \ldots\}$  and call F an odd [1, f]-factor of G if F is a factor of G with  $d_F(v)$  odd and  $d_F(v) \in [1, f(v)]$  for all vertices v in G. They prove that G has an odd [1, f]-factor if and only if G - X has at most  $\sum_{x \in X} f(x)$ components of odd cardinality for any subset  $X \subseteq V(G)$  [5]. Then, Topp and Vestergaard restrict the number of subsets to be considered above, and, as a consequence, proved that a graph of even order n in which no vertex v is the center of an induced  $K_{1,nf(v)+1}$  -star has an odd [1, f]-factor [15]. In [9, 10], Kouider and Maheo prove the existence of connected [a, b] factors in graphs of high degree. For even factors with degrees between 2 and b we establish a sufficient condition in [11].

**Theorem 2.** Let  $b \ge 2$  be an even integer and let G be a 2-edge connected graph with n vertices and with minimum degree  $\delta(G) \ge \min\{3, \frac{2n}{b+2}\}$ . Then G contains an even [2, b]-factor.

We shall now generalize this to even factors with degrees between a and b, where a is an even integer  $\geq 4$ .

### 2. Results

Let  $a, b, a \leq b$ , be even, positive integers. In the inequality (\*), we substitute e(X, Y) by |X||Y|, and derive a sufficient condition for existence of an even [a, b]-factor in G:

(\*\*) 
$$h(X,Y) - b|X| + a|Y| - \delta|Y| + |X||Y| \le 0.$$

We shall prove the following results.

**Theorem 3.** Let a, b be two even integers satisfying  $4 \leq a \leq b$ . Let G be a 2-edge connected graph of order n at least  $\max\{\frac{(a+b)^2}{b}, \frac{3(a+b)}{2}\}$ , and of minimum degree  $\delta$  at least  $\frac{an}{a+b}$ . Then G has an even [a,b]-factor.

**Example 1.** Take even integers a, b such that  $a \ge 12, b = 2a^2$ , let  $\delta = \frac{3a}{2} + 4$ and let G be the graph which consists of 2a - 2 disjoint copies of a complete graph  $K_{\delta+1}$ , each copy joined by one edge to a common vertex y. The order of G is  $n = (\frac{3a}{2} + 5)(2a - 2) + 1 = 3a^2 + 7a - 9$ , and it is easy to see that  $n \ge \max\{\frac{(a+b)^2}{b}, \frac{3}{2}(a+b)\}$ . The minimum degree of G is  $\delta = \frac{3a}{2} + 4$  and the inequality  $\delta \ge \frac{an}{a+b}$  follows from  $\frac{an}{a+b} = \frac{a(3a^2+7a-9)}{a+2a^2} = \frac{3a^2+7a-9}{2a+1} \le \frac{3a^2+7a}{2a} \le \frac{3}{2}a + \frac{7}{2}$ . So G is not 2-edge connected but satisfies all other conditions of Theorem 3. The graph G has no even [a, b]-factor F, because F must contain an edge from y to K, one of the complete graphs  $K_{\delta+1}$ , and the rectriction of F to K should contain exactly one odd vertex, which is impossible.

**Example 2.** For a positive integer  $k \ge 5$ , let a = 2k + 2 and b = ka. Let n = k(3k + 2) + 1. We consider a graph G of order n, composed of k vertex disjoint copies of the complete graph  $K_{3k+2}$ , and an external vertex  $x_0$  joined to each copy by 3 edges. This graph is 2-edge connected, its minimum degree is  $\delta = 3k \ge \frac{an}{a+b}$ , and  $n \ge \frac{(a+b)^2}{b}$ ,  $n \ge \frac{3b}{2}$ . In an even [a, b]-factor F of G the vertex  $x_0$  must be joined to at least 2k + 2 other vertices, so in F at least one of the  $K_{3k+2}$ 's, say K, is joined to  $x_0$  by exactly 3 edges. Thus the graph K should have a subgraph, namely  $K \cap F$ , with an odd number of odd vertices. Hence G has no even [a, b]-factor.

This example shows that even if G is 3-edge connected the conditions  $\delta \geq \frac{an}{a+b}$  and  $n \geq max\{\frac{(a+b)^2}{b}, 3b/2\}$  are not sufficient for existence of an even [a, b]-factor, even if a is much more smaller than b.

**Theorem 4.** Let  $a \ge 4$  and  $b \ge a$  be two even integers. Let G be a 2-edge connected graph of order  $n \ge \frac{(a+b)^2}{b}$  and of minimum degree at least  $\frac{an}{a+b} + \frac{a}{2}$ . Then G has an even [a, b]-factor.

In the following result, we have a weaker condition on the order, but a stronger one on the edge-connectivity.

**Theorem 5.** Let  $a \ge 4$  and  $b \ge a$  be two even integers, and let  $k \ge a + \min\{\sqrt{a}, \frac{b}{a}\}$ . Let G be a k-edge-connected graph of order  $n \ge \frac{(a+b)^2}{b}$  and of minimum degree at least  $\frac{an}{a+b}$ . Then G has an even [a, b]-factor.

**Example 3.** Let a, b, k be integers such that  $b > 3a^2$ , and  $k \le a - 1$ ; furthermore a, b are even and k is odd. We define a k-connected graph G as follows.

Let Y be a set k independant vertices, and consider a family of k + 2complete graphs  $H_i$  for  $1 \leq i \leq k+2$  such that  $H_i = K_{a+2}$  for  $i \leq k+1$ , and  $H_{k+2} = K_{b+3a-(k+1)(a+3)+1}$ . Each  $y \in Y$  is joined to exactly a + 1 vertices, one in  $H_i$  for each i,  $1 \leq i \leq k+1$ , and a-k vertices in  $H_{k+2}$  so that no two vertices of Y have a common neighbour. So  $d_H(y) = a + 1$ , for each  $y \in Y$ . The order n of G is 3a + b. As  $b > 3a^2$ , one can verify that  $\delta \geq \frac{an}{a+b}$ . Thus Gsatisfies all conditions in Theorem 5, except the one on k. Suppose that Ghas an even [a, b]-factor F. Now, let y be any vertex in Y. As  $d_G(y) = a + 1$ and a + 1 is odd, it follows that  $d_F(y) = a$ . Then necessarily, there exists a copy  $H_t$  for some  $t \leq k$  such that  $e_G(Y, H_t) = e_F(Y, H_t)$ . It follows that the restriction of the factor F to  $H_t$  has k odd vertices; as k is odd, that is impossible. So, the graph G has no even [a, b]-factor.

#### 3. Proofs

We shall use Claims 1–4 below for the proof of Theorem 3. First we establish the truth of (\*) for a large class of ordered pairs X, Y.

Let  $\tau(X,Y) = h(X,Y) - b|X| + a|Y| - \sum_{y \in Y} d_G(y) + e(X,Y).$ The hypotheses of Theorem 3 imply that  $\delta \ge \max\{\frac{3a}{2}, a + \frac{a^2}{b}\}.$  EVEN [a, b]-Factors in Graphs

Claim 1. Inequality (\*) holds if  $-b|X| + a|Y| \le 0$ .

**Proof.** Recall, that for any odd component C, b|V(C)| + e(C, Y) is odd; as b is even, that implies  $e(C, Y) \ge 1$ . Hence, between Y and each odd component of  $G - (X \cup Y)$  there is at least one edge, therefore  $h(X, Y) + e(X, Y) \le \sum_{y \in Y} d_G(y)$ , and (\*) follows as  $-b|X| + a|Y| \le 0$ .

Claim 2. Inequality (\*) holds if  $|Y| \ge a + b$ .

**Proof.** Let -b|X| + a|Y| = p. By Claim 1, we may assume p > 0. By definition of h(X, Y), we have  $|X| + |Y| + h(X, Y) \le n$ . Then we obtain

$$|X| = \frac{a|Y| - p}{b} \le \frac{a(n - h(X, Y) - |X|) - p}{b},$$

and thus

$$|X| \le \frac{a(n-h(X,Y)) - p}{a+b}$$

 $\operatorname{So}$ 

$$e(X,Y) \le |X||Y| \le \frac{a(n-h(X,Y))-p}{a+b}|Y|.$$

By hypothesis on  $\delta$  we have

$$-\sum_{y\in Y} d_G(y) \le -\delta|Y| \le -\frac{an}{a+b}|Y|.$$

That yields the inequality

$$\tau(X,Y) \le h(X,Y) + p - \frac{an}{a+b}|Y| + \frac{a(n-h(X,Y)) - p}{a+b}|Y|.$$

So now, since  $|Y| \ge a + b$ , we get

$$\tau(X,Y) \le h(X,Y) + p - \frac{a(h(X,Y) + p)}{a+b}|Y| \le (1-a)(h(X,Y) + p).$$

As  $a \ge 4$  and p > 0, we conclude that  $\tau(X, Y) \le 0$  and (\*) is proven. By Claims 1 and 2 we may henceforth assume  $0 \le \frac{b}{a}|X| < |Y| \le a + b - 1$ . **Proof of Theorem 3.** We assume  $0 \leq \frac{b}{a}|X| < |Y| \leq a + b - 1$  and, following the different values of |Y|, we proceed to prove that  $\tau(X, Y) \leq 0$ . As  $h(X, Y) \leq n - |X| - |Y|$ ,  $\tau(X, Y)$  is bounded as follows:

$$\tau(X,Y) \le h(X,Y) - b|X| + a|Y| - \delta|Y| + |X||Y|$$
$$\le n - (\delta - a + 1)|Y| + |X|(|Y| - b - 1),$$

and therefore, to prove  $\tau(X, Y) \leq 0$  it suffices to prove that

$$(***) n - (\delta - a + 1)|Y| + |X|(|Y| - b - 1) \le 0.$$

 $Case |Y| \ge b + 1.$ 

Let us set

$$\phi(|Y|) = n - (\delta - a + 1)|Y| + \frac{a}{b}|Y|(|Y| - b - 1).$$

As  $|X| < \frac{a}{b}|Y|$ , we see that (\*\*\*) will follow if  $\phi(|Y|) \le 0$ .

**Claim 3.**  $\phi(|Y|) \le 0$ .

**Proof.** For |Y| varying in the interval of integers, [b+1, a+b-1], the maximum value of the parabola  $\phi$  is attained at an endpoint of the interval. In both ends we shall show that  $\phi(|Y|) \leq 0$ .

$$\phi(b+1) = n - (\delta - a + 1)(b+1);$$

and as  $-\delta \leq -\frac{an}{a+b}$ , we get

$$\phi(b+1) \le n\frac{b-ab}{a+b} + ab + a - b - 1.$$

As  $-n \leq -\frac{(a+b)^2}{b}$ , we obtain

$$\phi(b+1) \le (a+b)(1-a) - (b+1)(1-a) = -(1-a)^2 \le 0.$$

At the other endpoint,

$$\phi(a+b-1) = n + \left(-(\delta - a + 1) + \frac{a}{b}(a-2)\right)(a+b-1).$$

As  $\delta \geq \frac{an}{a+b}$ , we get

$$\phi(a+b-1) \le n\frac{2a+b-a^2-ab}{a+b} + (a+b-1)(a^2-2a+ab-b)\frac{1}{b}.$$

Now the inequalities  $n \geq \frac{(a+b)^2}{b}$  and  $2a-a^2+b-ab=-a(a-2)-b(a-1) \leq 0$  imply

$$\phi(a+b-1) \le \frac{2a+b-a^2-ab}{b}(a+b-a-b+1)$$
  
$$\phi(a+b-1) \le \frac{-a(a-2)-b(a-1)}{b} \le 0.$$

This proves Claim 3.

Henceforth we may assume  $|Y| \leq b$  and  $|X| \leq a - 1$ , as  $|X| < \frac{a}{b}|Y|$ .

Let H be the set of odd components C of  $G - (X \cup Y)$ . Then,  $H = H_1 \cup H_2$  where  $H_1$  is the set of the odd components C having e(C, Y) = 1, and  $H_2$  is the set of those for which  $e(C, Y) \ge 3$ . Let us set h = h(X, Y) = |H| and  $h_i = h_i(X, Y) = |H_i|, i = 1, 2$ . So  $h = h_1 + h_2$ .

Claim 4.  $h_1 \leq \frac{n-|Y|}{\delta+1-|X|}$ .

**Proof of Claim 4.** A component C in  $H_1$  has at least two vertices. Otherwise  $C = \{c\}$  and, the degree of the vertex c could be at most |X| + 1; and, as  $|X| \le a - 1$ , then  $d_G(c) \le a$ ; that contradicts  $d_G(c) \ge \delta \ge \frac{3a}{2}$ . So the component C contains a vertex c' not joined to any vertex in Y, and hence having at least  $\delta - |X|$  neighbours in C, therefore  $|C| \ge \delta - |X| + 1$ and we obtain  $h_1 \le \frac{n-|Y|}{\delta+1-|X|}$ .

We continue with the proof of Theorem 3.

 $Case \ |Y| \le b \ \text{and} \ |X| = 0.$  To prove that  $\tau(X,Y) \le 0$  we shall show that

$$h(X,Y) + a|Y| - \sum_{y \in Y} d(y) \le 0.$$

As G has no bridge, and |X| = 0 necessarily  $h_1 = 0$ ,  $h = h_2$  and  $h \le \frac{1}{3} \sum_{y \in Y} d(y)$ . Then

$$\tau(X,Y) \le -\frac{2}{3} \sum_{y \in Y} d(y) + a|Y| \le |Y| \left(a - 2\frac{\delta}{3}\right).$$

As  $\delta \geq \frac{3a}{2}$ , we conclude  $\tau(X, Y) \leq 0$ . From now,  $|Y| \leq b$  and  $|X| \geq 1$ .

Case  $|Y| \leq b$  and  $1 \leq |X| \leq a-1$ . We note that  $\sum_{y \in Y} d(y) \geq e(Y, H) + e(X, Y)$ , and  $e(Y, H) \geq h_1 + 3h_2 = 3h - 2h_1$ , so

$$\begin{array}{ll} 3h & \leq & \displaystyle \sum_{y \in Y} d(y) - e(X,Y) + 2h_1; \\ \\ h & \leq & \displaystyle \frac{\sum_{y \in Y} d(y) - e(X,Y) + 2h_1}{3} \end{array}$$

By Claim 4, then

$$h \le \frac{\sum_{y \in Y} d(y) - e(X, Y)}{3} + \frac{2n}{3(\delta + 1 - |X|)}.$$

Recalling  $\tau(X,Y) = h - b|X| + a|Y| - \sum_{y \in Y} d(y) + e(X,Y)$ , we obtain the following upper bound for  $\tau(X,Y)$ .

$$\tau(X,Y) \le -2\frac{\sum_{y \in Y} d(y) - e(X,Y)}{3} + \frac{2n}{3(\delta + 1 - |X|)} - b|X| + a|Y|.$$

From  $e(X,Y) \leq |X| |Y|$  and  $\sum_{y \in Y} d(y) \geq \delta |Y|$  we obtain

$$\tau(X,Y) \le -2\frac{|Y|\delta}{3} + |X|(\frac{2|Y|}{3} - b) + a|Y| + \frac{2n}{3(\delta + 1 - |X|)}$$

As  $\delta \geq \frac{3a}{2}$ , this gives

$$au(X,Y) \le |X| \left(\frac{2|Y|}{3} - b\right) + \frac{2n}{3(\delta + 1 - |X|)}$$

Inserting  $|Y| \leq b$  yields

$$\tau(X,Y) \le -\frac{b|X|}{3} + \frac{2n}{3(\delta + 1 - |X|)}$$

Then  $\tau$  is strictly positive if and only if

$$b|X| < \frac{2n}{\delta + 1 - |X|};$$

in other words if

$$(****)$$
  $|X|(\delta+1-|X|) < \frac{2n}{b}.$ 

Let us consider the left side of this inequality as a function f(|X|) of |X|. We have assumed  $1 \le |X| \le a - 1 < \delta$ .

For |X| varying in the interval  $[1, \delta]$  the function f has its minimum for |X| = 1 and  $|X| = \delta$ , namely  $f(1) = f(\delta) = \delta$ . Hence inequality (\* \* \*) implies that  $\delta < \frac{2n}{b}$ . As  $\delta \ge \frac{an}{a+b}$ , we should have b(a-2) < 2a. But this does not hold for  $b \ge a \ge 4$ . So we conclude that  $\tau$  is nonpositive, and Theorem 3 is proven.

**Proof of Theorem 4.**  $\delta \geq \frac{a}{b}(a+b) + \frac{a}{2}$  implies  $\delta \geq \frac{a^2}{b} + \frac{3a}{2} \geq \max\left\{\frac{3a}{2}, a + \frac{a^2}{b}\right\}$ , and all arguments, including the argument for the case  $|Y| \leq b$ , can be carried through.

**Proof of Theorem 5.** Claims 1, 2 and 3 still hold with the hypotheses of Theorem 5, so the proof of Theorem 5 begins analogously to that of Theorem 3, and we reach the assumption  $0 \le \frac{b}{a}|X| < |Y| \le b$ . Now, we examine the missing case.

Case  $|Y| \leq b$ .

We know that  $0 \leq |X| \leq a - 1$  (as  $|X| < \frac{a}{b}|Y|$ ). Since G has edgeconnectivity at least k, each component of  $G - (X \cup Y)$  sends at least k - |X| edges to Y, so  $h(X, Y) \leq \frac{\sum_{y \in Y} d(y) - e(X, Y)}{k - |X|}$ . It follows that

$$\tau(X,Y) \leq \frac{\sum_{y \in Y} d(y) - e(X,Y)}{k - |X|} - b|X| + a|Y| + e(X,Y) - \sum_{y \in Y} d(y),$$
  
$$\tau(X,Y) \leq \frac{k - |X| - 1}{k - |X|} (e(X,Y) - \sum_{y \in Y} d(y)) - b|X| + a|Y|.$$

As  $0 \leq |X| \leq a-1$  and k > a we have  $\frac{k-|X|-1}{k-|X|} > 0$  and inserting  $e(X, Y) - \sum_{y \in Y} d(y) \leq |X||Y| - \delta|Y|$  we obtain

$$\tau(X,Y) \leq \frac{k - |X| - 1}{k - |X|} (|X||Y| - \delta|Y|) - b|X| + a|Y|,$$
  
$$\tau(X,Y) \leq |Y|(a - \frac{k - |X| - 1}{k - |X|} \delta) + (\frac{k - |X| - 1}{k - |X|} |Y| - b)|X|$$

The last term is nonpositive, since  $|Y| \leq b$ ; so to have  $\tau(X, Y) \leq 0$  it will suffice that

(i) 
$$\delta \ge a \frac{k - |X|}{k - |X| - 1}.$$

On one hand, as  $\delta \geq k$ , it is sufficient that  $k \geq a \frac{k-|X|}{k-|X|-1}$ ; as  $|X| \leq a-1$ , we see that the inequality (i) is satisfied if  $k \geq a + \sqrt{a}$ , because  $a \frac{k-|X|}{k-|X|-1} = a(1 + \frac{1}{k-|X|-1}) \leq a(1 + \frac{1}{\sqrt{a}}) = a + \sqrt{a} \leq k$ . On the other hand, we have

$$\delta \ge \frac{an}{a+b} \ge a(1+\frac{a}{b}).$$

If  $k \ge a + \frac{b}{a}$ , and as  $|X| \le a - 1$ , it follows that  $k - |X| - 1 \ge \frac{b}{a}$  and  $a\frac{k-|X|}{k-|X|-1} = a(1 + \frac{1}{k-|X|-1}) \le a(1 + \frac{a}{b}) \le \delta$  and hence, also in this case, the inequality (i) is satisfied; and  $\tau(X, Y) \le 0$ . This proves Theorem 5.

440

# References

- J. Akiyama and M. Kano, Factors and factorizations of graphs a survey, J. Graph Theory 9 (1985) 1–42.
- [2] A. Amahashi, On factors with all degrees odd, Graphs and Combin. 1 (1985) 111-114.
- [3] Mao-Cheng Cai, On some factor theorems of graphs, Discrete Math. 98 (1991) 223-229.
- [4] G. Chartrand and O.R. Oellermann, Applied and Algorithmic Graph Theory (McGraw-Hill, Inc., 1993).
- [5] Y. Cui and M. Kano, Some results on odd factors of graphs, J. Graph Theory 12 (1988) 327–333.
- [6] M. Kano, [a, b]-factorization of a graph, J. Graph Theory 9 (1985) 129–146.
- [7] M. Kano and A. Saito, [a, b]-factors of a graph, Discrete Math. 47 (1983) 113–116.
- [8] M. Kano, A sufficient condition for a graph to have [a, b]-factors, Graphs Combin. 6 (1990) 245–251.
- [9] M. Kouider and M. Maheo, Connected (a, b)-factors in graphs, 1998. Research report no. 1151, LRI, (Paris Sud, Centre d'Orsay). Accepted for publication in Combinatorica.
- [10] M. Kouider and M. Maheo, 2 edge-connected [2, k]-factors in graphs, JCMCC 35 (2000) 75–89.
- [11] M. Kouider and P.D. Vestergaard, On even [2, b]-factors in graphs, Australasian J. Combin. 27 (2003) 139–147.
- [12] Y. Li and M. Cai, A degree condition for a graph to have [a, b]-factors, J. Graph Theory 27 (1998) 1–6.
- [13] L. Lovász, Subgraphs with prescribed valencies, J. Comb. Theory 8 (1970) 391–416.
- [14] L. Lovász, The factorization of graphs II, Acta Math. Acad. Sci. Hungar. 23 (1972) 223–246.
- [15] J. Topp and P.D. Vestergaard, Odd factors of a graph, Graphs and Combin. 9 (1993) 371–381.

Received 22 April 2003 Revised 9 October 2003