Discussiones Mathematicae Graph Theory 24 (2004) 485–490

DOMINANT-MATCHING GRAPHS

IGOR' E. ZVEROVICH AND OLGA I. ZVEROVICH

RUTCOR – Rutgers Center for Operations Research, Rutgers University of New Jersey 640 Bartholomew Rd, Piscataway, NJ 08854-8003, USA e-mail: igor@rutcor.rutgers.edu

Abstract

We introduce a new hereditary class of graphs, the dominant-matching graphs, and we characterize it in terms of forbidden induced subgraphs.

Keywords and phrases: domination number, matching number, hereditary class of graphs.

2000 Mathematics Subject Classification: 05C69.

1. Dominant-Covering Graphs

Let G be a graph. The *neighborhood* of a vertex $x \in V(G)$ is the set $N_G(x) = N(x)$ of all vertices in G that adjacent to x. If vertices x and y of G are adjacent (respectively, non-adjacent), we shall use notation $x \sim y$ (respectively, $x \not\sim y$). For disjoint sets $X, Y \subseteq V(G)$, we write $X \sim Y$ (respectively, $X \not\sim Y$) to indicate that each vertex of X is adjacent to each vertex of Y (respectively, no vertex of X is adjacent to a vertex of Y).

A set $D \subseteq V(G)$ is called a *dominating set* in G if $V(G) = N[D] = \bigcup_{d \in D} N[d]$, where $N[d] = N(d) \cup \{d\}$ is the closed neighborhood of d. A *minimum* dominating set in G is a dominating set having the smallest cardinality. This cardinality is the *domination number* of G, denoted by $\gamma(G)$.

A set $C \subseteq V(G)$ is called a *vertex cover* in G if every edge of G is incident to at least one vertex in C. The minimum cardinality of a vertex cover in G is the *vertex covering number* of G, denoted by $\tau(G)$.

Definition 1. A graph G is called a *dominant-covering graph* if $\gamma(H) = \tau(H)$ for every isolate-free induced subgraph H of G.

Many similarly defined classes were characterized in terms of forbidden induced subgraphs by Zverovich [3], Zverovich [4], Zverovich and Zverovich [5], and Zverovich and Zverovich [6]. We give such a characterization for dominant-covering graphs, and then we extend it to dominant-matching graphs.

Theorem 1. A graph G is a dominant-covering graph if and only if G does not contain any of G_1, G_2, \ldots, G_{10} shown in Figure 1 as an induced subgraph.



Figure 1. Forbidden induced subgraphs for dominant-covering graphs.

Proof. Necessity. It is easy to check that the graphs $G_i \in \{G_1, G_2, \ldots, G_{10}\}$ (Figure 1) satisfies $2 = \gamma(G_i) < \tau(G_i)$, and therefore they are not dominant-covering. It follows that no one of them can be an induced subgraph of a dominant-covering graph.

Sufficiency. Let G be a minimal forbidden induced subgraph for the class of all dominant-covering graphs. Suppose that $G \notin \{G_1, G_2, \ldots, G_{10}\}$. By minimality, G does not contain any of G_1, G_2, \ldots, G_{10} as an induced subgraph. Also, each proper induced subgraph of G is a dominant-covering graph, therefore $\gamma(G) < \tau(G)$.

We consider a minimum dominating set D of G such that D covers the maximum possible number of edges of G [among all minimum dominating

sets of G]. If D covers all edges of G, then $\gamma(G) = \tau(G)$, a contradiction. Thus, we may assume that an edge e = uv is not covered by D.

Since D is a dominating set, there exist vertices w and x in D which are adjacent to u and v, respectively. If w = x then $G(u, v, w) \cong G_1$, a contradiction. Therefore $w \neq x$. Moreover, u is non-adjacent to x, and v is non-adjacent to w.

Let $D_u = (D \setminus \{w\}) \cup \{u\}$. We have $|D_u| = |D|$, and D_u covers the edges uv, uw and vx.

Case 1. D_u is not a dominating set.

Suppose that D_u does not dominate a vertex y of G. Since D is a dominating set, y is adjacent to w. Thus, the edge f = yw is covered by D, and it is not covered by D_u .

Case 2. D_u is a dominating set.

Clearly, D_u is a minimum dominating set. The choice of D implies that there exists an edge f which is covered by D and which is not covered by D_u . Obviously, f is incident to the vertex w, i.e., we may assume that f = yw for some vertex $y \notin \{u, v, x\}$.

In both cases, we have obtained that there exists some edge yw covered by D and not covered by D_u . If y is adjacent to u or x, then G contains G_1 or G_2 as an induced subgraph, a contradiction. Hence edge-set of the induced subgraph H = G(u, v, w, x, y) is one of the following:

Variant 1H: $E(H) = \{uv, uw, vx, wy\}$, or Variant 2H: $E(H) = \{uv, uw, vx, wy, vy\}$, or Variant 3H: $E(H) = \{uv, uw, vx, wy, wx\}$, or Variant 4H: $E(H) = \{uv, uw, vx, wy, wx, vy\}$.

Now we consider the set $D_v = (D \setminus \{x\}) \cup \{v\}$. By symmetry, there exists an edge g = zx which is covered by D and which is not covered by D_v . Again, we have four variants for the induced subgraph F = G(u, v, w, x, z):

Variant 1F: $E(H) = \{uv, uw, vx, xz\}$, or Variant 2F: $E(H) = \{uv, uw, vx, xz, uz\}$, or Variant 3F: $E(H) = \{uv, uw, vx, xz, wx\}$, or Variant 4F: $E(H) = \{uv, uw, vx, xz, wx, uz\}$. Note that the vertices y and z may or may not be adjacent. Combinations of Variants 1H, 2H, 3H, 4H and Variants 1F, 2F, 3F, 4F shows that the set $\{u, v, w, x, y, z\}$ induces one of G_3, G_4, \ldots, G_{10} , a contradiction.

2. Dominant-Matching Graphs

The matching number of a graph G is denoted by $\mu(G)$, i.e., $\mu(G)$ is the maximum cardinality of a matching in G.

Proposition 1 (see Lovász and Plummer [1]). $\mu(G) \leq \tau(G)$ for every graph G.

Proposition 2 (Volkmann [2]). $\gamma(G) \leq \mu(G)$ for every graph G without isolated vertices.

Definition 2. A graph G is called a *dominant-matching graph* if $\gamma(H) = \mu(H)$ for every isolate-free induced subgraph H of G.

Note that the class of all graphs such that $\mu(H) = \tau(H)$ for every induced subgraph H of G coincides with the class of all bipartite graphs, see e.g. Minimax König's Theorem in Lovász and Plummer [1]. Now we extend Theorem 1 by characterization of the dominant-matching graphs in terms of forbidden induced subgraphs.

Theorem 2. A graph G is a dominant-matching graph if and only if G does not contain any of G_3, G_4, \ldots, G_{10} (Figure 1) and H_1, H_2, H_3, H_4, H_5 (Figure 2) as an induced subgraph.

Proof. Necessity. It can be directly checked that

- $\gamma(H_i) = 1$ and $\mu(H_i) = 2$ for i = 1, 2, 3,
- $\gamma(H_j) = 2$ and $\mu(H_j) = 3$ for j = 4, 5, and
- $\gamma(G_k) = 2$ and $\mu(G_k) = 3$ for $k = 3, 4, \dots, 10$.

Therefore none of G_3, G_4, \ldots, G_{10} (Figure 1) and H_1, H_2, H_3, H_4, H_5 (Figure 2) can be an induced subgraph of a dominant-matching graph.



Figure 2. Some forbidden induced subgraphs for dominant-matching graphs.

Sufficiency. Suppose that the statement does not hold. We consider a minimal graph ${\cal G}$ such that

- G does not contain any of G_3, G_4, \ldots, G_{10} (Figure 1) and H_1, H_2, H_3, H_4, H_5 (Figure 2) as an induced subgraph, and
- G is not a dominant-matching graph.

The minimality of G means that each proper induced subgraph of G is a dominant-matching graph. If G does not contain both G_1 and G_2 (Figure 1) induced subgraphs, then G is a dominant-covering graph by Theorem 1. Hence $\gamma(G) = \tau(G)$. Proposition 1 and Proposition 2 imply that $\gamma(G) = \mu(G)$, a contradiction to the choice of G.

Thus, it is sufficient to consider two cases where either G_1 or G_2 is an induced subgraph of G. By minimality of G, $\gamma(G) < \mu(G)$, and G is a connected graph.

Case 1. G_1 is an induced subgraph of G.

Since $\gamma(G) < \mu(G)$, $G \neq G_1$. By connectivity of G, there exists a vertex $u \in V(G) \setminus V(G_1)$ that is adjacent to at least one vertex of G_1 . Clearly, the

set $V(G_1) \cup \{u\}$ induces one of H_1, H_2 or H_3 (Figure 2), a contradiction to the choice of G.

Case 2. G_2 is an induced subgraph of G. As before, there exists a vertex $u \in V(G) \setminus V(G_2)$ that is adjacent to at least one vertex of G_2 . We may assume that G has no induced G_1 [see Case 1]. Hence the set $V(G_2) \cup \{u\}$ induces either H_4 or H_5 (Figure 2), a contradiction to the choice of G.

Acknowledgement

We thank the referee for valuable suggestions and remarks.

References

- L. Lovász and M. Plummer, Matching Theory, North-Holland Math. Stud. 121, Annals Discrete Math. 29 (North-Holland Publ. Co., Amsterdam-New York; Akad. Kiadó, Budapest, 1986) xxvii+544 pp.
- [2] L. Volkmann, On graphs with equal domination and edge independence numbers, Ars Combin. 41 (1995) 45–56.
- [3] I.E. Zverovich, k-bounded classes of dominant-independent perfect graphs, J. Graph Theory 32 (1999) 303–310.
- [4] I.E. Zverovich, Perfect connected-dominant graphs, Discuss. Math. Graph Theory 23 (2003) 159–162.
- [5] I.E. Zverovich and V.E. Zverovich, A semi-induced subgraph characterization of upper domination perfect graphs, J. Graph Theory 31 (1999) 29–49.
- [6] I.E. Zverovich and V.E. Zverovich, An induced subgraph characterization of domination perfect graphs, J. Graph Theory 20 (1995) 375–395.

Received 14 July 2003 Revised 17 November 2003