PROBLEMS COLUMN

Discussiones Mathematicae Graph Theory 25 (2005) 217–218

## ESTIMATION OF CUT-VERTICES IN EDGE-COLOURED COMPLETE GRAPHS

## Adam Idzik

Akademia Świętokrzyska 15 Świętokrzyska street, 25–406 Kielce, Poland and Institute of Computer Science

Polish Academy of Sciences 21 Ordona street, 01–237 Warsaw, Poland

e-mail: adidzik@ipipan.waw.pl

All graphs considered here are finite simple graphs, i.e., graphs without loops, multiple edges or directed edges. For a graph G = (V, E), where Vis a vertex set and E is an edge set, we write sometimes V(G) for V and E(G) for E to avoid ambiguity. We shall write  $G \setminus v$  instead of  $G_{V \setminus \{v\}} =$  $(V \setminus \{v\}, E \cap 2^{V \setminus \{v\}})$ , the subgraph induced by  $V \setminus \{v\}$ . A vertex  $v \in V(G)$ is called a *cut-vertex* of G if G is connected and  $G \setminus v$  is not. By a *k-edgecolouring* of a graph we mean any finite partition of the set of its edges into k subsets. A graph (V, E) with a given k-edge-colouring  $(E^1, \dots, E^k)$  $(E^i \cap E^j = \emptyset$  for  $i \neq j; i, j \in \{1, \dots, k\}$  and  $\bigcup_{i \in \{1, \dots, k\}} E^i = E)$  is denoted by  $(V, E^1, \dots, E^k)$ . The graphs  $(V, E^i)$  are called monochromatic subgraphs of  $(V, E^1, \dots, E^k), i \in \{1, \dots, k\}$ . As usual, by  $K_m$  we denote the complete graph with m vertices.

Let  $c(G^i)$  denote the number of cut-vertices of  $G^i$  in a monochromatic subgraph  $G^i = (V, E^i)$  of a k-edge-coloured complete graph  $K_m = (V, E^1, \dots, E^k)$   $(i \in \{1, \dots, k\}).$ 

Given a k-edge-coloured graph  $G = (V, E^1, \dots, E^k)$ , we define  $F^i = E \setminus E^i$ ,  $G^i = (V, E^i)$ ,  $\overline{G}^i = (V, F^i)$ , where  $E = \bigcup_{i \in \{1, \dots, k\}} E^i$  and  $i \in \{1, \dots, k\}$ . Here  $G^i$  is a monochromatic subgraph of G and  $\overline{G}^i$  its complement in G. **Theorem** (Idzik, Tuza, Zhu). Let  $(E^1, \dots, E^k)$  be a k-edge-colouring of  $K_m$   $(k \ge 2, m \ge 4)$ , such that all the graphs  $\bar{G}^1, \dots, \bar{G}^k$  are connected.

- (i) If one of the subgraphs  $G^1, \dots, G^k$  is 2-connected, say  $G^i$ , then  $c(\bar{G}^i) \leq m-2$  and  $c(\bar{G}^j) = 0$  for  $j \neq i$   $(i, j \in \{1, \dots, k\})$ .
- (ii) If none of the graphs  $G^1, \dots, G^k$  is 2-connected, and one of them is connected, say  $G^i$ , then  $c(\overline{G}^i) \leq 2$   $(i \in \{1, \dots, k\})$ .
- (iii) If none of the graphs  $G^1, \dots, G^k$  is 2-connected, and one of them is disconnected, say  $G^i$ , then  $c(\bar{G}^i) \leq 1$   $(i \in \{1, \dots, k\})$ .

**Problem.** Let  $(E^1, \dots, E^k)$  be a k-edge-colouring of  $K_m$   $(k \ge 2, m \ge 4)$ . What is the cardinality of the set of the sum of cut-vertices of  $\overline{G}^i$  in the case none of  $G^i$  is 2-connected and (a) two of  $G^i$  are connected or (b) two of  $G^i$ are disconnected and  $c(\overline{G}^i) = 1$   $(i \in \{1, \dots, k\})$ ?

Observe that in both cases (a) and (b) all the graphs  $\bar{G}^1, \dots, \bar{G}^k$  are connected.

This problem is related to some theorems presented in [1] and [2].

## References

- J. Bosák, A. Rosa and Š. Znám, On decompositions of complete graphs into factors with given diameters, in: P. Erdős and G. Katona, eds., Theory of Graphs, Proceedings of the Colloquium Held at Tihany, Hungary (Academic Press, New York, 1968) 37–56.
- [2] A. Idzik and Z. Tuza, Heredity properties of connectedness in edge-coloured complete graphs, Discrete Math. 235 (2001) 301–306.

Received 21 November 2003