

DECOMPOSITIONS INTO TWO PATHS

ZDZISŁAW SKUPIEŃ

Faculty of Applied Mathematics
AGH University of Science and Technology
al. Mickiewicza 30, 30–059 Kraków, Poland
e-mail: skupien@uci.agh.edu.pl

Abstract

It is proved that a connected multigraph G which is the union of two edge-disjoint paths has another decomposition into two paths with the same set, U , of endvertices provided that the multigraph is neither a path nor cycle. Moreover, then the number of such decompositions is proved to be even unless the number is three, which occurs exactly if G is a tree homeomorphic with graph of either symbol $+$ or \perp . A multigraph on n vertices with exactly two traceable pairs is constructed for each $n \geq 3$. The Thomason result on hamiltonian pairs is used and is proved to be sharp.

Keywords: graph, multigraph, path decomposition, hamiltonian decomposition, traceable.

2000 Mathematics Subject Classification: 05C70, 05C35, 05C38, 05C45.

1. Introduction

Investigations presented in what follows have been inspired by discussion within GRAPHNET [1] in February 2001 on a *two-path conjecture* presented then and there by Ken W. Smith of Central Michigan University. The conjecture says that a connected graph G which is the edge-disjoint union of two paths of length n has at least one more subgraph which is a path of length n . In four days the discussion concluded with a note by Doug West in which he presented a proof (based on Thomason's paper [6]) of the

result which we state below in terms of decompositions. For notation and terminology, see [7].

The above two-path conjecture of Smith brings to mind a result due to Sloane [4]. It answers in the affirmative Shen Lin's question of 1965 whether every 4-regular graph which is hamiltonian decomposable has another hamiltonian cycle. Lin's paper [2] originated related investigations into hamiltonian decompositions.

Recall that a decomposition of a graph G is a collection of edge-disjoint subgraphs whose union is equal to G . A decomposition is called *hamiltonian* (*traceable*) if all decomposition parts are hamiltonian cycles (hamiltonian paths), the decomposition is called a *hamiltonian pair* (*traceable pair*) if the number of parts in question is two. Let $h_2(G)$ and $t_2(G)$ be the *numbers* of respectively hamiltonian and traceable pairs of G , the numbers being 0 if $\Delta(G) > 4$. Similarly, let $p_2(G)$ stand for the *number of decompositions* of G into two nontrivial paths.

In what follows by a graph, G , we mean a multigraph (which is loopless), the phrase *simple graph* is used to emphasize that multiple (or parallel) edges do not appear. The degree of a vertex is the number of incident edges.

Theorem A (D. West). *If G is connected and decomposable into two paths of length k (where $k > 1$), then G is decomposable into a different pair of paths. In particular, one of these two, different from the original pair, has length at least k .*

The following theorem is a part of Thomason's related result.

Theorem B. *Let G be a multigraph with three or more vertices that has a hamiltonian pair. Then the number $h_2(G)$ of hamiltonian decompositions of G is even and at least four. Moreover, for any two edges of G , the number of hamiltonian pairs in which the two edges are in the same part is also even.*

Note that if G has a pair of parallel edges, a simple switch produces the pair of new decomposition parts. Nevertheless, there are large multigraphs with only few (i.e., two) traceable pairs. Our main result follows.

Theorem 1. *Let G be a connected multigraph that is decomposable into two nontrivial paths whose set of endvertices is denoted by U . If G is neither a path nor a cycle, then G is the union of a different pair of edge-disjoint paths with the same set U of endvertices. In fact, the number $p_2(G)$ of such*

decompositions is then even unless G is homeomorphic with the graph of either symbol $+$, or \perp , and then $p_2 = 3$.

Corollary 2. *The number of traceable pairs among multigraphs of order $n \geq 3$ is even.* ■

The following result shows that Thomason's lower bound $h_2(G) \geq 4$ in his theorem above is sharp.

Proposition 3. *For each $n \geq 3$ there are two n -vertex multigraphs M_n and M_n'' which have exactly four hamiltonian pairs and two traceable ones, respectively.*

2. Proofs and Examples

Proof of Theorem 1. Let P and Q denote the original paths whose union is G . Consider the only interesting case that the largest vertex valency, $\Delta(G)$, is 3 or 4 and G is not homeomorphic with $+$ or \perp . Then both paths P and Q are nontrivial and G has three or more edges. Moreover, $2 \leq |U| \leq 4$, the vertices in U have degree 3 or less, and U includes all vertices of G of odd degree (1 or 3). Furthermore, at least two vertices of G are of degree larger than one.

Case 1. G has exactly one vertex of degree bigger than 2 and $|U| = 3$. Then paths P and Q share one endvertex and intersect at another vertex which is of degree 3 or 4 in G . Hence $\delta(G) = 1$. It is easily seen that $p_2(G) = 2$.

Case 2. $|U| = 2$. Then both vertices in U are of degree two in G , G has a vertex of degree 4 and no vertex of odd degree. Let \hat{G} be obtained from G by joining vertices in U by two parallel edges, say e, f . Let G' be the 4-regular homeomorph of \hat{G} (obtained by contracting an edge incident to a degree-2 vertex, one after another until no such an edge remains). Then $p_2(G)$ is equal to the number of hamiltonian decompositions of G' in which a fixed length-2 path P_3 containing the edge e and its neighbor is in one decomposition part. Therefore $p_2(G)$ is even by Theorem B. Moreover, $p_2(G) = h_2(G')/2$.

Case 3. $|U| = 3$ and G has two or more vertices of degree bigger than 2. Hence U comprises a vertex, x , of degree 2 and two vertices, say y, z , of odd

degree. Let $\hat{G} = G + \{xy, xz\}$ and let G' be the 4-regular homeomorph of \hat{G} . Then $p_2(G) = h_2(G') - h'_2$ where h'_2 counts the hamiltonian decompositions of G' such that one part has preimage in \hat{G} containing the path yxz . Hence, by Theorem B, $p_2(G)$ is even.

Case 4. $|U| = 4$. Then all vertices in U are of odd degree (1 or 3) and G has at least two vertices with degrees in the set $\{3, 4\}$. Add to G a new vertex, say w , together with four edges joining w to all vertices in U . Let \hat{G} and G' be the resulting multigraph and its 4-regular homeomorph, respectively. Then two-path decompositions of G (which must keep U fixed) are in one-one correspondence with hamiltonian pairs of G' whence $p_2(G)$ is even. ■

Proof of Proposition 3. Given the cycle C_n with $n \geq 5$ and a path $P_4 = stuv$ contained in C_n , let the multigraph M_n be obtained from the square C_n^2 of C_n by removing the two crossing chords su and tv and by doubling of edges st and uv . Thus M_n is a 4-regular multigraph with two pairs of parallel edges. Note that contracting any two parallel edges of M_n with $n \geq 6$ results in M_{n-1} . Assume that multigraphs M_4 and M_3 are obtained if this contracting is applied to M_5 and then to M_4 , respectively. Hence $M_3 = {}^2K_3$, the doubled triangle. Let M_n'' be obtained from M_n by removing a pair of parallel edges. Hence $M_3'' = {}^2P_3$ and M_4'' is the join of the 2-cycle C_2 and $2K_1$. Assume that, for each $n \geq 4$, notation in M_n'' is chosen so that degree-2 vertices are u and v (or u', v') and 2st are the two parallel edges. Note that there exists a map $M_n'' \mapsto M_{n+1}''$ for $n \geq 4$ in which the degree-2 vertex u with neighbors t and, say, v_1 ($v_1 = s$ if $n = 4$) is removed and replaced by two new vertices, say u', v' , together with four edges $tu', u'v, v'v, v'v_1$. It is enough to show that $t_2(M_n'') = 2$. This equality is easily seen for $n = 3, 4$. Use the map $M_n'' \mapsto M_{n+1}''$ to show by induction on $n \geq 4$ that in each traceable pair of M_n'' the part containing the edge tu is either the v - u section of the cycle C_n or its switching at 2st . ■

3. Concluding Remarks

More examples of multigraphs M on n vertices (inclusive of the above examples M_n'') with the smallest possible nonzero number of traceable pairs $t_2(M) = 2$, and with $|U| = 2$, are given in author's paper [3] for each $n \geq 7$. Then $p_2(M) = 2$, with vertices in U being the only possible endvertices of

decomposition parts. Simple graphs G of each order $n \geq 5$ with $t_2(G) = 4$ and $|U| = 3$ are given in [3], too.

At the other extreme, $t_2(2P_n) = \frac{1}{2}h_2(2C_n) = 2^{n-2}$ and this is not the largest value of t_2 among n -vertex simple graphs. It is a challenging problem to find (good estimates of) the largest value of t_2 (and/or h_2) among simple graphs (or multigraphs) on n vertices.

Acknowledgements

The author thanks both referees for their helpful remarks.

References

- [1] <http://listserv.nodak.edu/archives/graphnet.html>
- [2] S. Lin, *Computer solutions of the traveling salesman problem*, Bell System Tech. J. **44** (1965) 2245–2269.
- [3] Z. Skupień, *Sparse hamiltonian 2-decompositions together with numerous Hamilton cycles*, submitted.
- [4] N.J.A. Sloane, *Hamiltonian cycles in a graph of degree four*, J. Combin. Theory **6** (1969) 311–312.
- [5] K.W. Smith, *Two-path conjecture*, in: [1], Feb. 16, 2001.
- [6] A.G. Thomason, *Hamiltonian cycles and uniquely edge colourable graphs*, in: B. Bollobás, ed., *Advances in Graph Theory* (Proc. Cambridge Combin. Conf., 1977), Ann. Discrete Math. **3** (1978) (North-Holland, Amsterdam, 1978) pp. 259–268.
- [7] D.B. West, *Introduction to Graph Theory* (Prentice Hall, Upper Saddle River, NJ, 1996).
- [8] D. West, in: [1], Feb. 20, 2001.

Received 8 March 2004
Revised 2 November 2004