## ARC-TRANSITIVE AND *s*-REGULAR CAYLEY GRAPHS OF VALENCY FIVE ON ABELIAN GROUPS

Mehdi Alaeiyan

Department of Mathematics Iran University of Science and Technology Narmak, Tehran 16844, Iran e-mail: alaeiyan@iust.ac.ir

### Abstract

Let G be a finite group, and let  $1_G \notin S \subseteq G$ . A Cayley di-graph  $\Gamma = \operatorname{Cay}(G, S)$  of G relative to S is a di-graph with a vertex set G such that, for  $x, y \in G$ , the pair (x, y) is an arc if and only if  $yx^{-1} \in S$ . Further, if  $S = S^{-1} := \{s^{-1} | s \in S\}$ , then  $\Gamma$  is undirected.  $\Gamma$  is conected if and only if  $G = \langle s \rangle$ . A Cayley (di)graph  $\Gamma = \operatorname{Cay}(G, S)$  is called normal if the right regular representation of G is a normal subgroup of the automorphism group of  $\Gamma$ . A graph  $\Gamma$  is said to be arc-transitive, if  $\operatorname{Aut}(\Gamma)$  is transitive on an arc set. Also, a graph  $\Gamma$  is s-regular if  $\operatorname{Aut}(\Gamma)$  acts regularly on the set of s-arcs.

In this paper, we first give a complete classification for arc-transitive Cayley graphs of valency five on finite Abelian groups. Moreover, we classify s-regular Cayley graph with valency five on an abelian group for each  $s \geq 1$ .

**Keywords:** Cayley graph, normal Cayley graph, arc-transitive, *s*-regular Cayley graph.

2000 Mathematics Subject Classification: 05C25, 20B25.

## 1. INTRODUCTION

For a group G, and a subset S of G such that  $1_G \notin S$ , a Cayley graph  $\operatorname{Cay}(G, S)$  of G relative to S is defined as a graph with a vertex set G and edge set E consisting of those ordered pairs (x, y) from G for which  $yx^{-1} \in S$ . If S is symmetric, that is, if  $S^{-1} = \{s^{-1} : s \in S\}$  is equal to S,

then (x, y) is an edge if and only if (y, x) is an edge, and  $\operatorname{Cay}(G, S)$  is said to be undirected. For a finite, simple and undirect graph  $\Gamma$ , we use  $V(\Gamma), E(\Gamma)$ , and  $\operatorname{Aut}(\Gamma)$  to denote its vertex set, edge set and full automorphism group respectively is said to be vertex-transitive and edge-transitive, if  $\operatorname{Aut}(\Gamma)$ acts transitively on  $V(\Gamma)$ , and  $E(\Gamma)$ , respectively. Moreover, for a positive integer s, an s-arc of  $\Gamma$  is an (s + 1)-tuple  $(v_1, v_2, \ldots, v_s)$  of vertices such that  $\{v_{i-1}, v_i\} \in E(\Gamma)$  for  $1 \leq i \leq s$  and if  $s \geq 2$ , then  $v_{i-1} \neq v_{i+1}$  for  $1 \leq i \leq s - 1$ . We call  $\Gamma$  s-arc-transitive, if  $\operatorname{Aut}(\Gamma)$  acts transitively on  $V(\Gamma)$ and on the set of s-arcs; and  $\Gamma$  is called an s-transitive graph if  $\Gamma$  is s-arctransitive but not (s + 1)-arc-transitive. For the case s = 1, we simply use  $A(\Gamma)$  to denote its 1-arc set and call 1-arc-transitive graphs arc-transitive. An arc-transitive graph  $\Gamma$  is said to be s-regular if for any two s-arcs in  $\Gamma$ , there is a unique automorphism of  $\Gamma$  mapping one to the other. Also, an arc-transitive graph  $\Gamma$  is said to be one regular if  $|\operatorname{Aut}(\Gamma)| = |A(\Gamma)|$ .

In [11] Ming-Yao Xu and Jing Xu classified all arc-transitive Cayley graphs of valency at most four on Abelian groups and in [12], M.Y. Xu classified all one-regular circulant graphs of valency 4. Ming-Yao Xu, Hyo-Seob Sim and Young- Gheel Baik [13] classified all arc-transitive circulant graphs and digraphs of order  $p^m$ , where p is an odd prime. For the case m = 1, that is, for the group  $G = Z_p$ , C.Y. Chao [5] gave such a classification for undirected case in 1971. In 1972 Berggen [4] simplified Chao's proof; also Chao and Wells [6] did the same thing for the directed case in 1973. On the other hand, Alspach Conder, Marusic [1] classified all 2-arc-transitive circulant graphs. The purpose of this paper is to investigate arc-transitive Cayley graphs of valency five on an Abelian group, that is, the arc-transitive graphs whose automorphism groups have an Abelian regular subgroup.

The groups- and graph-theoretic notation and terminology are standard; see [1, 2, 7, 10], for example.

We will denote the semi-directed product of group H by K with H.K.

**Theorem 1.1.** Let G be an Abelian group and let S be a subset of G such that  $1_G \notin S$ . Suppose that  $\Gamma = \operatorname{Cay}(G, S)$  is a connected undirected Cayley graph of group G on S.

- (a) Let  $\Gamma$  be non-normal. Then all arc-transitive Cayley graphs  $\Gamma$  with valency five are as follows:
  - (1)  $G = Z_4 \times Z_2^3 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle \times \langle d \rangle$ ,  $S = \{a, a^{-1}, b, c, d\}$ ,  $\Gamma = K_2 \times Q_4 = Q_5$ ,  $\operatorname{Aut}(\Gamma) = S_2 wr S_5$ .

- (2)  $G = Z_4^2 \times Z_2 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle, S = \{a, a^{-1}, b, b^{-1}, c\}, \Gamma = C_4 \times Q_3 = Q_5, \operatorname{Aut}(\Gamma) = S_2 wr S_5.$
- (3)  $G = Z_4 \times Z_2^2 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle, \ S = \{a, a^{-1}, b, c, a^2 b c\}, \ \Gamma = Q_4^d,$ Aut $(\Gamma) = S_2^4.S_5.$
- (4)  $G = Z_6 = \langle a \rangle$ ,  $S = \{a, a^2, a^3, a^4, a^5\}$ ,  $\Gamma = C_3[K_2] = K_6$ , Aut $(\Gamma) = S_6$ .
- (5)  $G = Z_{10} = \langle a \rangle$ ,  $S = \{a, a^3, a^7, a^9, a^5\}$ ,  $\Gamma = K_{5,5}$ ,  $Aut(\Gamma) = S_5 wr S_2$ .
- (6)  $G = Z_6 \times Z_2 = \langle a \rangle \times \langle b \rangle, S = \{a, a^{-1}, a^2b, a^{-2}b, b\}, \Gamma = K_{6,6} 6K_2,$ Aut $(\Gamma) = S_6 \times S_2.$

(b) Let  $\Gamma$  be normal. Then  $\Gamma$  is arc-transitive if one of the following holds:

- (1)  $G = Z_2^4 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle \times \langle d \rangle, \ S = \{a, b, c, d, abcd\}, \ \Gamma = Q_4^d,$ Aut $(\Gamma) = S_2^4.S_5.$
- (2)  $G = Z_2^5 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle \times \langle d \rangle \times \langle e \rangle, S = \{a, b, c, d, e\}, and \Gamma = Q_5,$ Aut $(\Gamma) = S_2 wr S_5.$

The rest of this paper is organized as follows. In Section 2, we give some preliminaries and in Section 3, we prove Theorem 1.1. In the last section, we will classify s-regular Cayley graphs with valency five on an Abelian group for each  $s \ge 1$ .

## 2. PRIMARY ANALYSIS

For a graph  $\Gamma$ , we denote the automorphism group of  $\Gamma$  by Aut( $\Gamma$ ). The following propositions are basic.

**Proposition 2.1.** Let  $\Gamma = Cay(G, S)$  be a Cayley graph of G on S.

- (1) Aut( $\Gamma$ ) contains the right regular representation G, so  $\Gamma$  is vertex- transitive.
- (2)  $\Gamma$  is connected if and only if  $G = \langle S \rangle$ .
- (3)  $\Gamma$  is undirected if and only if  $S^{-1} = S$ .

Let  $\Gamma = \operatorname{Cay}(G, S)$  be a Cayley graph of G on S, and let

$$\operatorname{Aut}(G,S) = \{ \alpha \in \operatorname{Aut}(G) | S^{\alpha} = S \}.$$

Obviously,  $\operatorname{Aut}(\Gamma) \geq G\operatorname{Aut}(G, S)$  write  $A = \operatorname{Aut}(\Gamma)$ . We have,

**Proposition 2.2** [8, 11].

- (1)  $N_A(G) = G.\operatorname{Aut}(G, S).$
- (2)  $A = G.\operatorname{Aut}(G, S)$  is equivalent to  $G \triangleleft A$ .

**Proposition 2.3** [9]. A graph  $\Gamma$  is arc-transitive if it is vertex-transitive and the stabilizer  $G_u$  of a vertex u acts transitively on the neighborhood  $\Gamma_1(u)$  of u in  $\Gamma$ .

**Definition 2.4.** A Cayley graph  $\Gamma = \operatorname{Cay}(G, S)$  is called normal if  $G \trianglelefteq \operatorname{Aut}(\Gamma)$ .

**Proposition 2.5.** Let  $\Gamma = \text{Cay}(G, S)$  be a normal Cayley graph on G relative to S, Then  $\Gamma$  is arc-transitive if and only if Aut(G, S) acts transitively on the neighborhood  $\Gamma_1(1)$  of 1 in  $\Gamma$ .

For the normality of Cayley graphs of valency five on Abelian groups we have the following:

**Theorem 2.6** [3]. Let  $\Gamma = \operatorname{Cay}(G, S)$  be a connected undirected Cayley graph of an Abelian group G on S with valency 5. Then  $\Gamma$  is normal except when one of the following cases holds:

- (1)  $G = Z_2^4 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle \times \langle d \rangle, S = \{a, b, c, d, abc\} and \Gamma = K_2 \times K_{4,4}$ .
- (2)  $G = Z_4 \times Z_2^2 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle, S = \{a, a^{-1}, a^2, b, c\}$  and  $\Gamma = C_4 \times K_4$ .
- (3)  $G = Z_4 \times Z_2^2 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle, S = \{a, a^{-1}, b, c, a^2b\}$  and  $\Gamma = K_2 \times K_{4,4}$ .
- (4)  $G = Z_4 \times Z_2^3 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle \times \langle d \rangle, S = \{a, a^{-1}, b, c, d\}$  and  $\Gamma = K_2 \times Q_4 = Q_5$ .
- (5)  $G = Z_6 \times Z_2^2 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle, S = \{a, a^{-1}, a^3, b, c\} and \Gamma = K_{3,3} \times C_4.$
- (6)  $G = Z_m \times Z_2^2 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle$  with  $m \ge 3, S = \{a, a^{-1}, ab, a^{-1}b, c\}$ and  $\Gamma = K_2 \times C_m[2K_1].$
- (7)  $G = Z_{4m} \times Z_2 = \langle a \rangle \times \langle b \rangle$  with  $m \ge 3, S = \{a, a^{-1}, a^{2m-1}, a^{2m+1}, b\}$ and  $\Gamma = K_2 \times C_m[2K_1].$
- (8)  $G = Z_{10} = \langle a \rangle, S = \{a^2, a^4, a^6, a^8, a^5\}$  and  $\Gamma = K_2 \times K_5$ .
- (9)  $G = Z_{10} \times Z_2 = \langle a \rangle \times \langle b \rangle, S = \{a, a^{-1}, a^3, a^7, b\}, \Gamma = K_2 \times (K_{5,5} 5K_2).$
- (10)  $G = Z_m \times Z_4 = \langle a \rangle \times \langle b \rangle$  with  $m \ge 3, S = \{a, a^{-1}, b, b^{-1}b^2\}$  and  $\Gamma = C_m \times K_4.$

- (11)  $G = Z_m \times Z_6 = \langle a \rangle \times \langle b \rangle$  with  $m \ge 3, S = \{a, a^{-1}, b, b^{-1}, b^3\}$  and  $\Gamma = C_m \times K_{3,3}.$
- (12)  $G = Z_m \times Z_4 \times Z_2 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle$  with  $m \ge 3, S = \{a, a^{-1}, b, b^{-1}, c\}$ and  $\Gamma = C_m \times Q_3$ .
- (13)  $G = Z_2^3 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle, S = \{a, b, c, ab, ac\} and \Gamma = K_2[2K_2].$
- (14)  $G = Z_4 \times Z_2 = \langle a \rangle \times \langle b \rangle, S = \{a, a^{-1}, b, a^2, a^2b\}$  and  $\Gamma = K_2[2K_2].$
- (15)  $G = Z_4 \times Z_2^2 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle, S = \{a, a^{-1}, b, c, a^2 b c\}$  and  $\Gamma = Q_4^d$ .
- (16)  $G = Z_{2m} = \langle a \rangle$  with  $m \ge 3, S = \{a, a^{-1}, a^{m+1}, a^{m-1}, a^m\}$  and  $\Gamma = C_m[K_2]$ .
- (17)  $G = Z_{2m} \times Z_2 = \langle a \rangle \times \langle b \rangle$  with  $m \ge 2, S = \{a, a^{-1}, ab, a^{-1}b, b\}$  and  $\Gamma = C_{2m}[K_2].$
- (18)  $G = Z_{2m} \times Z_2 = \langle a \rangle \times \langle b \rangle$  with  $m \ge 2, S = \{a, a^{-1}, ab, a^{-1}ba^m\}$  and  $\Gamma = C_{2m}^d [2K_1].$
- (19)  $G = Z_{10} = \langle a \rangle, S = \{a, a^3, a^7, a^9, a^5\}$  and  $\Gamma = K_{5,5}$ .
- (20)  $G = Z_6 \times Z_2 = \langle a \rangle \times \langle b \rangle, S = \{a, a^{-1}, a^2b, a^{-2}b, b\}$  and  $\Gamma = K_{6,6} 6K_2$ .
- (21)  $G = Z_{2m} \times Z_4 = \langle a \rangle \times \langle b \rangle$  with  $m \ge 2, S = \{a, a^{-1}, b, b^{-1}, a^m b^2\}$  and  $\Gamma = Q_3 \times C_m$ .
- (22)  $G = Z_{6m} = \langle a \rangle$  with m odd and  $m \geq 3, S = \{a^2, a^{-2}, a^m, a^{5m}, a^{3m}\}$ and  $\Gamma = K_{3,3} \times_c C_m$ .
- (23)  $G = Z_{6m} \times Z_2 = \langle a \rangle \times \langle b \rangle$  with  $m \ge 2, S = \{a, a^{-1}, ba^m, ba^{-m}, ba^{3m}\}$ and  $\Gamma = K_{3,3} \times_c C_{2m}$ .

Let X and Y be two graphs. The direct product  $X \times Y$  is defined as a graph with a vertex set  $V(X \times Y) = V(X) \times V(Y)$  such that for any vertex  $u = [x_1, y_1]$  and  $v = [x_2, y_2]$  in  $V(X \times Y)$ , [u, v] is an edge in  $X \times Y$  whenever  $x_1 = x_2$  and  $[y_1, y_2] \in E(Y)$  or  $y_1 = y_2$  and  $[x_1, x_2] \in E(X)$ . Two graphs are called relatively prime if they have no nontrivial common direct factor. The *lexicographic product* X[Y] is defined as a graph vertex set V(X[Y]) = $V(X) \times V(Y)$  such that for any two vertices  $u = [x_1, y_1]$  and  $v = [x_2, y_2]$ in V(X[Y]), [u, v] is an edge in X[Y] whenever  $[x_1, x_2] \in E(X)$  or  $x_1 = x_2$ and  $[y_1, y_2] \in E(Y)$ . Let  $V(Y) = \{y_1, y_2, \ldots, y_n\}$ . Then there is a natural embedding nX in X[Y], where for  $1 \leq i \leq n$ , the *i*th copy of X is a subgraph induced on the vertex subset  $\{(x, y_i) | x \in V(X)\}$  in X[Y]. The deleted lexicographic product X[Y] - nX is a graph obtained by deleting all the edges of (this natural embedding of) nX from X[Y]. Let  $\Gamma$  be a graph and  $\alpha$  a permutation of  $V(\Gamma)$ , and  $C_n$  a circuit of length n. The *twisted product*  $\Gamma \times_{\alpha} C_n$  of  $\Gamma$  by  $C_n$  with respect to  $\alpha$  is defined by

$$V(\Gamma \times_{\alpha} C_n) = V(\Gamma) \times V(C_n) = \{(x, i) \mid x \in V(\Gamma), i = 0, 1, \dots, n-1\},$$
  

$$E(\Gamma \times_{\alpha} C_n) = \{[(x, i), (x, i+1)] | x \in V(\Gamma), i = 0, 1, \dots, n-2\}$$
  

$$\cup \{[(x, n-1), (x^{\alpha}, o)] \mid x \in V(\Gamma)\}$$
  

$$\cup \{[(x, i), (y, i)] | [x, y] \in E(\Gamma), i = 0, 1, \dots, n-1\}.$$

Now we introduce some graphs which appear in our main theorem. The graph  $Q_4^d$  denotes a graph obtained by connecting all long diagonals of 4cube  $Q_4$ , that is, connecting all vertex u and v in  $Q_4$  such that d(u, v) = 4. The graph  $K_{m,m} \times_c C_n$  is a twisted product of  $K_{m,m}$  by  $C_n$  such that c is a cycle permutation on each part of the complete bipartite graph  $K_{m,m}$ . The graph  $Q_3 \times_d C_n$  is a twisted product of  $Q_3$  by  $C_n$  such that d transposes each pair elements on long diagonals of  $Q_3$ . The graph  $C_{2m}^d[2K_1]$  is defined by:

$$V(C_{2m}^{d}[2K_{1}]) = V(C_{2m}[2K_{1}]),$$
  

$$E(C_{2m}^{d}[2K_{1}]) = E(C_{2m}[2K_{1}])$$
  

$$\cup \{[(x_{i}, y_{j}), (x_{i+m}, y_{j})] \mid i = 0, 1, \dots, m-1, j = 1, 2\},$$

where  $V(C_{2m}) = \{x_0, x_1, \dots, x_{2m-1}\}$  and  $V(2K_1) = \{y_1, y_2\}.$ 

## 3. The Proof of Theorem

In this section, our objective is to show all arc-transitive Cayley graphs of Abelian groups with valency five.

First, we want to show that some cases of Theorem 2.5 are satisfied by Theorem 1.1.

In the following cases we shall assume  $G = \operatorname{Aut}(\Gamma)$ .

In the cases (1) and (3), let  $V(K_2) = \{y_1, y_2\}$  and let  $V(K_{4,4}) = \{x_1, x_2, x_3, x_4, x'_1, x'_2, x'_3, x'_4\}$  such that  $(x_i, x'_j) \in E(K_{4,4})$  for  $1 \le i, j \le 4$ . We obtain that  $f \notin G_{(y_1, x_1)}$  such that  $f(y_2, x_1) = (y_1, x_1)'$ ]), so by Proposition 2.3  $\Gamma$  is not arc-transitive. In the cases (2) and (10), let  $V(C_m) = \{1, 2, 3, \ldots, m\}$  and let  $V(K_4) = \{x_1, x_2, x_3, x_4\}$ . We also obtain that  $f \notin G_{(2,x_1)}$  such that  $f(2, x_4) = (3, x_1)$ , so by Proposition 2.3  $\Gamma$  is not arc-transitive.

In the cases (5) and (11), let  $V(K_{3,3}) = \{x_1, x_2, x_3, x'_1, x'_2, x'_3\}$  and let  $V(C_4) = \{y_1, y_2, y_3, y_4\}$ . We have  $(x_i, x'_j) \in E(K_{3,3})$ , for  $1 \le i, j \le 4$ , and  $(y_i, y_{i+1}) \in E(C_4)$ . We obtain  $f \notin G_{(x_1, y_1)}$  such that  $f(x'_1, y_1) = (x_1, y_2)$ , so by Proposition 2.3  $\Gamma$  is not arc-transitive.

In the cases (6) and (7),  $\Gamma = K_2 \times C_m[2K_1]$  contains two copies X and Y from  $C_m[2K_1]$ . Let  $V(X) = \{x_1, x_2, \ldots, x_m; y_1, y_2, \ldots, y_m \text{ and } V(Y) = \{x'_1, x'_2, \ldots, x'_m; y'_1, y'_2, \ldots, y'_m \text{ such that vertices } x_1, x_2, \ldots, x_m \text{ form a circuit} on a copy of X. We find that <math>f \notin G_{(x_1)}$  such that  $f(x_2) = x'_1$ , so by Proposition 2.3,  $\Gamma$  is not arc-transitive.

In the case (8), we obtained that for  $m \ge 4$ ,  $K_2 \times K_m$  is not arctransitive. Neither is the graph  $K_2 \times K_{4,4}$ .

In the case (9), let  $V(K_{5,5} - 5K_2) = \{x_1, x_2, \dots, x_5, x'_1, x'_2, \dots, x'_5\},$  $V(K_2) = \{y_1, y_2\}$  such that  $(x_i, x'_j) \in E(K_{5,5} - 5K_2)$  for  $i \neq j, 1 \leq i, j \leq 5$ . We find that  $f \notin G_{(y_1, x_1)}$  such that  $f(y_1, x'_2) = (x_1, y_2)$ , so by Proposition 2.3  $\Gamma$  is not arc-transitive.

In the cases (12) for  $[m \neq 4]$  and (21), let  $V(C_m) = \{0, 2, 3, \ldots, m-1\}$ and  $Q_3$  contain two circuits  $C_4, C'_4$  with set of vertices  $V(C_4) = \{x_1, x_2, x_3, x_4\}$  and  $V(C'_4) = \{y_1, y_2, y_3, y_4\}$ , respectively. In addition  $(x_i, x'_i) \in E(Q_3)$  for  $1 \leq i \leq 4$ . We obtained that  $f \notin G_{(x_1,0)}$  such that  $f(x'_1, 0) = (x_1, 0)$ , so by Proposition 2.3  $\Gamma$  is not arc-transitive.

In the cases (13) and (14), let  $V(K_2) = \{x, y\}$  and  $V(2K_2) = \{1, 2, 3, 4\}$ , and also  $E(2K_2)$  contain two edges (1, 2), (3, 4). We find that  $f \notin G_{(y,1)}$  such that f(x, 1) = (y, 2), so by Proposition 2.3  $\Gamma$  is not arc-transitive.

In the case (16) for  $[m \neq 3]$ , let  $V(C_m) = \{1, 2, ..., m\}$  and  $V(K_2) = \{x, y\}$ . We find that  $f \notin \operatorname{Aut}(\Gamma)$  such that f([(2, y), (3, x)]) = [(3, x), (3, y)].

The case (17) is also the special case of (16), since  $2m \neq 3$ .

In the case (18), we find that  $f \notin G_{(x_0,y_2)}$  such that  $f(x_1,y_2) = (x_m,y_2)$ , so by Proposition 2.3  $\Gamma$  is not arc-transitive.

In the cases (22) and (23), let  $V(C_m) = \{0, 1, ..., m-1\}, V(K_{3,3}) = \{x_1, x_2, x_3, x'_1, x'_2, x'_3\}$  and also  $(x_i, x'_j) \in E(K_{3,3})$  for  $1 \le i, j \le 3$ . We find that  $f \notin G_{(x_1,0)}$  such that  $f(x'_1, 0) = (x_1, 1)$ , so by Proposition 2.3  $\Gamma$  is not arc-transitive.

In the case (4),  $\Gamma = K_2 \times Q_4 \simeq C_4 \times Q_3$  and  $Q_3$  is arc-transitive, then by combination of functions we conclude that  $\Gamma$  is arc-transitive.

- In the cases of (6) and (7) the case (12) for [m = 4] is similarly the case (4).
  - In the case (15), we will obtain similarly graph  $\Gamma = Q_4$ .

In the case (16), for m = 3 we have  $\Gamma \simeq K_6$ .

The case (19) is obvious and in the case (20),  $\Gamma = K_{6,6} - 6K_2$ , and we will obtain the same result in graph  $K_{6,6}$ . Thus we complete the proof of Theorem 1.1(a).

For the normal case, since |S| = 5, S contains at least one element of order 2. Since Aut(G, S) is transitive on S all five elements in S are of order 2. Then we have one of the following cases:

- (1)  $G = Z_2^3 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle, S = \{a, b, c, ab, ac\}, \Gamma = K_2[2K_2].$
- (2)  $G = Z_2^4 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle \times \langle d \rangle, S = \{a, b, c, d, abc\}, \Gamma = K_2 \times K_{4,4}.$
- (3)  $G = Z_2^4 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle \times \langle d \rangle, S = \{a, b, c, d, ab\}, \Gamma = K_4 \times C_4.$
- (4)  $G = Z_2^4 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle \times \langle d \rangle, S = \{a, b, c, d, abcd\}, \Gamma = Q_4^d.$
- (5)  $G = Z_2^5 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle \times \langle d \rangle \times \langle e \rangle, S = \{a, b, c, d, e\}, \Gamma = Q_5.$

Note that graphs of the case (1) and (2) are non-normal. In the case (2) of non-normal graphs we showed that graph  $\Gamma = K_4 \times C_4$  is not arc-transitive, and in the case (10) of non-normal graphs we showed that  $\Gamma = Q_4^d$  is arc-transitive. In the final non-normal case we obtained that the graph  $Q_5$  is arc-transitive.

# 4. s-Regular Cayley Graph with Valency Five on Abelian Groups

Let  $\Gamma = \operatorname{Cay}(G, S)$  be a Cayley graph on G with respect to S and let  $A = \operatorname{Aut}(\Gamma)$ . Denote by  $A_1$  the stabilizer of identity 1 of G in A and by  $\operatorname{Aut}(G, S)$  the subgroup of A fixing S setwise. Then we have:

**Theorem 4.1** [14, Proposition 1.5].  $\Gamma$  is normal if and only if  $A_1 = \operatorname{Aut}(G, S)$ .

By noting that all Cayley graphs are vertex-transitive, one can easily prove the following lemma.

**Lemma 4.2.** All s-regular ( $s \ge 1$ ) Cayley graphs are connected.

The following theorem gives a classification of s-regular Cayley graphs with valency five on Abelian groups for each  $s \ge 1$ .

**Theorem 4.3.** Let  $\Gamma$  be an s-regular Cayley graph with valency five on an Abelian group for some  $s \geq 1$ . Then s = 2 or 3. Furthermore,  $\Gamma$  is 2-regular if and only if  $\Gamma$  is isomorphic to  $Q_4^d$ , or  $Q_5$ , or  $K_6$ , or  $K_{6,6} - 6K_2$ ; and is 3-regular if and only if  $\Gamma$  is isomorphic to the complete bipartite graph  $K_{5,5}$ .

**Proof.** Let G be an Abelian group. Assume that  $\Gamma = \operatorname{Cay}(G, S)$  is an s-regular Cayley graph with valency five for some  $s \geq 1$ . Then by Lemma 4.2  $\Gamma$  is connected. By Theorem 1.1, the only non-normal arc-transitive Cayley graphs  $\operatorname{Cay}(G, S)$  are the  $\Gamma_1 = Q_5$ ,  $\Gamma_2 = Q_4^d$ ,  $\Gamma_3 = K_6$ ,  $\Gamma_4 = K_{5,5}$ , and  $\Gamma_5 = K_{6,6} - 6K_2$ . The graphs  $\Gamma_1$ ,  $\Gamma_2$ ,  $\Gamma_3$ , and  $\Gamma_5$  are 2-regular and the graph  $\Gamma_4$  is 3-regular. Thus, we assume the  $\Gamma = \operatorname{Cay}(G, S)$  is normal from now on. Since  $\Gamma$  is of valency 5,  $S = S^{-1}$  contains at least one involution in G. As  $\Gamma$  is arc-transitive, so the group  $\operatorname{Aut}(G, S)$  acts transitive on S. Hence S consists of five involutions. Since S generates the group G, we have  $G = Z_2^3$ , or  $G = Z_2^4$ , or  $G = Z_2^5$ . By Theorem 1.1, the only normal arc-transitive Cayley graphs  $\operatorname{Cay}(G, S)$  with valency 5 are

(1) 
$$G = Z_2^4 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle \times \langle d \rangle, S = \{a, b, c, d, abcd\}, \Gamma = Q_4^d.$$

(2) 
$$G = Z_2^5 = \langle a \rangle \times \langle b \rangle \times \langle c \rangle \times \langle d \rangle \times \langle e \rangle, S = \{a, b, c, d, e\}, \text{ and } \Gamma = Q_5,$$

and each of them is 2-regular.

#### References

- B. Alspach, M. Conder, D. Marusic and Ming-Yao Xu, A classification of 2-arc-transitive circulant, J. Algebraic Combin. 5 (1996) 83–86.
- [2] N. Biggs, Algebraic Graph Theory (Cambridge University Press, 1974).
- [3] Y.G. Baik, Y.Q. Feng and H.S. Sim, The normality of Cayley graphs of finite Abelian groups with valency 5, System Science and Mathematical Science 13 (2000) 420–431.
- [4] J.L. Berggren, An algebraic characterization of symmetric graph with p point, Bull. Aus. Math. Soc. 158 (1971) 247–256.
- [5] C.Y. Chao, On the classification of symmetric graph with a prime number of vertices, Trans. Amer. Math. Soc. 158 (1971) 247–256.
- [6] C.Y. Chao and J. G. Wells, A class of vertex-transitive digraphs, J. Combin. Theory (B) 14 (1973) 246–255.
- [7] J.D. Dixon and B. Mortimer, Permutation Groups (Springer-Verlag, 1996).

- [8] C.D. Godsil, On the full automorphism group of a graph, Combinatorica 1 (1981) 243-256.
- [9] C.D. Godsil and G. Royle, Algebric Graph Theory (Springer-Verlag, 2001).
- [10] H. Wielandt, Finite Permutation Group (Academic Press, New York, 1964).
- [11] Ming-Yao Xu and Jing Xu, Arc-transitive Cayley graph of valency at most four on Abelian Groups, Southest Asian Bull. Math. 25 (2001) 355–363.
- [12] Ming-Yao Xu, A note on one-regular graphs of valency four, Chinese Science Bull. 45 (2000) 2160–2162.
- [13] Ming-Yao Xu, Hyo-Seob Sim and Youg-Gheel Baik, Arc-transitive circulant digraphs of odd prime-power order, (summitted).
- [14] Ming-Yao Xu, Automorphism groups and isomorphisms of Cayley digraphs, Discrete Math. 182 (1998) 309–319.

Received 29 November 2005 Revised 5 June 2006