Discussiones Mathematicae
Graph Theory 27 (2007) 29–38

NEW SUFFICIENT CONDITIONS FOR HAMILTONIAN AND PANCYCLIC GRAPHS*

Ingo Schiermeyer

Fakultät für Mathematik und Informatik
Technische Universität Bergakademie Freiberg
09596 Freiberg, Germany

and

Mariusz Woźniak

Faculty of Applied Mathematics
AGH University of Science and Technology
Mickiewicza 30, 30–059 Kraków, Poland

Abstract

For a graph G of order n we consider the unique partition of its vertex set $V(G) = A \cup B$ with $A = \{v \in V(G) : d(v) \geq n/2\}$ and $B = \{v \in V(G) : d(v) < n/2\}$. Imposing conditions on the vertices of the set B we obtain new sufficient conditions for hamiltonian and pancyclic graphs.

Keywords: hamiltonian graphs, pancyclic graphs, closure.

2000 Mathematics Subject Classification: 05C38, 05C45.

1. Introduction

We use [4] for terminology and notation not defined here and consider finite and simple graphs only.

A graph of order n is called *hamiltonian* if it contains a cycle of length n and is called *pancyclic* if it contains cycles of all lengths from 3 to n.

*The work was partially done while the last author was visiting TU Bergakademie in Freiberg. This stay was partially supported by Deutscher Akademischer Austauschdienst (DAAD).

Let $\omega(G)$ denote the number of components of a graph G. A graph G is called 1-tough if, for every nonempty proper subset S of $V(G)$, we have $\omega(G-S) \leq |S|$.

Various sufficient conditions for a graph to be hamiltonian have been given in terms of vertex degrees. Recall some of them.

Theorem 1 (Dirac [5]). *Let G be a graph of order $n \geq 3$. If $\delta(G) \geq n/2$, then G is hamiltonian.*

Theorem 2 (Ore [9]). *Let G be a graph of order $n \geq 3$. If $d(u)+d(v) \geq n$ for every pair of nonadjacent vertices $u, v \in V(G)$, then G is hamiltonian.*

In [7] Theorem 2 was extended as follows.

Theorem 3 (Flandrin, Li, Marczyk, Woźniak [7]). *Let $G = (V, E)$ be a 2-connected graph on n vertices with minimum degree δ. If $uv \in E(G)$ for every pair of vertices $u, v \in V(G)$ with $d(u) = \delta$ and $d(v) < n/2$, then G is hamiltonian.*

With respect to its vertex degrees, the vertex set of every graph G has a unique partition $V(G) = A \cup B$ with $A = \{v \in V(G) : d(v) \geq n/2\}$ and $B = \{v \in V(G) : d(v) < n/2\}$. In terms of A and B we make the following observations:

- If a graph G satisfies Dirac's condition then $B = \emptyset$.
- If a graph G satisfies Ore's condition, then $G[B]$ is complete and $|B| \leq \delta + 1$.
- If a graph G satisfies the condition of Theorem 3, then $G[B]$ is connected, $G[u \in B : d(u) = \delta]$ is complete and $|B| \leq \delta + 1$.

2. Results

We define three classes of graphs $\mathcal{G}_1$, $\mathcal{G}_2$ and $\mathcal{G}_3$ as follows.

Let $\mathcal{G}_1$ be the class of all 2-connected graphs G such that $uv \in E(G)$ for every pair of vertices $u, v \in B$ with $d(u) = \delta(G)$.

Let $\mathcal{G}_2$ be the class of all 2-connected graphs G such that there exists a vertex $u \in B$ with $d(u) = \delta(G)$ and $uv \in E(G)$ for all vertices $v \in B - \{u\}$.

Let $\mathcal{G}_3$ be the class of all 2-connected graphs G such that $|B| \leq \delta(G)+1$ and $\Delta(G[B]) \geq \min\{2, |B|-1\}$.



Figure 1. Graph $\mathcal{F}_{n,\delta}$

For all n, δ with $2 \leq \delta \leq \frac{n-1}{2}$ define $\mathcal{F}_{n,\delta}$ as a graph of order n, minimum degree δ and vertex set $V(\mathcal{F}_{n,\delta}) = \{u_0, u_1, \ldots, u_\delta, w_1, \ldots, w_{n-\delta-1}\}$ such that $d(u_0) = \delta$, $N(u_0) = \{u_1, \ldots, u_\delta\}$, vertices $u_1, \ldots, u_\delta$ are independent, vertices $w_1, \ldots, w_{n-\delta-1}$ induce a clique and $u_i w_j \in E(G)$ for all $1 \leq i \leq \delta$ and $1 \leq j \leq \delta - 1$. Now, for $S = \{u_0, w_1, \ldots, w_{\delta-1}\}$ we have

$$\omega(\mathcal{F}_{n,\delta} - S) = \delta + 1 > \delta = |S|.$$

Hence, $\mathcal{F}_{n,\delta}$ is not 1-tough and therefore not hamiltonian.

For all n, δ with $2 \leq \delta \leq \frac{n-1}{2}$ define $\mathcal{H}_{n,\delta}$ as a supergraph of $\mathcal{F}_{n,\delta}$ such that $V(\mathcal{H}_{n,\delta}) = V(\mathcal{F}_{n,\delta})$ and $E(\mathcal{H}_{n,\delta}) = E(\mathcal{F}_{n,\delta}) \cup \{u_0 w_i : 1 \leq i \leq n - \delta - 1\}$.

Hence, $\mathcal{H}_{n,\delta}$ is not 1-tough and therefore not hamiltonian, too.

Theorem 3 can be now restated as follows.

Theorem 3 (restated)**.**
If $G \in \mathcal{G}_1$, then G is hamiltonian. ■

Using closure operations we obtain the following extension of Theorem 3.

Theorem 4. *If $G \in \mathcal{G}_2$, then G is hamiltonian or $G \subset \mathcal{F}_{n,\delta}$.*

The proof of the above theorem is given in Section 3. It provides a further extension which can be formulated as follows.

Theorem 5. *If $G \in \mathcal{G}_3$, then G is hamiltonian or $G \subset \mathcal{H}_{n,\delta}$.*

Since both $\mathcal{F}_{n,\delta}$ and $\mathcal{H}_{n,\delta}$ are not 1-tough, we obtain the following corollary.

Corollary 6. *If $G \in \mathcal{G}_3$ is 1-tough, then G is hamiltonian.*

Bondy suggested the interesting "meta-conjecture" in [2] that almost any nontrivial condition on graphs which implies that the graph is hamiltonian also implies that the graph is pancyclic (there may be a family of exceptional graphs). He proved the following result concerning Ore's condition.

Theorem 7 ([2])**.** *Let G be a graph of order $n \geq 3$. If $d(u) + d(v) \geq n$ for every pair of nonadjacent vertices $u, v \in V(G)$, then G is pancyclic or isomorphic to the complete bipartite graph $K_{\frac{n}{2},\frac{n}{2}}$.*

In [7] it was shown that Theorem 7 can be extended as follows.

Theorem 8. *If $G \in \mathcal{G}_1$, then G is pancyclic or $G \cong K_{\frac{n}{2},\frac{n}{2}}$.* ■

Theorem 3 extends the following result of Jin, Liu and Wang [8].

Corollary 9 ([8])**.** *Let G be a 2-connected graph of order $n \geq 3$. If $d(u) + d(v) \geq n + \delta$ for every pair of nonadjacent vertices $u, v \in V(G)$, then G is pancyclic or $G \cong K_{\frac{n}{2},\frac{n}{2}}$.* ■

Concerning pancyclicity we will prove the following theorems.

Theorem 10. *If $G \in \mathcal{G}_2$, then G is pancyclic or $G \cong K_{\frac{n}{2},\frac{n}{2}}$ or $G \subset \mathcal{F}_{n,\delta}$.*

Theorem 11. *If $G \in \mathcal{G}_3$, then G is pancyclic or bipartite or $G \subset \mathcal{H}_{n,\delta}$.*

3. Proofs

3.1. Hamiltonicity

The closure concept of Bondy and Chvátal [3] is based on the following result of Ore [9].

Theorem 12 (Ore [9])**.** *Let G be a graph on n vertices such that the edge $e = uv$ does not belong to $E(G)$ and $d(u) + d(v) \geq n$. Then, the graph G is hamiltonian if and only if the graph $G + e$ is hamiltonian.*

By successively joining pairs of nonadjacent vertices having degree sum at least n as long as this is possible (in the new graph(s)), the unique so called n-closure $cl_n(G)$ is obtained. Using Theorem 12 it is easy to prove the following result.

Theorem 13 (Bondy and Chvátal [3]). *Let G be a graph of order $n \geq 3$. Then G is hamiltonian if and only if $cl_n(G)$ is hamiltonian.*

Corollary 14 (Bondy and Chvátal [3]). *Let G be a graph of order $n \geq 3$. If $cl_n(G)$ is complete ($cl_n(G) = K_n$), then G is hamiltonian.*

Ainouche and Christofides [1] established the following generalization of Theorem 12.

Theorem 15 (Ainouche and Christofides [1]). *Let G be a 2-connected graph on n vertices such that the edge $e = uv$ does not belong to $E(G)$. Let $T = T(u, v) = \{w \in V(G) \setminus (N[u] \cup N[v])\}$ and let $t = |T|$. Suppose that*

$$(*) \qquad d(w) \geq t + 2 \quad \text{for all vertices of } T.$$

Then, the graph G is hamiltonian if and only if the graph $G + e$ is hamiltonian.

In [1] the corresponding (unique) closure of G is called the 0-dual closure $cl^*(G)$. Since Theorem 15 is more general than Theorem 12 (cf. [1]), $G \subseteq cl_n(G) \subseteq cl^*(G)$. The counterpart of Corollary 14 is

Corollary 16 (Ainouche and Christofides [1]). *Let G be a 2-connected graph of order n. If $cl^*(G)$ is complete ($cl^*(G) = K_n$), then G is hamiltonian.*

***Proof of Theorem* 4.** Observe first that if $\delta(G) \geq \frac{n}{2}$ then G is hamiltonian by Dirac's theorem. So, assume that $B \neq \emptyset$.

Step 0. Applying the Bondy-Chvátal closure to the set A we get the graph G_0 with the set A complete.

Step 1. By using cl^* we are able to add to G_0 all edges connecting the vertex u with the set A. Indeed, it suffices to verify the hypothesis of Theorem 15. Suppose there exists a vertex $x \in A$ such that $ux \notin E$. Since u is adjacent to all vertices of B and x is adjacent (in G_0) to all vertices of A, we have $T_{G_0}(u, x) = \emptyset$. Denote the graph obtained in this step by G_1.

Step 2. Let $x \in B, x \neq u$. We put $a(x) = a_G(x) = |N_G(x) \cap A|$. Denote by B' the vertices of B with $a(x) < \delta - 1$. Consider now a vertex $x \in B'$ and

a vertex $y \in A$ such that $xy \notin E$. Then x has at least one neighbour in B different from u. This implies $|T(x,y)| \leq \delta - 2$. Hence, the condition $(*)$ of Theorem 15 is satisfied. This means that we can add all edges between B' and A. We denote the graph obtained in this step by G_2.

Step 3. Denote by B_1 the vertices of B different from u that are joined to all vertices of A in G_2. Note that $B' \subset B_1$. We put $B_2 = B - (B_1 \cup \{u\})$. Let $\xi = |B_1|$ and $\eta = |B_2|$. We have $1 + \xi + \eta = |B| \leq \delta + 1$. Consider now a vertex $x \in B_2$ and $y \in A$ such that $xy \notin E$. By Step 2, $a_{G_2}(x) \geq \delta - 1$. Since the vertices of A as well as the vertices of B_1 and the vertex u are the neighbours of y we get $|T(x,y)| \leq \eta - 1$. So, $t + 2 \leq \eta + 1$. If the condition $(*)$ of Theorem 15 is not fulfilled then $\delta \leq \eta$. This implies in particular that $\xi = 0$ and $\delta = \eta$. Moreover, x has no neighbour in B other than u, for otherwise $|T(x,y)| \leq \eta - 2$ and $(*)$ would be satisfied. Observe that either

(a) the above statements concern all vertices of B_2 (see Step 5), or

(b) we can add all edges between B_2 and A.

In the later case we can continue the closure operation (see Step 4 below).

Step 4. Denote by G_3 the graph obtained in Step 3b. Let x, y be two vertices of B such that $xy \notin E$. Then at most $\delta - 2$ vertices of B belong to $T(x,y)$ and we can finish the closure operation with the conclusion that $\mathrm{cl}^*(G) = K_n$.

Step 5. Suppose now that no edge can be added in Step 3. Then B consists of the vertex u and its δ neighbours, say $u_1, u_2, \ldots, u_\delta$, forming an independent set. This implies that each of the vertices $u_1, u_2, \ldots, u_\delta$ sends at least $\delta - 1$ edges to A. Suppose, that there exists a vertex $x \in A$ such that $u_i x \notin E$ and $u_j x \in E$ for some $j \neq i$.. Then $|T(u_i, x)| \leq \delta - 2$ and the edge $u_i x$ can be added to G_2. Denote by G_5 the graph obtained from G_2 by adding all edges as above. We may conclude that in G_5 all vertices $u_1, u_2, \ldots, u_\delta$ have the same neighbourhood. It is now easy to see that only in the case where this neighbourhood contains exactly $\delta - 1$ vertices of A the graph G_5 is not hamiltonian. Observe that in this case $G_5 \subset \mathcal{F}_{n,\delta}$. ■

***Proof of Theorem* 5.** As in the previous proof observe first that if $\delta(G) \geq \frac{n}{2}$ then G is hamiltonian by Dirac's theorem. So, assume that $B \neq \emptyset$. Applying the Bondy-Chvátal closure to the set A we get the graph G_0 with the set A complete. It is easy to verify the hamiltonicity of the

graph G_0 if $|B| \leq 2$. So, suppose $|B| \geq 3$. Then $\Delta(G[B]) \geq 2$. Let u be a vertex of B having the maximum number of neighbours in B. Consider a vertex $x \in A$ with $ux \notin E(G_0)$. Then $|T(u,x)| \leq \delta - 2$. This implies that the operation cl* can be applied. That means we can add all edges between u and A. Denote the graph obtained in this way by G_1.

Suppose now that there exists a vertex $x \in B$ such that $ux \notin E(G_0)$. It is easy to see that $|T(u,x)| \leq \delta - 3$. Therefore, we can add all edges between u and other vertices of B. It suffices now to observe that the graph obtained in this way has the same properties as the graph G_1 in the proof of the previous theorem. Now we can follow that proof. ■

In terms of the 0-dual closure Theorem 5 can be restated as follows.

Theorem 5 (restated).
If $G \in \mathcal{G}_3$, then G is hamiltonian or $cl^*(G) = (\overline{K}_\delta \cup K_{n-2\delta}) + K_\delta$.

3.2. Pancyclicity

For the proof of Theorem 11 we will apply the following three theorems.

Theorem 17 (Faudree, Häggkvist, Schelp [6]). *Every hamiltonian graph of order n and size $e(G) > \frac{(n-1)^2}{4} + 1$ is pancyclic or bipartite.* ■

Lemma 18 (Bondy [2]). *Let G be a hamiltonian graph of order n with a Hamilton cycle $v_1v_2 \ldots v_nv_1$ such that $d(v_1) + d(v_n) \geq n + 1$. Then G is pancyclic.*

Theorem 19 (Schmeichel-Hakimi [10]). *If G is a hamiltonian graph of order $n \geq 3$ with a Hamilton cycle $v_1v_2...v_nv_1$ such that $d(v_1) + d(v_n) \geq n$, then G is either*

- *pancyclic,*
- *bipartite, or*
- *missing only an $(n-1)$-cycle.*

Moreover, in the last case we have $d(v_{n-2}), d(v_{n-1}), d(v_2), d(v_3) < n/2$.

Remark. Actually, the Schmeichel-Hakimi result gives some more information about the possible adjacency structure near the vertices v_1 and v_n, but the above version is sufficient for our proof.

***Proof of Theorem* 11.** If $\delta \geq n/2$, then G is pancyclic or $G \cong K_{\frac{n}{2},\frac{n}{2}}$ by Theorem 7. Hence we may assume that $2 \leq \delta \leq \frac{n-1}{2}$. If $G \in \mathcal{H}_{n,\delta}$, then G is not hamiltonian and thus not pancyclic. Hence we may further assume that G is hamiltonian.

If $\delta = \frac{n-1}{2}$, then

$$e(G) \geq \frac{1}{2}\left(\frac{n+1}{2} \cdot \frac{n-1}{2} + \frac{n-1}{2} \cdot \frac{n+1}{2}\right) = \frac{n^2-1}{4} > \frac{(n-1)^2}{4} + 1$$

for all $n \geq 5$. Thus G is pancyclic or bipartite by Theorem 17 (since n is odd, G cannot be bipartite).

If $\delta = \frac{n-2}{2}$, then

$$e(G) \geq \frac{1}{2}\left(\frac{n}{2} \cdot \frac{n-2}{2} + \frac{n}{2} \cdot \frac{n}{2}\right) = \frac{n^2-n}{4} > \frac{(n-1)^2}{4} + 1$$

for all $n \geq 6$. Thus G is pancyclic or bipartite by Theorem 17. In the later case we conclude that $G \cong K_{\frac{n}{2},\frac{n}{2}} - \frac{n}{4}K_2$.

If $2 \leq \delta \leq \frac{n-3}{2}$, then $|A| \geq \frac{n+1}{2} > |B|$, since $|B| \leq \delta + 1 \leq \frac{n-1}{2}$. In this case the third alternative of Theorem 19 cannot occur since a simple counting argument gives $|A| \leq |B|$, a contradiction. Hence G is pancyclic or bipartite by Theorem 19. ■

***Proof of Theorem* 10.** Since $\mathcal{G}_2 \subset \mathcal{G}_3$ we can apply Theorem 11. If $G \subset \mathcal{H}_{n,\delta}$ then we conclude that $G \subset \mathcal{F}_{n,\delta}$, since there is a vertex $u \in B$ with $d(u) = \delta$. Suppose $G \not\subset \mathcal{F}_{n,\delta}$. Then G is hamiltonian by Theorem 4. Thus, if G is bipartite, then G is balanced bipartite with partite sets V_1 and V_2. Suppose $u \in V_1$ for a vertex u with $d(u) = \delta$ and $uv \in E(G)$ for all vertices $v \in B - \{u\}$. Since $\delta < n/2$, there exists a vertex $w \in V_2$ with $w \notin N(u)$. But then $d(w) \leq \frac{n}{2} - 1 < \frac{n}{2}$, a contradiction. Thus G cannot be bipartite. Therefore, by Theorem 11, G is pancyclic. ■

4. Concluding Remarks

Our results presented in Section 2 all imply that $|B| \leq \delta + 1$ for the considered graphs. Thus it is a natural question to study hamiltonicity (and pancyclicity) of graphs with $|B| \leq \delta + k$ for some positive integer $k \geq 2$.

For all n, δ and k with $2 \leq \delta \leq \frac{n-1}{2}$ and $1 \leq k \leq \delta - 1$ define $\mathcal{I}_{n,\delta,k}$ as a graph of order n, minimum degree δ and vertex set

$$V(\mathcal{I}_{n,\delta,k}) = \{u_1, \ldots, u_k, v_1, \ldots, v_\delta, w_1, \ldots, w_{n-\delta-k}\}$$

such that $d(u_i) = n - 1$ for $1 \leq i \leq k$, the vertices $\{v_1, \ldots, v_\delta\}$ are independent, $G[\{w_1, \ldots, w_{n-\delta-k}\}]$ is complete and $v_i w_j \in E(G)$ for all $1 \leq i \leq \delta$ and $1 \leq j \leq \delta - k$.

Now, for $S = \{u_1, \ldots, u_k, w_1, \ldots, w_{\delta-k}\}$ we have

$$\omega(\mathcal{I}_{n,\delta,k} - S) = \delta + 1 > \delta = |S|.$$

Hence, $\mathcal{I}_{n,\delta,k}$ is not 1-tough and thus not hamiltonian. Note that $\mathcal{I}_{n,\delta,1} = \mathcal{H}_{n,\delta}$.

Following the proof of Theorem 5 we have obtained the following theorem.

Theorem 20. *Let G be a 2-connected graph of order n. If for some k with $1 \leq k \leq \delta - 1$*

(i) *$G[B]$ is complete for $|B| \leq k + 1$ or*

(ii) *there are at least k vertices of degree at least $k + 1$ in B for $k + 2 \leq |B| \leq \delta + k$,*

then G is hamiltonian or $\subset \mathcal{I}_{n,\delta,k}$. ■

References

[1] A. Ainouche and N. Christofides, *Semi-independence number of a graph and the existence of hamiltonian circuits*, Discrete Appl. Math. **17** (1987) 213–221.

[2] J.A. Bondy, *Pancyclic graphs* I, J. Combin. Theory **11** (1971) 80–84.

[3] J.A. Bondy and V. Chvátal, *A method in graph theory*, Discrete Math. **15** (1976) 111–136.

[4] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications (Elsevier North Holland, New York, 1976).

[5] G.A. Dirac, *Some theorems on abstract graphs*, Proc. London Math. Soc. **2** (1952) 69–81.

[6] R.J. Faudree, R.Häggkvist and R.H. Schelp, *Pancyclic graphs — connected Ramsey number*, Ars Combin. **11** (1981) 37–49.

[7] E. Flandrin, H. Li, A. Marczyk and M. Woźniak, *A note on a new condition implying pancyclism*, Discuss. Math. Graph Theory **21** (2001) 137–143.

[8] G. Jin, Z. Liu and C. Wang, *Two sufficient conditions for pancyclic graphs*, Ars Combinatoria **35** (1993) 281–290.

[9] O. Ore, *Note on hamilton circuits*, Amer. Math. Monthly **67** (1960) 55.

[10] E.F. Schmeichel and S.L. Hakimi, *A cycle structure theorem for hamiltonian graphs*, J. Combin. Theory (B) **45** (1988) 99–107.

Received 1 June 2005
Revised 28 April 2006