Discussiones Mathematicae Graph Theory 27 (2007) 83–91

TREES WITH EQUAL RESTRAINED DOMINATION AND TOTAL RESTRAINED DOMINATION NUMBERS

JOANNA RACZEK

Department of Discrete Mathematics Faculty of Applied Physics and Mathematics Gdańsk University of Technology Narutowicza 11/12, 80–952 Gdańsk, Poland

 $\mathbf{e}\text{-}\mathbf{mail:}$ gardenia@pg.gda.pl

Abstract

For a graph G = (V, E), a set $D \subseteq V(G)$ is a total restrained dominating set if it is a dominating set and both $\langle D \rangle$ and $\langle V(G) - D \rangle$ do not have isolated vertices. The cardinality of a minimum total restrained dominating set in G is the total restrained domination number. A set $D \subseteq V(G)$ is a restrained dominating set if it is a dominating set and $\langle V(G) - D \rangle$ does not contain an isolated vertex. The cardinality of a minimum restrained dominating set in G is the restrained domination number. We characterize all trees for which total restrained and restrained domination numbers are equal.

Keywords: total restrained domination number, restrained domination number, trees.

2000 Mathematics Subject Classification: 05C05, 05C69.

1. Introduction

Let G = (V(G), E(G)) be a simple graph with |V(G)| = n(G). The neighbourhood $N_G(u)$ of a vertex u is the set of all vertices adjacent to u in G and the closed neighbourhood of u is $N_G[u] = N_G(u) \cup \{u\}$. For a set $D \subseteq V(G)$ the closed neighbourhood of D is defined to be $\bigcup_{u \in D} N_G[u]$. The private neighbourhood of a vertex u with respect to a set $D \subseteq V(G)$, where $u \in D$, is the set $PN_G[u, D] = N_G[u] - N_G[D - \{u\}]$. If $v \in PN_G[u, D]$,

then we say that v is a private neighbour of u with respect to the set D. The degree $d_G(u)$ of a vertex u is the number of edges incident to u in G, that is $d_G(u) = |N_G(u)|$. Let $\Omega(G)$ be the set of all leaves of G, that is the set of vertices degree 1. A vertex which is a neighbour of a leaf is called a support vertex. Let S(G) be the set of all support vertices in G. The diameter diam(G) of a connected graph G is the maximum distance between two vertices of G, that is diam $(G) = \max_{u,v \in V(G)} d_G(u,v)$. We say that a set $D \subseteq V(G)$ is independent, if the induced subgraph $\langle D \rangle$ has no edge.

A set $D \subseteq V(G)$ is a dominating set of G if for every vertex $v \in V(G) - D$ there exists a vertex $u \in D$ such that v and u are adjacent. The minimum cardinality of a dominating set in G is the domination number denoted $\gamma(G)$. A minimum dominating set of a graph G is called a $\gamma(G)$ -set.

A set $D \subseteq V(G)$ is a restrained dominating set of G (RDS) if D is a dominating set and the induced subgraph $\langle V(G) - D \rangle$ does not contain an isolated vertex. The cardinality of a minimum restrained dominating set in G is the restrained domination number and is denoted by $\gamma_r(G)$. A minimum RDS of a graph G is called a $\gamma_r(G)$ -set. The concept of restrained domination was introduced by Telle and Proskurowski [6], albeit indirectly, as a vertex partitioning problem. Restrained domination was studied further for example by Domke *et al.* [1, 2].

The total restrained domination number of a graph was defined by Ma, Chen and Sun [5]. A set $D \subseteq V(G)$ is a *total restrained dominating set of* G(TRDS) if it is a dominating set and the induced subgraphs $\langle D \rangle$ and $\langle V(G) - D \rangle$ do not contain isolated vertices. The cardinality of a minimum total restrained dominating set in G is the *total restrained domination number* and is denoted by $\gamma_r^t(G)$. A minimum TRDS of a graph G is called a $\gamma_r^t(G)$ set. We note that every graph G without an isolated vertex has a (total) restrained dominating set, since D = V(G) is such a set.

For any graph theoretical parameters λ and μ , we define G to be (λ, μ) graph if $\lambda(G) = \mu(G)$. Henning has written an extensive series of papers which give constructive characterizations of trees for which certain domination parameters are equal (see, for example [4]). In this paper we provide a constructive characterization of (γ_r, γ_r^t) -trees. For any unexplained terms and symbols see [3].

2. A Characterization of (γ_r, γ_r^t) -Trees

As a consequence of the definitions of the restrained and total restrained

domination numbers we have the following observations.

Observation 1. Let G be a graph without isolated vertices. Then

- (i) every leaf is in every $\gamma_r^t(G)$ -set;
- (ii) every support vertex is in every $\gamma_r^t(G)$ -set;
- (iii) every leaf is in every $\gamma_r(G)$ -set;
- (iv) $\gamma(G) \leq \gamma_r(G) \leq \gamma_r^t(G)$.

Observation 2. Let T be a (γ_r, γ_r^t) -tree. Then each $\gamma_r^t(T)$ -set is a $\gamma_r(T)$ -set.

Let \mathcal{T}_1 and \mathcal{T}_2 be the following two operations defined on a tree T.

- **Operation** \mathcal{T}_1 . Assume $x \in V(T)$ is a support vertex. Then add a vertex y and the edge xy.
- Operation \mathcal{T}_2 . Assume $x \in V(T)$ is a support vertex. Then add a path $P_4 = (y_1, y_2, y_3, y_4)$ and the edge xy_1 .

Let \mathcal{T} be the family of trees such that $\mathcal{T} = \{T : T \text{ is obtained from } P_3 \text{ by a finite sequence of Operations } \mathcal{T}_1 \text{ or } \mathcal{T}_2\} \cup \{P_2, P_6\}$. We show first that each tree in the family \mathcal{T} has equal restrained domination number and total restrained domination number.

Lemma 3. If T belongs to the family \mathcal{T} , then T is a (γ_r, γ_r^t) -tree.

Proof. We proceed by induction on the number s(T) of operations required to construct the tree T. If s(T) = 0, then $T \in \{P_2, P_3, P_6\}$ and clearly T is a (γ_r, γ_r^t) -tree. Assume now that T is a tree with s(T) = k for some positive integer k and each tree $T' \in \mathcal{T}$ with s(T') < k is a (γ_r, γ_r^t) -tree. Then Tcan be obtained from a tree T' belonging to \mathcal{T} by operation \mathcal{T}_1 or \mathcal{T}_2 . We now consider two possibilities depending on whether T is obtained from T'by Operation \mathcal{T}_1 or \mathcal{T}_2 .

Case 1. T is obtained from T' by Operation \mathcal{T}_1 . Suppose T is obtained from T' by adding a vertex y and the edge xy, where $x \in V(T')$ is a support vertex. Thus y belongs to every $\gamma_r(T)$ -set and every $\gamma_r^t(T)$ -set. Hence $\gamma_r(T) = \gamma_r(T') + 1$ and $\gamma_r^t(T) = \gamma_r^t(T') + 1$. Since $\gamma_r(T') = \gamma_r^t(T')$ and $\gamma_r(T) \leq \gamma_r^t(T)$, we conclude that $\gamma_r(T) = \gamma_r^t(T)$.

Case 2. T is obtained from T' by Operation \mathcal{T}_2 . Suppose T is obtained from T' by adding a path (y_1, y_2, y_3, y_4) and the edge xy_1 , where $x \in V(T')$

is a support vertex. Then x and y_3 are support vertices in T and y_4 is a leaf. Hence x, y_3 and y_4 belong to every $\gamma_r^t(T)$ -set and for this reason $\gamma_r^t(T) \ge \gamma_r^t(T') + 2$. On the other hand, any $\gamma_r^t(T')$ -set may be extended to a TRDS of T by adding to it y_3 and y_4 . Thus $\gamma_r^t(T) = \gamma_r^t(T') + 2$.

Now let D be a $\gamma_r(T)$ -set. Then $y_4 \in D$ and $N_T[y_2] \cap D \neq \emptyset$. For this reason $\gamma_r(T) \geq \gamma_r(T') + 2$. On the other hand, $\gamma_r(T) \leq \gamma_r^t(T) = \gamma_r^t(T') + 2 = \gamma_r(T') + 2$. We conclude that $\gamma_r(T) = \gamma_r(T') + 2$ and consequently, $\gamma_r(T) = \gamma_r^t(T)$.

We now show that every (γ_r, γ_r^t) -tree belongs to the family \mathcal{T} . It is clear that P_2 is a (γ_r, γ_r^t) -tree and P_2 belongs to the family \mathcal{T} . Therefore from now on we consider only trees T with $n(T) \geq 3$.

Lemma 4. Let T be a (γ_r, γ_r^t) -tree with $n(T) \geq 3$ and let D_r^t be a minimum total restrained dominating set of T. If $u, v \in D_r^t$ and $uv \in E(T)$, then either u or v is a leaf.

Proof. It is possible to see that the statement is true for all trees T with diameter 2 and 3. For this reason we consider only trees with diameter at least 4. Suppose T is a (γ_r, γ_r^t) -tree, $u, v \in D_r^t$, $uv \in E(T)$ and neither u nor v is a leaf. We consider three cases.

Case 1. u is an isolated vertex in $\langle (V(T) - D_r^t) \cup \{u\} \rangle$ and v is an isolated vertex in $\langle (V(T) - D_r^t) \cup \{v\} \rangle$. Since neither u nor v is a leaf, we conclude that $D_r^t - \{u, v\}$ is a RDS of T of cardinality smaller than $\gamma_r(T)$, a contradiction.



Figure 1. Illustration for *Case* 2 of the proof of Lemma 4.

Case 2. Both $\langle (V(T) - D_r^t) \cup \{u\} \rangle$ and $\langle (V(T) - D_r^t) \cup \{v\} \rangle$ are without isolated vertices. Then since T is a (γ_r, γ_r^t) -tree, we conclude that $D_r^t - \{u\}$ and $D_r^t - \{v\}$ are not dominating sets of T. Therefore, both u and v have a private neighbour with respect to D_r^t . Let $U_0 = \{u\}$ and $V_0 = \{v\}$ and

denote by U_1 and V_1 the sets of private neighbours of u and v with respect to D_r^t , respectively. Of course, $U_1 \cap V_1 = \emptyset$ and $U_1 \cup V_1$ is an independent set of vertices, because T is a tree. Since D_r^t is a TRDS, each vertex of $U_1 \cup V_1$ has a neighbour in $V(T) - D_r^t$. Denote by U_2 and V_2 the sets of all vertices of $V(T) - D_r^t$ which are neighbours of vertices of U_1 and V_1 , respectively. Observe that $U_2 \cap V_2 = \emptyset$, $U_1 \cap U_2 = \emptyset$, $V_1 \cap V_2 = \emptyset$ and $U_2 \cup V_2$ is an independent set of vertices. Since T is a tree, no two vertices of $U_1 \cup V_1$ have common neighbour in $U_2 \cup V_2$, so $|U_1| \leq |U_2|$ and $|V_1| \leq |V_2|$. Moreover, since D_r^t is a dominating set of T, each vertex of $U_2 \cup V_2$ has a neighbour in D_r^t . Denote by U_3 and V_3 the sets of all vertices belonging to D_r^t which are neighbours of vertices of U_2 and V_2 , respectively. Since T is a tree, $(U_3 \cup V_3) \cap \{u, v\} = \emptyset, U_3 \cap V_3 = \emptyset, U_3 \cup V_3$ is an independent set of vertices, $|U_2| \leq |U_3|$ and $|V_2| \leq |V_3|$. Finally, since D_r^t is a TRDS of T, each vertex of $U_3 \cup V_3$ has a neighbour in D_r^t . Denote by U_4 and V_4 the sets of all vertices belonging to D_r^t which are neighbours of vertices of U_3 and V_3 , respectively. Since T is a tree, $(U_4 \cup V_4) \cap \{u, v\} = \emptyset$, $(U_4 \cup V_4) \cap (U_3 \cup V_3) = \emptyset$, $U_4 \cap V_4 = \emptyset$, $U_4 \cup V_4$ is an independent set of vertices, $|U_3| \leq |U_4|$ and $|V_3| \leq |V_4|$. Define U_5 to be the set of vertices of $V(T) - U_2$ which are private neighbours with respect to D_r^t of vertices belonging to U_3 and define V_5 to be the set of vertices of $V(T) - V_2$ which are private neighbours with respect to D_r^t of vertices belonging to V_3 . Denote by U_6 and V_6 the sets of all vertices of $V(T) - D_r^t$ which are neighbours of vertices of U_5 and V_5 , respectively, and so on.

Generally, let k be a non-negative integer. Define U_{4k+5} to be the set of vertices of $V(T) - U_{4k+2}$ which are private neighbours with respect to D_r^t of vertices belonging to U_{4k+3} and define V_{4k+5} to be the set of vertices of $V(T) - V_{4k+2}$ which are private neighbours with respect to D_r^t of vertices belonging to V_{4k+3} . Since D_r^t is a TRDS, each vertex of $U_{4k+1} \cup V_{4k+1}$, where $k \ge 0$, has a neighbour in $V(T) - D_r^t$. Let U_{4k+2} be the set of all vertices of $V(T) - D_r^t$ which are neighbours of vertices of U_{4k+1} and let V_{4k+2} be the set of all vertices of all vertices of $V(T) - D_r^t$ which are neighbours of vertices of $U_{4k+2} \cup V_{4k+2}$ be the set of all vertices of all vertices of V_{4k+3} the set of all vertices belonging to D_r^t which are neighbours of $V_{4k+2} \cup V_{4k+2}$ has a neighbour in D_r^t . Denote by U_{4k+3} the set of all vertices belonging to D_r^t which are neighbours of vertices of V_{4k+2} and denote by V_{4k+3} the set of all vertices belonging to D_r^t which are neighbours of vertices of U_{4k+2} . Finally, since D_r^t is a TRDS of T, each vertex of $U_{4k+3} \cup V_{4k+3}$ has a neighbour in D_r^t . Denote by U_{4k+4} and V_{4k+4} the sets of all vertices belonging to D_r^t which are neighbours of vertices of V_{4k+3} has a neighbour in D_r^t . Denote by U_{4k+4} and V_{4k+4} the sets of all vertices belonging to D_r^t which are neighbours of vertices of U_{4k+4} the sets of all vertices belonging to D_r^t which are neighbours of vertices belonging to D_r^t . Denote by U_{4k+4} and V_{4k+4} the sets of all vertices belonging to D_r^t which are neighbours of vertices of $U_{4k+3} \cup V_{4k+3}$ has a neighbour in D_r^t . Denote by U_{4k+4} and V_{4k+4} the sets of all vertices belonging to D_r^t which are neighbours of vertices of U_{4k+3} and V_{4k+3} , respectively. Since T is a finite tree, there exist

the smallest integer *i* such that $U_{4i+5} = \emptyset$ and the smallest integer *j* such that $V_{4i+5} = \emptyset$.

Since T is a tree, we conclude that no two vertices of $U_{4k+1} \cup V_{4k+1}$ have common neighbour in $U_{4k+2} \cup V_{4k+2}$. This implies that $|U_{4k+1}| \leq |U_{4k+2}|$ and $|V_{4k+1}| \leq |V_{4k+2}|$. Similarly, $|U_{4k+2}| \leq |U_{4k+3}|$ and $|V_{4k+2}| \leq |V_{4k+3}|$. Further, $|U_{4k+3}| \leq |U_{4k+4}|$ and $|V_{4k+3}| \leq |V_{4k+4}|$. Moreover, every two of defined sets are disjoint.

Now consider the set $D = D_r^t - (U_3 \cup U_7 \cup \cdots \cup U_{4i+3} \cup V_3 \cup V_7 \cup \cdots \cup V_{4j+3} \cup \{u, v\}) \cup U_1 \cup U_5 \cup \cdots \cup U_{4i+1} \cup V_1 \cup V_5 \cup \cdots \cup V_{4j+1}$. It is possible to observe that D is a dominating set of T and $\langle V(T) - D \rangle$ does not contain an isolated vertex. Hence D is a RDS of T. Moreover $|D| < |D_r^t|$, which implies that T is not a (γ_r, γ_r^t) -tree, a contradiction.

Case 3. Either $\langle (V(T) - D_r^t) \cup \{u\} \rangle$ or $\langle (V(T) - D_r^t) \cup \{v\} \rangle$ contains an isolated vertex, say u is an isolated vertex in $\langle (V(T) - D_r^t) \cup \{u\} \rangle$. Then since T is a (γ_r, γ_r^t) -tree, we conclude that $D_r^t - \{v\}$ is not a dominating set of T. Let j and $V_0, V_1, \ldots, V_{4j+5}$ have the same meaning and properties as in previous case. Consider the set $D = D_r^t - (V_3 \cup V_7 \cup \cdots \cup V_{4j+3} \cup \{u, v\}) \cup V_1 \cup V_5 \cup \cdots \cup V_{4j+1}$. It is easy to observe that D is a dominating set of T and $\langle V(T) - D \rangle$ does not contain an isolated vertex. Hence D is a RDS of T. Moreover $|D| < |D_r^t|$, which implies that T is not a (γ_r, γ_r^t) -tree, a contradiction.

This proves the statement.

The above Lemma together with Lemma 1 imply what follows.

Corollary 5. If T is a (γ_r, γ_r^t) -tree with $n(T) \ge 3$, then $\Omega(T) \cup S(T)$ is the unique $\gamma_r^t(T)$ -set and $\gamma_r(T) = \gamma_r^t(T) = |\Omega(T) \cup S(T)|$.

Corollary 6. If T is a (γ_r, γ_r^t) -tree with $n(T) \ge 3$, then S(T) is a $\gamma(T)$ -set and $\gamma(T) = |S(T)|$.

Corollary 7. If T is a (γ_r, γ_r^t) -tree with $n(T) \ge 3$, then $\gamma_r^t(T) = \gamma(T) + |\Omega(T)|$.

Lemma 8. Let T be a (γ_r, γ_r^t) -tree with $n(T) \ge 3$. If $u, v \in S(T)$, then $d_T(u, v) \ge 3$.

Proof. It is possible to verify that the statement is true for all trees with diameter between 2 and 5. For this reason we consider only trees with diameter at least 6.

Let T be a (γ_r, γ_r^t) -tree with $n(T) \geq 3$ and let D_r^t be a $\gamma_r^t(T)$ -set. By Corollary 5, $u, v \in D_r^t$ and by Lemma 4, u and v are not adjacent. Suppose that $d_T(u, v) = 2$ and let x be the neighbour of u and v in T. Lemma 4 implies that x is not a support vertex and as x is not a leaf, $x \notin D_r^t$. Since both $\langle (V(T) - D_r^t) \cup \{u\} \rangle$ and $\langle (V(T) - D_r^t) \cup \{v\} \rangle$ are without isolated vertices and T is a (γ_r, γ_r^t) -tree, we deduce that $D_r^t - \{u\}$ and $D_r^t - \{v\}$ are not dominating sets of T. Therefore, both u and v have a private neighbour with respect to D_r^t . Let j and $V_0, V_1, \ldots, V_{4j+5}$ have the same meaning and properties as in the proof of Lemma 4. Consider the set $D = D_r^t - (V_3 \cup V_7 \cup \cdots \cup V_{4j+3} \cup \{v\}) \cup V_1 \cup V_5 \cup \cdots \cup V_{4j+1}$. It is possible to observe that D is a dominating set of T and $\langle V(T) - D \rangle$ does not contain an isolated vertex. Hence D is a RDS of T. Moreover $|D| < |D_r^t|$, which implies that T is not a (γ_r, γ_r^t) -tree, a contradiction.

Corollary 9. If T is a (γ_r, γ_r^t) -tree with $n(T) \ge 3$, then each vertex of V(T) - S(T) has exactly one neighbour in S(T).

Corollary 10. If T is a (γ_r, γ_r^t) -tree with $n(T) \ge 3$, then S(T) is the unique $\gamma(T)$ -set.

Lemma 11. If T is a (γ_r, γ_r^t) -tree with $n(T) \geq 3$, then T belongs to the family \mathcal{T} .

Proof. It is easily seen that the statement is true for all trees with with diameter between 2 and 5. For this reason we consider only trees with diameter at least 6.

Let T be a (γ_r, γ_r^t) -tree and assume that the result holds for all trees on n(T) - 1 and fewer vertices. We proceed by induction on the number of vertices of a (γ_r, γ_r^t) -tree. Let $P = (s_0, s_1, \ldots s_l), l \ge 6$, be a longest path in T and let D_r^t be a $\gamma_r^t(T)$ -set. We consider two cases.

Case 1. $d_T(s_1) > 2$. In this case s_1 is a neighbour of at least two leaves of T. Denote $T' = T - s_0$. Of course $D_r^t - \{s_0\}$ is a TRDS of T', so $\gamma_r^t(T') \leq \gamma_r^t(T) - 1$. Moreover, any $\gamma_r^t(T')$ -set may be extended to a $\gamma_r^t(T)$ -set by adding to it s_0 , so $\gamma_r^t(T') = \gamma_r^t(T) - 1$. By similar arguments it may be concluded that $\gamma_r(T') = \gamma_r(T) - 1$. Hence, $\gamma_r(T') = \gamma_r^t(T')$. Consequently, T' is a (γ_r, γ_r^t) -tree and by induction hypothesis, $T' \in \mathcal{T}$. As s_1 is a support vertex in T', we deduce that T may be obtained from T' by Operation \mathcal{T}_1 . Case 2. $d_T(s_1) = 2$. Then Corollary 5 and Lemma 8 imply that $d_T(s_2) = 2$ and s_3 is not a support vertex. Moreover, s_3 is a neighbour of exactly one support vertex, say x.

Suppose that $x \neq s_4$. Then s_4 is not a support vertex, but s_4 is a neighbour of exactly one support vertex, say y. Denote $A = N_T(s_3) - \{x\} - V(P)$ and observe that since x is a support vertex, Lemma 8 implies that $A \cap S(T) = \emptyset$. Corollary 9 says that each vertex of A has exactly one neighbour in S(T). Let A' be the set of neighbours of vertices of A which belong to S(T). Hence $s_0, s_1, x, y \in D_r^t$ and $s_2, s_3, s_4 \notin D_r^t$. Consider the set $D = D_r^t - \{s_1, y\} - A' \cup \{s_3\}$. It is easy to observe that D is a dominating set in T and $\langle V(T) - D \rangle$ does not contain an isolated vertex. Hence D is a RDS of T. Moreover $|D| < |D_r^t|$ even when $A = \emptyset$, which implies that T is not a (γ_r, γ_r^t) -tree, a contradiction. Therefore s_4 is the unique support vertex in $N_T(s_3)$.

Now suppose that $d_T(s_3) > 2$. Denote $A = N_T(s_3) - V(P)$ and observe that since $d_T(s_3) > 2$, $A \neq \emptyset$. Moreover, since s_4 is a support vertex, $A \cap S(T) = \emptyset$. Let A' be the set of neighbours of vertices of A which belong to S(T). Then $s_0, s_1, s_4 \in D_r^t$ and $s_2, s_3 \notin D_r^t$. Consider the set $D = ((D_r^t - \{s_1\}) - A') \cup \{s_3\}$. It is easy to observe that D is a dominating set of T and $\langle V(T) - D \rangle$ does not contain an isolated vertex. Hence D is a RDS of T. Moreover $|D| < |D_r^t|$, which implies that T is not a (γ_r, γ_r^t) -tree, a contradiction. Therefore $d_T(s_3) = 2$ and s_4 is the unique neighbour of s_3 belonging to S(T).

Denote $T' = T - \{s_0, s_1, s_2, s_3\}$. Of course s_0 and s_1 belong to every $\gamma_r^t(T)$ -set. For this reason, $\gamma_r^t(T') \leq \gamma_r^t(T) - 2$. Since s_4 is a support vertex in T', any $\gamma_r^t(T')$ -set may be extended to a TRDS of T by adding to it s_0 and s_1 , so $\gamma_r^t(T') = \gamma_r^t(T) - 2$. Further, $\gamma_r(T') \leq \gamma_r^t(T') = \gamma_r^t(T) - 2 = \gamma_r(T) - 2$ and any $\gamma_r(T')$ -set may be extended to a RDS of T by adding to it s_0 and s_3 . Hence $\gamma_r(T') = \gamma_r(T) - 2$ and so $\gamma_r(T') = \gamma_r^t(T')$. Consequently, T' is a (γ_r, γ_r^t) -tree and by induction hypothesis, $T' \in \mathcal{T}$. As s_4 is a support vertex in T', we conclude that T may be obtained from T' by Operation \mathcal{T}_2 .

As an immediate consequence of Lemmas 4 and 11 we have the following characterization of (γ_r, γ_r^t) -trees.

Theorem 12. A tree T is a (γ_r, γ_r^t) -tree if and only if T belongs to the family \mathcal{T} .

References

- G.S. Domke, J.H. Hattingh, S.T. Hedetniemi, R.C. Laskar and L.R. Marcus, Restrained domination in graphs, Discrete Math. 203 (1999) 61–69.
- [2] G.S. Domke, J.H. Hattingh, S.T. Hedetniemi and L.R. Marcus, *Restrained domination in trees*, Discrete Math. **211** (2000) 1–9.
- [3] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, Fundamentals of domination in graphs (Marcel Dekker, New York, 1998).
- [4] M.A. Henning, Trees with equal average domination and independent domination numbers, Ars Combin. 71 (2004) 305–318.
- [5] D. Ma, X. Chen and L. Sun, On total restrained domination in graphs, Czechoslovak Math. J. 55 (2005) 165–173.
- [6] J.A. Telle and A. Proskurowski, Algorithms for vertex partitioning problems on partial k-trees, SIAM J. Discrete Math. 10 (1997) 529–550.

Received 7 November 2005 Revised 21 August 2006