THE WIENER NUMBER OF KNESER GRAPHS

RANGASWAMI BALAKRISHNAN AND S. FRANCIS RAJ

Srinivasa Ramanujan Centre, SASTRA University Kumbakonam-612 001, India

> e-mail: mathbala@satyam.net.in e-mail: francisraj_s@yahoo.com

Abstract

The Wiener number of a graph G is defined as $\frac{1}{2} \sum d(u, v)$, where $u, v \in V(G)$, and d is the distance function on G. The Wiener number has important applications in chemistry. We determine the Wiener number of an important family of graphs, namely, the Kneser graphs.

Keywords: Wiener number, Kneser graph, odd graph.

2000 Mathematics Subject Classification: 05C12.

1. INTRODUCTION

Let G = (V, E) be a simple connected undirected graph with |V(G)| = nand |E(G)| = m. Given two distinct vertices u, v of G, let d(u, v) denote the distance (= number of edges in a shortest path between u and v in G). The Wiener number (also called Wiener Index) W(G) of the graph G is defined by

$$W = \frac{1}{2} \sum_{u,v \in V(G)} d(u,v).$$

Given the structure of an organic compound, the corresponding (molecular) graph is obtained by replacing the atoms by vertices and covalent bonds by edges. The Wiener number is one of the oldest molecular-graphbased structure-descriptors, first proposed by the American chemist Harold Wiener [18], as an aid to determine the boiling point of paraffins. The study of Wiener number is one of the current areas of research in mathematical chemistry (see, for example, [14] and [19]). For more details on the computation of Wiener number and its applications to chemistry, see [8]. Some recent articles in the topic are [2,5,6]. One of the important families of graphs is the family of Kneser graphs. There are a good number of papers in problems dealing with the coloring parameters of Kneser graphs. See for instance, [9–12, 16]. Extremal problems concerning Kneser graphs are considered in [7]. In this paper, we obtain an explicit expression, based on mathematical induction, for the Wiener number of the Kneser graphs. Our notation and terminology are as in [3].

We recall the definition of a Kneser graph.

Let n and k be positive integers, m = 2n + k, where $k \ge 1$. We denote by [m] the set $\{1, 2, \ldots, m\}$ and by $\binom{[m]}{n}$ the collection of all n-subsets of [m]. The Kneser graph KG(m, n) has vertex set $\binom{[m]}{n}$, in which two vertices are adjacent iff they are disjoint. It is to be noted that Kneser graphs are vertextransitive but not distance-regular and therefore not distance-transitive.

2. Some Basic Results

Lemma 2.1 (Stahl [15]). If A, B are two distinct vertices of KG(m, n), with d(A, B) = 2p, then $|A \cap B| \ge n - kp$.

A consequence of Lemma 2.1 is Lemma 2.2.

Lemma 2.2. If A, B are two distinct vertices of KG(m, n), with d(A, B) = 2p + 1, then $|A \cap B| \le kp$.

Proof. Let A, B, C be vertices of KG(m, n) such that d(A, C) = 2p, d(C, B) = 1 and d(A, B) = 2p + 1 so that C is the vertex preceding B in a A-B distance path in G. By Lemma 2.1, $|A \cap C| \ge n - kp$. But since $|B \cap C| = \emptyset$, $|A \cap B| \le n - (n - kp) = kp$.

Using Lemmas 2.1 and 2.2, Valencia-Pabon and Vera [13] have determined the diameter of G.

Lemma 2.3 ([13]). The diameter of the Kneser graph KG(m,n) is $\lceil \frac{n-1}{k} \rceil + 1$.

3. WIENER NUMBER OF KG(m, n)

We now compute the Wiener number of the Kneser graph KG(m, n).

We first observe that for any two distinct vertices of KG(m, n), $|A \cap B| \in \{0, 1, \ldots, n-1\}$. Further $d(A_0, A) = 0$ if and only if $|A_0 \cap A| = n$, and $d(A_0, A) = 1$ if and only if $|A_0 \cap A| = 0$.

Lemma 3.1. Let G = KG(m, n) be a Kneser graph with diameter D. Fix $A_0 \in V(G)$. Then for any $A \in V(G)$, $p \ge 1$,

- (i) $d(A_0, A) = 2p < D$, if and only if $|A_0 \cap A| \in N_{2p} := \{n - kp, n - kp + 1, \dots, n - k(p - 1) - 1\},\$
- (ii) $d(A_0, A) = 2p + 1 < D$, if and only if $|A_0 \cap A| \in N_{2p+1} := \{k(p-1) + 1, k(p-1) + 2, \dots, kp\}, and$
- (iii) $d(A_0, A) = D$, if and only if

$$|A_0 \cap A| \in N_D := \begin{cases} \left\{ k(\frac{D}{2} - 1) + 1, \dots, n - k(\frac{D}{2} - 1) - 1 \right\} & \text{if } D \text{ is even,} \\ \left\{ k(\frac{D-1}{2} - 1) + 1, \dots, n - k(\frac{D-1}{2}) - 1 \right\} & \text{if } D \text{ is odd.} \end{cases}$$

We set $N_0 = \{n\}$ and $N_1 = \{0\}$. Before we prove Lemma 3.1, we observe the following:

Observation 3.2. (a) $|N_0| = 1$, $|N_1| = 1$ and $|N_i| = k$ when 2 < i < D.

- (b) The sets N_{2p+1} are successive disjoint intervals (intervals of positive integers) that are increasing from 0 to n as p increases from 1, and the sets N_{2p} are successive disjoint intervals that are decreasing from n to 0 again as p increases from 1, that is, for i < j and for x ∈ N_i and y ∈ N_j, x < y if i and j are odd and x > y if i and j are even.
- (c) If $p \neq q$ and p < D and q < D, then $N_p \cap N_q = \phi$.

We prove (c); (a) and (b) are obvious from the definition of the sets N_i .

Proof of (c). If p and q are both even or both odd, there is nothing to prove. So let one of them be odd and the other even, say, p odd and q even. Also let i = largest even integer less than D, and j = largest odd integer less than D, and j = 2e + 1. This implies that $i = j \pm 1$. First, let i = j - 1, so that i = 2e. Again, from the definition of i and j, we observe

that $p \leq j = 2e + 1$, $q \leq i = 2e$ and therefore by (b) of Observation 3.2, it suffices to prove that $N_{2e} \cap N_{2e+1} = \emptyset$. Indeed, we verify that ke < n - ke - 1. This is true, since $ke \geq n - ke - 1$ implies that $n - 1 \leq 2ke \leq k(D - 2)$, a contradiction to the fact that $D = \lceil \frac{n-1}{k} \rceil + 1$. Similarly, we verify the result when i = j + 1.

Proof of Lemma 3.1. We prove (i) and (ii) by induction on i where $d(A_0, A) = i < D$.

Let $d(A_0, A) = 2$. By Lemma 2.1, $|A_0 \cap A| \ge n - k$. If $|A_0 \cap A| = n$, then $d(A_0, A) = 0$, a contradiction. Hence $n > |A_0 \cap A| \ge n - k$ and so $|A_0 \cap A| \in N_2$.

Conversely, let $|A_0 \cap A| \in N_2 = \{n - k, n - k + 1, \dots, n - 1\}$, so that $|A_0 \cap A| = n - k + r$, where $0 \leq r < k$. Consequently, $|A_0 \cup A| = 2n - (n - k + r) = n + k - r$; and hence $|[m] \setminus (A_0 \cup A)| = (2n + k) - (n + k - r) = n + r$ where $0 \leq r < k$. From these n + r elements, we can find a vertex Z in V(G) such that $|A_0 \cap Z| = 0 = |A \cap Z|$. Therefore, $d(A_0, Z) = d(A, Z) = 1$ and so $d(A_0, A) \leq 2$. Clearly, $d(A_0, A) < 2$ is not possible and therefore $d(A_0, A) = 2$.

Next, let $d(A_0, A) = 3$. By Lemma 2.2, $|A_0 \cap A| \le k$. If $|A_0 \cap A| = 0$, then $d(A_0, A) = 1$ which is not true. Therefore $|A_0 \cap A| \in N_3 = \{1, 2, ..., k\}$.

Conversely, let $|A_0 \cap A| \in N_3 = \{1, 2, \dots, k\}$, so that $|A_0 \cap A| = k - r$ for $0 \leq r < k$. Hence if $X = A_0 \cap A$, then $A_0 = X \cup \{\alpha_1, \alpha_2, \dots, \alpha_{n-k+r}\}$, and $A = X \cup \{\beta_1, \beta_2, \dots, \beta_{n-k+r}\}$, where $\{\alpha_1, \alpha_2, \dots, \alpha_{n-k+r}\}$ is disjoint from $\{\beta_1, \beta_2, \dots, \beta_{n-k+r}\}$. Further let $C = [m] \setminus (A_0 \cup A)$. Then |C| =(2n + k) - (2n - k + r) = 2k - r, where $0 \leq r < k$. Choose k elements c_1, \dots, c_k from C and set $B = (C \setminus \{c_1, \dots, c_k\}) \cup \{\alpha_1, \dots, \alpha_{n-k+r}\}$. Then |B| = (2k - r) - k + (n - k + r) = n and so $B \in V(G)$, $|A_0 \cap B| =$ n - k + r and $|A \cap B| = 0$. Since $0 \leq r < k$, we have $|A_0 \cap B| \in N_2$ and therefore, $d(A_0, B) = 2$. Also d(A, B) = 1. Hence $d(A_0, A) \leq 3$, which gives $d(A_0, A) = 3$. Thus we have established the result when $d(A_0, A) = 2$ or 3.

Assume now that $d(A_0, A) = i > 3$ and that the result is true if $d(A_0, A) \in \{2, 3, ..., i - 1\}, i < D$.

Case (i). Let *i* be odd so that i = 2p + 1 < D. Let $d(A_0, A) = 2p + 1$. By Lemma 2.2, $|A_0 \cap A| \le kp$. If $|A_0 \cap A| \le k(p-1)$, by the induction hypothesis, $d(A_0, A) \le 2(p-1) + 1$ which is a contradiction. Thus $|A_0 \cap A| \in N_{2p+1} = \{k(p-1) + 1, \dots, kp\}$.

Conversely, let $|A_0 \cap A| \in N_{2p+1} = \{k(p-1) + 1, \dots, kp\}$, so that $|A_0 \cap A| = kp - r$ where $0 \le r < k$. Set $Y' = A_0 \cap A$, so that $A_0 =$

The Wiener Number of Kneser Graphs

 $Y' \cup \{a_1, \ldots, a_{n-kp+r}\}$, and $A = Y' \cup \{b_1, \ldots, b_{n-kp+r}\}$ where the sets $\{a_1, \ldots, a_{n-kp+r}\}$ and $\{b_1, \ldots, b_{n-kp+r}\}$ are disjoint. Further, let $C' = [m] \setminus (A_0 \cup A)$. Then, |C'| = 2n + k - (2n - kp + r) = k(p+1) - r. Now choose k elements c'_1, c'_2, \ldots, c'_k from C' and set $B' = (C' \setminus \{c'_1, \ldots, c'_k\}) \cup \{a_1, \ldots, a_{n-kp+r}\}$. Hence |B'| = n and so $B' \in V(G)$, $|A_0 \cap B'| = n - kp + r$ and $|A \cap B'| = 0$. Since, $0 \le r < k$, by the induction hypothesis, $d(A_0, B') = 2p$ and d(A, B') = 1 and so, $d(A_0, A) \le 2p + 1$. If $d(A_0, A) < 2p + 1$, by the induction hypothesis, $|A_0 \cap A| \ne kp - r$, for any r in $0, 1, 2, \ldots, k - 1$, which is a contradiction. Thus $d(A_0, A) = 2p + 1$.

Case (ii). Let *i* be even so that $i = 2p < D, p \ge 2$. Let $d(A_0, A) = 2p$. By Lemma 2.1, $|A_0 \cap A| \ge n - kp$. If $|A_0 \cap A| \ge n - k(p-1)$, by the induction hypothesis, $d(A_0, A) \le 2(p-1)$, which is a contradiction. Thus $|A_0 \cap A| \in N_{2p} = \{n - kp, n - kp + 1, \dots, n - k(p-1) - 1\}$.

Conversely, let $|A_0 \cap A| \in N_{2p}$, so that $|A_0 \cap A| = n - kp + r$, where $0 \leq r < k$. Set $Y'' = A_0 \cap A$, $A_0 = Y'' \cup \{x_1, x_2, \dots, x_{kp-r}\}$, and $A = Y'' \cup \{y_1, y_2, \dots, y_{kp-r}\}$ where the set $\{x_1, x_2, \dots, x_{kp-r}\}$ is disjoint from $\{y_1, y_2, \dots, y_{kp-r}\}$. Further, if $B'' = Y'' \cup \{x_1, \dots, x_k, y_{k+1}, \dots, y_{kp-r}\}$, then |B''| = n and so $B'' \in V(G)$. Also $|A_0 \cap B''| = n - k(p-1) + r$, and $|A \cap B''| = n - k$. Since $0 \leq r < k$, by the induction hypothesis, $d(A_0, B'') = 2(p-1)$ and d(A, B'') = 2 and so $d(A_0, A) \leq 2p$. If $d(A_0, A) < 2p$, by the induction hypothesis, $|A_0 \cap A| \neq n - kp + r$ for any r in $0, 1, \dots, k-1$, which is a contradiction. Thus $d(A_0, A) = 2p$.

We have settled all the cases for which $d(A_0, A) \leq D - 1$. We now dispose of the case when $d(A_0, A) = D$. There are two possibilities according to whether D is even or odd.

Case (a). D is even. Let $d(A_0, A) = D$, so that $d(A_0, A) \not\leq D - 1$. Consequently,

$$|A_0 \cap A| \notin \bigcup_{i=2}^{D-1} N_i \cup \{0,n\} = \bigcup_{\substack{i \text{ even}\\2 \le i \le D-1}} N_i \bigcup_{\substack{j \text{ odd}\\2 \le j \le D-1}} N_j \cup \{0,n\}.$$

In 2 to D-1, there are $\frac{D}{2}-1$ odd numbers and $\frac{D}{2}-1$ even numbers. Hence

$$\bigcup_{\substack{i \text{ even} \\ 2 \le i \le D-1}} N_i = \bigcup_{i=1}^{\frac{D}{2}-1} N_{2i} = \left\{ n - k \left(\frac{D}{2} - 1\right), \dots, n-1 \right\},$$

and

$$\bigcup_{\substack{j \text{ odd} \\ 2 \le j \le D-1}} N_j = \bigcup_{j=1}^{\frac{D}{2}-1} N_{2j+1} = \left\{1, 2, \dots, k\left(\frac{D}{2}-1\right)\right\},\$$

and hence

$$A_0 \cap A| \in \left\{ k \left(\frac{D}{2} - 1 \right) + 1, \dots, n - k \left(\frac{D}{2} - 1 \right) - 1 \right\}.$$

Conversely, let

$$|A_0 \cap A| \in \left\{ k \left(\frac{D}{2} - 1 \right) + 1, \dots, n - k \left(\frac{D}{2} - 1 \right) - 1 \right\}.$$

Then $|A_0 \cap A| \notin \bigcup_{i=2}^{D-1} N_i \cup \{0, n\}$, and so $d(A_0, A) = D$.

Case (b). D is odd. In this case,

$$|A_0 \cap A| \notin \bigcup_{i=2}^{D-1} N_i \cup \{0,n\} = \bigcup_{\substack{i \text{ even}\\2 \le i \le D-1}} N_i \bigcup_{\substack{j \text{ odd}\\2 \le j \le D-1}} N_j \cup \{0,n\}.$$

In 2 to D-1, there are $\frac{D-1}{2}-1$ odd numbers and $\frac{D-1}{2}$ even numbers. Hence,

$$\bigcup_{\substack{i \text{ even} \\ 2 \le i \le D-1}} N_i = \bigcup_{i=1}^{\frac{D-1}{2}} N_{2i} = \left\{ n - k \left(\frac{D-1}{2} \right), \dots, n-1 \right\},$$

and

$$\bigcup_{\substack{j \text{ odd} \\ 2 \le j \le D-1}} N_j = \bigcup_{j=1}^{\frac{D-1}{2}-1} N_{2j+1} = \left\{1, 2, \dots, k\left(\frac{D-1}{2}-1\right)\right\}.$$

Therefore, $|A_0 \cap A| \in \{k(\frac{D-1}{2}-1)+1, \dots, n-k(\frac{D-1}{2})-1\}$. Conversely, if $|A_0 \cap A| \in \{k(\frac{D-1}{2}-1)+1, \dots, n-k(\frac{D-1}{2})-1\}$, then $|A_0 \cap A| \notin \bigcup_{i=2}^{D-1} N_i \cup \{0, n\}$, and hence $d(A_0, A) = D$.

Remark 3.3. Let $A_0 \in V(KG(m,n))$. Let $0 \leq j \leq n$. Then the number of vertices A of KG(m,n) such that $|A_0 \cap A| = j$ is equal to $\binom{n}{j}\binom{n+k}{n-j}$.

224

Theorem 3.4. The Wiener number W of the Kneser graph KG(m, n) is given by

$$W = \frac{1}{2} \binom{2n+k}{n} \left[\sum_{i=0}^{\lfloor \frac{D-1}{2} \rfloor} (2i) \sum_{j=n-ki}^{\min\{n-k(i-1)-1,n\}} \binom{n}{j} \binom{n+k}{n-j} + \sum_{i=0}^{\lfloor \frac{D-1}{2} \rceil-1} (2i+1) \sum_{j=\max\{k(i-1)+1,0\}}^{ki} \binom{n}{j} \binom{n+k}{n-j} + S \right],$$

where

$$S = \begin{cases} D \sum_{j=k(\frac{D}{2}-1)+1}^{n-k(\frac{D}{2}-1)-1} \binom{n}{j} \binom{n+k}{n-j} & \text{if } D \text{ is even,} \\ \\ D \sum_{j=k(\frac{D-1}{2}-1)+1}^{n-k(\frac{D-1}{2})-1} \binom{n}{j} \binom{n+k}{n-j} & \text{if } D \text{ is odd.} \end{cases}$$

(Note: the min and max symbols are used in the summation to take care of the case i = 0.)

Proof. Let A_0 be a fixed vertex of KG(m, n). By Lemma 3.1 and Remark 3.3, we see that the number of vertices at distance l < D is given by

$$\sum_{j=n-ki}^{\min\{n-k(i-1)-1,n\}} \binom{n}{j} \binom{n+k}{n-j}, \text{ if } l=2i, \text{ and}$$
$$\sum_{j=\max\{k(i-1)+1,0\}}^{ki} \binom{n}{j} \binom{n+k}{n-j}, \text{ if } l=2i+1.$$

and the number of vertices at distance ${\cal D}$ is given by

$$\sum_{\substack{j=k(\frac{D}{2}-1)+1\\j=k(\frac{D}{2}-1)+1}}^{n-k(\frac{D}{2}-1)-1} \binom{n}{j}\binom{n+k}{n-j}, \text{ if } D \text{ is even, and}$$
$$\sum_{\substack{j=k(\frac{D-1}{2}-1)+1\\j=k(\frac{D-1}{2}-1)+1}}^{n-k(\frac{D}{2}-1)-1} \binom{n}{j}\binom{n+k}{n-j}, \text{ if } D \text{ is is odd.}$$

Since KG(m, n) is vertex-transitive, we get the expression given in Theorem 3.4 for W.

We now deduce the Wiener number of the odd graphs O_k . The graph O_k is the Kneser graph KG(2k-1, k-1) so that it is obtained by setting k = 1and n = k - 1 in KG(m, n). By Lemma 2.3, the diameter of $O_k = D = \lfloor \frac{k-2}{1} \rfloor + 1 = k - 1$. Substituting k = 1 and n = k - 1 in Theorem 3.4, the Wiener number of odd graphs is given by

$$W(O_k) = \frac{1}{2} \binom{2k-1}{k-1} \left[\sum_{i=0}^{\lfloor \frac{D-1}{2} \rfloor} (2i) \sum_{j=k-1-i}^{k-1-i} \binom{k-1}{j} \binom{k}{k-1-j} + \sum_{i=0}^{\lfloor \frac{D-1}{2} \rfloor - 1} (2i+1) \sum_{j=i}^{i} \binom{k-1}{j} \binom{k}{k-1-j} + S \right]$$

where

(A)

$$S = \begin{cases} D \sum_{\substack{j=\frac{D}{2} \\ p=\frac{D}{2}}}^{k-1-\frac{D}{2}} \binom{k-1}{j} \binom{k}{k-1-j} & \text{if } D \text{ is even,} \\ D \sum_{\substack{j=\frac{D-1}{2} \\ p=\frac{D-1}{2}}}^{k-2-\frac{D-1}{2}} \binom{k-1}{j} \binom{k}{k-1-j} & \text{if } D \text{ is odd,} \end{cases}$$

$$= \frac{1}{2} \binom{2k-1}{k-1} \left[\sum_{i=0}^{\lfloor \frac{k-2}{2} \rfloor} (2i) \binom{k-1}{k-1-i} \binom{k}{i} + \sum_{i=0}^{\lfloor \frac{k-2}{2} \rfloor-1} (2i+1) \binom{k-1}{i} \binom{k-1}{k-1-i} + S \right],$$

where

$$S = \begin{cases} D\binom{k-1}{k-1-\frac{D}{2}}\binom{k}{\frac{D}{2}} & \text{if } D \text{ is even}, \\ \\ D\binom{k-1}{\left(\frac{D-1}{2}\right)}\binom{k}{k-1-\left(\frac{D-1}{2}\right)} & \text{if } D \text{ is odd}. \end{cases}$$

Now,

$$\binom{k-1}{k-1-i}\binom{k}{i} = \frac{(k-1)!}{i!(k-1-i)!} \frac{k!}{i!(k-i)!}$$
$$= \frac{k}{k-i} \frac{(k-1)!^2}{(i!)^2(k-i-1)!^2} = \frac{k(k-1)^2 \cdots (k-i+1)^2(k-i)}{1^2 \cdots i^2}$$

and

$$\binom{k-1}{i}\binom{k}{k-1-i} = \frac{(k-1)!}{i!(k-i-1)!} \frac{k!}{(1+i)!(k-1-i)!}$$
$$= \frac{k}{i+1} \frac{(k-1)!^2}{(i!)^2(k-i-1)!^2} = \frac{k(k-1)^2 \cdots (k-i)^2}{1^2 \cdots i^2(1+i)}.$$

When D is even, we can add S to the first summation by taking $i = \left\lceil \frac{k-2}{2} \right\rceil$ in (A) and when D is odd, to the second summation by taking $i = \left\lfloor \frac{k-2}{2} \right\rfloor$ in (A). This gives

$$W(O_k) = \frac{1}{2} \binom{2k-1}{k-1} \left[\sum_{i=0}^{\lceil \frac{k-2}{2} \rceil} (2i) \frac{k(k-1)^2 \cdots (k-i+1)^2 (k-i)}{1^2 \cdot 2^2 \cdots i^2} + \sum_{i=0}^{\lfloor \frac{k-2}{2} \rfloor} (2i+1) \frac{k(k-1)^2 \cdots (k-i)^2}{1^2 \cdot 2^2 \cdots i^2 \cdot (1+i)} \right].$$

Expression (B) has also been established by Tilakam [17] using intersection arrays [4]. Another equivalent expression for $W(O_k)$ is given in [1].

Acknowledgement

This research was supported by the Department of Science and Technology, Government of India grant DST SR/S4/MS: 234/04 dated March 31, 2006.

References

- R. Balakrishanan, N. Sridharan and K. Viswanathan, The Wiener index of odd graphs, Indian Inst. Sci. 86 (2006) 527–531.
- [2] R. Balakrishanan, K. Viswanathan and K.T. Raghavendra, Wiener index of two special trees, MATCH Commun. Math. Comupt. Chem. 57 (2007) 385–392.

- [3] R. Balakrishnan and K. Ranganathan, A Textbook of Graph Theory (Springer, New York, 2000).
- [4] N.L. Biggs, Algebraic Graph Theory (Cambridge University Press, London, 1974).
- [5] A.A. Dobrynin, R. Entringer and I. Gutman, Wiener index of trees: theory and applications, Acta Appl. Math. 66 (2001) 211–249.
- [6] A.A. Dobrynin, I. Gutman, S. Klavžar and P. Žigert, Wiener index of hexagonal systems, Acta Appl. Math. 72 (2002) 247–294.
- [7] P. Frankl and Z. Füredi, *Extremal problems concerning Kneser graphs*, J. Combin. Theory (B) 40 (1986) 270–284.
- [8] I. Gutman and O. Polansky, Mathematical Concepts in Organic Chemistry (Springer-Verlag, Berlin, 1986).
- H. Hajabolhassan and X. Zhu, Circular chromatic number of Kneser graphs, J. Combin. Theory (B) 881 (2003) 299–303.
- [10] A. Johnson, F.C. Holroyd, and S. Stahl, Multichormatic numbers, star chromatic numbers and Kneser graphs, J. Graph Theory 26 (1997) 137–145.
- [11] K.W. Lih and D.F. Liu, Circular chromatic number of some reduced Kneser graphs, J. Graph Theory 41 (2002) 62–68.
- [12] L. Lovász, Kneser's conjecture, chromatic number and homotopy, J. Combin. Theory (A) 25 (1978) 319–324.
- [13] M. Valencia-Pabon and J.-C. Vera, On the diameter of Kneser graphs, Discrete Math. 305 (2005) 383–385.
- [14] S.-P. Eu, B. Yang, and Y.-N Yeh, Generalised Wiener indices in hexagonal chains, Intl., J., Quantum Chem. 106 (2006) 426–435.
- [15] S. Stahl, n-tuple coloring and associated graphs, J. Combin. Theory (B) 20 (1976) 185–203.
- [16] S. Stahl, The multichromatic number of some Kneser graphs, Discrete Math. 185 (1998) 287–291.
- [17] K. Tilakam, Personal communication.
- [18] H. Wiener, Structural determination of Paraffin boiling points, J. Amer. Chem. Soc. 69 (1947) 17–20.
- [19] L. Xu and X. Guo, Catacondensed hexagonal systems with large Wiener numbers, MATCH Commun. Math. Comput. Chem. 55 (2006) 137–158.

Received 21 May 2007 Revised 18 February 2008 Accepted 20 February 2008