Discussiones Mathematicae Graph Theory 28 (2008) 335–343

# A NOTE ON DOMINATION PARAMETERS IN RANDOM GRAPHS

ANTHONY BONATO

Department of Mathematics Wilfrid Laurier University Waterloo, ON, Canada, N2L 3C5 e-mail: abonato@rogers.com

AND

CHANGPING WANG

Department of Mathematics Ryerson University Toronto, ON, Canada, M5B 2K3 e-mail: cpwang@ryerson.ca

#### Abstract

Domination parameters in random graphs G(n, p), where p is a fixed real number in (0, 1), are investigated. We show that with probability tending to 1 as  $n \to \infty$ , the total and independent domination numbers concentrate on the domination number of G(n, p).

**Keywords:** domination, random graphs, independent domination, total domination.

2000 Mathematics Subject Classification: 05C69, 05C80.

### 1. INTRODUCTION

Domination is a central topic in graph theory, with a number of applications in computer science and engineering. A set S of vertices in a graph Gis a *dominating set* of G if each vertex not in S is joined to some vertex of S.

The authors gratefully acknowledge support from NSERC and MITACS.

The domination number  $\gamma(G)$  is the minimum cardinality of a dominating set of G. The concentration of the domination number of random graphs G(n, p) was investigated in [8]. Other contributions to domination in random graph theory include [2, 6, 7]. For background on random graphs and domination, the reader is directed to [1, 5] and [3, 4], respectively. We say that an event holds asymptotically almost surely (a.a.s.) if the probability that it holds tends to 1 as n tends to infinity. All logarithms are in base e unless otherwise stated, and we use the notation  $\mathbb{L}n = \log_{1/(1-p)} n$ .

**Theorem 1** ([8]). For  $p \in (0, 1)$  fixed, a.a.s.  $\gamma(G(n, p))$  equals

$$|\mathbb{L}n - \mathbb{L}((\mathbb{L}n)(\log n))| + 1 \text{ or } |\mathbb{L}n - \mathbb{L}((\mathbb{L}n)(\log n))| + 2.$$

Despite the fact that deterministic graphs of order n may have domination number equalling  $\Theta(n)$  (such as a path  $P_n$  with  $\gamma(P_n) = \lceil n/3 \rceil$ ), Theorem 1 demonstrates that *a.a.s.* G(n,p) has domination number equalling  $(1+o(1))\mathbb{L}n = \Theta(\log n)$ .

A set S is said to be an *independent dominating set* of G if S is both an independent set and a dominating set of G (that is, S is a maximal independent set). A *total dominating set* S in a graph G is a subset of V(G)satisfying that every  $v \in V(G)$  is joined to at least one vertex in S. The *independent domination number* of G, written  $\gamma_i(G)$ , is the minimum order of an independent dominating set of G; the total domination number, written  $\gamma_t(G)$ , is defined analogously. It is straightforward to see that  $\gamma(G) \leq \gamma_i(G)$ and  $\gamma(G) \leq \gamma_t(G)$ . However, the domination number may be of much smaller order than either the independent or total domination numbers; see for example, [3, 4]. As proved in [9], there are cubic graphs where the difference between  $\gamma_i$  and  $\gamma$  is  $\Theta(n)$ .

Our goal in this note is to demonstrate that in G(n, p) with p fixed, asymptotically the independent and total domination numbers concentrate on  $(1 + o(1))\mathbb{L}n$ . In particular, we prove the following theorems.

**Theorem 2.** A.a.s.  $\gamma_t(G(n,p))$  equals

$$|\mathbb{L}n - \mathbb{L}((\mathbb{L}n)(\log n))| + 1 \text{ or } |\mathbb{L}n - \mathbb{L}((\mathbb{L}n)(\log n))| + 2.$$

Theorem 3. A.a.s. we have that

$$|\mathbb{L}n - \mathbb{L}((\mathbb{L}n)(\log n))| + 1 \le \gamma_i(G(n, p)) \le |\mathbb{L}n|.$$

As the proofs of the theorems are technical—though elementary—we present them in the next section. For both proofs, we compute the asymptotic expected value of each domination parameter, and then analyze its variance. The second moment method (see Chapter 4 of [1], for example) completes the proofs.

All graphs we consider are finite, undirected, and simple. If A is an event in a probability space, then we write  $\mathbb{P}(A)$  for the probability of A in the space. We use the notation  $\mathbb{E}(X)$  and Var(X) for the expected value and variance of a random variable X on G(n, p), respectively. Throughout, n is a positive integer, all asymptotics are as  $n \to \infty$ , and  $p \in (0, 1)$  is a fixed real number.

### 2. Proofs of Theorems 2 and 3

The proofs are presented in the following two subsections. We note the following facts from [8]. For  $r \ge 1$ , let  $X_r$  be the number of dominating sets of size r. Fix an r-set  $S_1$ . Denote by S(j) the set of r-sets which intersect  $S_1$  in j elements. Let  $I_1$  and  $I^j$  be indicator random variables, where the events  $I_1 = 1$  and  $I^j = 1$  represent that  $S_1$  and  $S_j \in S(j)$  are dominating sets, respectively. Let

$$A = \binom{n}{r} \sum_{j=0}^{r-1} \binom{r}{j} \binom{n-r}{r-j} \mathbb{E}(I_1 I^j).$$

**Lemma 4** ([8]). The random variable  $X_r$  satisfies the following properties.

(1)  $\mathbb{E}(X_r) = \binom{n}{r} (1 - (1 - p)^r)^{n-r}.$ (2) For  $r \ge \lfloor \mathbb{L}n - \mathbb{L}((\mathbb{L}n)(\log n)) \rfloor + 2$ , we have that  $\mathbb{E}(X_r) \to \infty$  as  $n \to \infty$ . (3) For  $r \ge \lfloor \mathbb{L}n - \mathbb{L}((\mathbb{L}n)(\log n)) \rfloor + 2$ ,

$$A \le \mathbb{E}^{2}(X_{r})\left(1 + 2r(1-p)^{r} - \frac{r^{2}}{n}\right)(1 + o(1)) + rg(1)\binom{n}{r},$$

where

(2.1) 
$$g(1) = \frac{2rn^{r-1}}{(r-1)!} \exp\left(n(1-p)^{2r-1} - 2(1-p)^r\right).$$

## 2.1. Proof of Theorem 2

For r a positive integer, the random variable  $X_r^t$  denotes the number of total dominating sets of size r. By Chebyshev's inequality, the proof of the theorem will follow once we show that  $\mathbb{E}(X_r^t) \to \infty$  as  $n \to \infty$ , and  $Var(X_r^t) = o(\mathbb{E}^2(X_r^t))$ . (See, for example, Section 4.3 of [1].)

**Lemma 5.** If  $r = \lfloor \mathbb{L}n - \mathbb{L}((\mathbb{L}n)(\log n)) \rfloor + 2$ , then

$$\mathbb{E}(X_r^t) = \binom{n}{r} (1 - (1 - p)^r)^{n - r} (1 - (1 - p)^{r - 1})^r (1 + o(1)).$$

**Proof.** For  $1 \leq j \leq {n \choose r}$ , denote by  $E_j$  the event that the subgraph induced by a given *r*-set  $S_j$  has no isolated vertices. We have that

$$\mathbb{E}(X_r^t) = \binom{n}{r} (1 - (1 - p)^r)^{n-r} \mathbb{P}(E_j).$$

It is not hard to show that for all j,  $\mathbb{P}(E_j) \geq 1 - r(1-p)^{r-1}$ , and so  $\lim_{n\to\infty} \mathbb{P}(E_j) = 1$ . The proof follows since for  $r = \lfloor \mathbb{L}n - \mathbb{L}((\mathbb{L}n)(\log n)) \rfloor + 2$ ,

$$\lim_{n \to \infty} (1 - (1 - p)^{r-1})^r = 1.$$

We next show that for a certain value of r, the expected value of  $X_r^t$  concentrates on the expected value of  $X_r$ .

**Lemma 6.** If 
$$r = \lfloor \mathbb{L}n - \mathbb{L}((\mathbb{L}n)(\log n)) \rfloor + 2$$
, then  $\mathbb{E}(X_r^t) = (1+o(1))\mathbb{E}(X_r)$ .

**Proof.** By Lemmas 4 and 5, we have that

$$\frac{\mathbb{E}(X_r^t)}{\mathbb{E}(X_r)} = (1 - (1 - p)^{r-1})^r (1 + o(1)).$$

Hence,

$$\lim_{n \to \infty} \frac{\mathbb{E}(X_r^t)}{\mathbb{E}(X_r)} = \lim_{n \to \infty} (1 - (1 - p)^{r-1})^r (1 + o(1)) = 1.$$

By Lemmas 4 and 6, the proof of the following lemma is immediate.

**Lemma 7.** If 
$$r = \lfloor \mathbb{L}n - \mathbb{L}((\mathbb{L}n)(\log n)) \rfloor + 2$$
, then  $\mathbb{E}(X_r^t) \to \infty$  as  $n \to \infty$ .

338

We now analyze the variance of the random variable  $X_r^t$ .

**Lemma 8.** If  $r = \lfloor \mathbb{L}n - \mathbb{L}((\mathbb{L}n)(\log n)) \rfloor + 2$ , then  $Var(X_r^t) = o(\mathbb{E}^2(X_r^t))$ .

**Proof.** For  $1 \leq j \leq {n \choose r}$ , let  $I^j$  be the corresponding indicator random variables. Hence,

$$X_r^t = \sum_{j=1}^{\binom{n}{r}} I^j.$$

By the linearity of expectation, we have that

(2.2)  
$$\mathbb{E}((X_r^t)^2) = \sum_{j=1}^{\binom{n}{r}} \mathbb{E}\left(\left(I^j\right)^2\right) + 2\sum_{\substack{j\neq i}}^{\binom{n}{r}} \mathbb{E}\left(I^i I^j\right)$$
$$= \mathbb{E}\left(X_r^t\right) + 2\sum_{\substack{j\neq i}}^{\binom{n}{r}} \mathbb{E}\left(I^i I^j\right).$$

We fix an r-set  $S_1$ . For  $0 \le j \le r-1$ , denote by S(j) the set of r-sets which intersect  $S_1$  in j elements. Let  $I_1^t$  and  $I^{jt}$  be the indicator random variables, where the events  $I_1^t = 1$  and  $I^{jt} = 1$  represent that  $S_1$  and  $S_j \in S(j)$  are total dominating sets, respectively. Then

$$2\sum_{j\neq i}^{\binom{n}{r}} \mathbb{E}(I^i I^j) = \binom{n}{r} \sum_{j=0}^{r-1} \binom{r}{j} \binom{n-r}{r-j} \mathbb{E}(I_1^t I^{jt}).$$

Together with (2.2), we obtain that

(2.3) 
$$\mathbb{E}((X_r^t)^2) = \mathbb{E}(X_r^t) + \binom{n}{r} \sum_{j=0}^{r-1} \binom{r}{j} \binom{n-r}{r-j} \mathbb{E}(I_1^t I^{jt})$$
$$= \mathbb{E}(X_r^t) + A^t,$$

where  $A^t = \binom{n}{r} \sum_{j=0}^{r-1} \binom{r}{j} \binom{n-r}{r-j} \mathbb{E}(I_1^t I^{jt})$ . As each total dominating set is a dominating set,  $A^t \leq A$ . By Lemmas 4 and 6, we therefore have that

(2.4) 
$$A^t \leq rg(1)\binom{n}{r} + (1+o(1))\mathbb{E}^2(X_r^t)\left(1+2r(1-p)^r - \frac{r^2}{n}\right),$$

where g(1) is given in (2.1).

By (2.3) and (2.4) we have that

(2.5) 
$$\frac{Var(X_r^t)}{\mathbb{E}^2(X_r^t)} \leq \frac{1}{\mathbb{E}(X_r^t)} + (1+o(1))\left(2r(1-p)^r - \frac{r^2}{n}\right) + \frac{rg(1)\binom{n}{r}}{\mathbb{E}^2(X_r^t)}.$$

To show that  $Var(X_r^t) = o(\mathbb{E}^2(X_r^t))$ , it suffices by Lemma 7 to show that

$$\frac{rg(1)\binom{n}{r}}{\mathbb{E}^2(X_r^t)} = o(1).$$

For sufficiently large n we have that

$$\frac{rg(1)\binom{n}{r}}{\mathbb{E}^{2}(X_{r}^{t})} = \frac{r \times 2r \frac{n^{r-1}}{(r-1)!} \exp\left(n\left((1-p)^{2r-1}-2(1-p)^{r}\right)\right)}{\binom{n}{r}\left((1-(1-p)^{r})^{n-r}\left(1-(1-p)^{r-1}\right)^{r}\right)^{2}} (1+o(1))$$

$$\leq \frac{3r^{3}}{n} \frac{\left(1-2(1-p)^{r}+(1-p)^{2r-1}\right)^{n}}{\left(1-(1-p)^{r}\right)^{2n-2r}\left(1-(1-p)^{r-1}\right)^{2r}} (1+o(1))$$

$$\leq \frac{3r^{3}}{n} \left(1+\frac{p(1-p)^{2r-1}}{\left(1-(1-p)^{r}\right)^{2}}\right)^{n-r} \left(1+\frac{2p(1-p)^{r-1}}{\left(1-(1-p)^{r-1}\right)^{2}}\right)^{r} (1+o(1)),$$

where the first equality follows by (2.1) and since  $\exp(x) \sim 1 + x$  if x is close to 0.

Since  $1 + x \leq \exp(x)$  for  $x \geq 0$ , we obtain that

$$\frac{rg(1)\binom{n}{r}}{\mathbb{E}^{2}(X_{r}^{t})} \leq \frac{3r^{3}}{n} \exp\left(\frac{(n-r)p(1-p)^{2r-1}}{(1-(1-p)^{r})^{2}} + \frac{2rp(1-p)^{r-1}}{(1-(1-p)^{r-1})^{2}}\right)(1+o(1))$$
$$\leq \frac{3r^{3}}{n} \exp\left((1+o(1))p\frac{(\mathbb{L}n)(\log n))^{2}}{n}\right)(1+o(1)) = o(1),$$

as  $r = \lfloor \mathbb{L}n - \mathbb{L}((\mathbb{L}n)(\log n)) \rfloor + 2.$ 

## 2.2. Proof of Theorem 3

We use the following lemma, whose proof is straightforward and so is omitted. For  $r \ge 1$ , let  $X_r^I$  be the random variable which denotes the number of independent dominating sets of size r.

340

Lemma 9. (1) For all  $r \ge 1$ 

$$\mathbb{E}(X_r^I) = \binom{n}{r} (1 - (1 - p)^r)^{n - r} (1 - p)^{\binom{r}{2}}.$$

(2) Let  $\lambda \in (\frac{1}{2}, 1)$  be fixed. For  $\lfloor \mathbb{L}n \rfloor + 1 \leq r \leq \lfloor 2\lambda \mathbb{L}n \rfloor$ , as  $n \to \infty$  we have that  $\mathbb{E}(X_r^I) \to \infty$ .

Our final lemma estimates the variance of  $X_r^I$ .

**Lemma 10.** Let  $p \in (0,1)$  and  $\lambda \in (\frac{1}{2},1)$  be fixed. For  $\lfloor \mathbb{L}n \rfloor + 1 \leq r \leq \lfloor 2\lambda \mathbb{L}n \rfloor$ ,

$$Var\left(X_{r}^{I}\right) = \mathbb{E}^{2}(X_{r}^{I})O\left(\frac{(\log n)^{4}}{n^{1-\lambda}}\right).$$

By Chebyshev's inequality and Lemmas 9 and 10, we have that

$$\mathbb{P}(\gamma_i(G) > r) = \mathbb{P}(X_r^I = 0) \le \mathbb{P}(|X_r^I - \mathbb{E}(X_r^I)| \ge \mathbb{E}(X_r^I))$$
$$\le \frac{Var(X_r^I)}{\mathbb{E}^2(X_r^I)} = o(1).$$

The assertion of Theorem 3 follows, therefore, once we prove Lemma 10.

**Proof of Lemma 10.** We denote by  $\mathbb{E}((X_r^I)^2)$  the expectation of the number of ordered pairs of independent domination sets of size r in  $G \in G(n, p)$ . The expectation satisfies

(2.6) 
$$\mathbb{E}\left(\left(X_{r}^{I}\right)^{2}\right) = \sum_{j} \binom{n}{r} \binom{r}{j} \binom{n-r}{r-j} \left(1 - (1-p)^{r}\right)^{2(n-2r+j)} \times \left(1 - (1-p)^{r-j}\right)^{2(r-j)} (1-p)^{2\binom{r}{2} - \binom{j}{2}}.$$

The explanations for the terms in the equation (2.6) are as follows. The vertices of the first independent dominating set  $S_1$  may be chosen in  $\binom{n}{r}$  ways. The independent dominating sets  $S_1$  and  $S_2$  may have j elements in common. These vertices may be chosen in  $\binom{r}{j}$  ways. The rest of r - j vertices of  $S_2$  may have to be chosen from  $V(G) \setminus S_1$ , which gives the  $\binom{n-r}{r-j}$  term. Every vertex not in  $S_1 \cup S_2$  must be joined to one of  $S_1$  and one of  $S_2$ , and so we obtain the term  $(1 - (1 - p)^r)^{2(n-2r+j)}$ . Every vertex in  $S_1 \setminus S_2$ 

must be joined to one of  $S_2 \setminus S_1$ , and every vertex in  $S_2 \setminus S_1$  must be joined to one of  $S_1 \setminus S_2$ , and so we have the term  $(1 - (1 - p)^{r-j})^{2(r-j)}$ . Both sets  $S_1$  and  $S_2$  are independent, which supplies the last term.

Observe that  $(1-p)^{r-j} \ge (1-p)^r$ . Hence, by (2.6) and Lemma 9 (1), we have that

(2.7)  

$$\mathbb{E}\left(\left(X_{r}^{I}\right)^{2}\right) \leq \mathbb{E}^{2}\left(X_{r}^{I}\right)\frac{1}{\binom{n}{r}}\left(\binom{n-r}{r}\right) + r\binom{n-r}{r-1} + \sum_{j=2}^{r}\binom{r}{j}\binom{n-r}{r-j}\left(1-p\right)^{-\binom{j}{2}}.$$

By the choice of r it follows that

(2.8) 
$$\frac{1}{\binom{n}{r}} \left( \binom{n-r}{r} + r\binom{n-r}{r-1} \right) = \left( 1 - \frac{r^2}{n} \right) \left( 1 + O\left(\frac{(\log n)^4}{n^2}\right) \right)$$
$$+ \frac{r^2}{n} + O\left(\frac{(\log n)^3}{n^2}\right)$$
$$= 1 + O\left(\frac{(\log n)^4}{n^2}\right),$$

and

(2.9) 
$$\frac{1}{\binom{n}{r}} \sum_{j=2}^{r} \binom{r}{j} \binom{n-r}{r-j} (1-p)^{-\binom{j}{2}} = O\left(\frac{(\log n)^4}{n^{1-\lambda}}\right).$$

By (2.7), (2.8), and (2.9) the assertion follows.

#### References

- [1] N. Alon and J. Spencer, The Probabilistic Method (Wiley, New York, 2000).
- [2] P.A. Dreyer, Applications and variations of domination in graphs, Ph.D. Dissertation, Department of Mathematics (Rutgers University, 2000).
- [3] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, Fundamentals of Domination in Graphs (Marcel Dekker, New York, 1998).
- [4] T.W. Haynes, S.T. Hedetniemi and P.J. Slater (eds.), Domination in Graphs: Advanced Topics (Marcel Dekker, New York, 1998).

- [5] S. Janson, T. Luczak and A. Ruciński, Random Graphs (John Wiley and Sons, New York, 2000).
- [6] C. Kaiser and K. Weber, Degrees and domination number of random graphs in the n-cube, Rostock. Math. Kolloq. 28 (1985) 18–32.
- [7] K. Weber, Domination number for almost every graph, Rostock. Math. Kolloq. 16 (1981) 31–43.
- [8] B. Wieland and A.P. Godbole, On the domination number of a random graph, The Electronic Journal of Combinatorics 8 (2001) #R37.
- [9] I.E. Zverovich and V.E. Zverovich, The domination parameters of cubic graphs, Graphs and Combinatorics 21 (2005) 277–288.

Received 11 January 2008 Revised 3 March 2008 Accepted 3 March 2008