THE SIGNED MATCHINGS IN GRAPHS

CHANGPING WANG

Department of Mathematics Ryerson University Toronto, ON, Canada, M5B 2K3

e-mail: cpwang@ryerson.ca

Abstract

Let G be a graph with vertex set V(G) and edge set E(G). A signed matching is a function $x : E(G) \to \{-1, 1\}$ satisfying $\sum_{e \in E_G(v)} x(e) \leq 1$ for every $v \in V(G)$, where $E_G(v) = \{uv \in E(G) | u \in V(G)\}$. The maximum of the values of $\sum_{e \in E(G)} x(e)$, taken over all signed matchings x, is called the signed matching number and is denoted by $\beta'_1(G)$. In this paper, we study the complexity of the maximum signed matching problem. We show that a maximum signed matching can be found in strongly polynomial-time. We present sharp upper and lower bounds on $\beta'_1(G)$ for general graphs. We investigate the sum of maximum size of signed matchings and minimum size of signed 1-edge covers. We disprove the existence of an analogue of Gallai's theorem. Exact values of $\beta'_1(G)$ of several classes of graphs are found.

Keywords: signed matching, signed matching number, maximum signed matching, signed edge cover, signed edge cover number, strongly polynomial-time.

2000 Mathematics Subject Classification: 05C70, 05C85.

1. INTRODUCTION

Structural and algorithmic aspects of covering vertices by edges have been extensively studied in graph theory. A matching (edge cover) of a graph G is a set C of edges of G such that each vertex of G is incident to at most (at least) one edge of C. Let b be a fixed positive integer. A simple

b-matching (simple *b*-edge cover) of a graph *G* is a set *C* of edges of *G* such that each vertex of *G* is incident to at most (at least) *b* edges of *C*. The maximum (minimum) size of a simple *b*-matching (simple *b*-edge cover) of *G* is called *b*-matching number (*b*-edge cover number), denoted by $\beta_b(G)$ ($\rho_b(G)$). The (simple) *b*-matching problems have been widely studied in, for instance, [1, 5, 6, 8]. For an excellent survey of results on matchings, edge covers, *b*-matchings and *b*-edge covers, see Schrijver [9].

Let G be a graph with vertex set V(G), edge set E(G) and maximum degree $\Delta(G)$. For a vertex $v \in V(G)$, let $E_G(v) = \{uv \in E(G) : u \in V(G)\}$ denote the set of edges of G incident to v. The degree, d(v), of v in G is $|E_G(v)|$. For a vertex $v \in V(G)$, v is called odd (even) if d(v) is odd (even). A k-factor of G is a k-regular spanning subgraph of G. In particular, F is a 1-factor of G if and only if E(F) is a perfect matching in G. Let $f: E(G) \to \mathbb{R}$ be a real-valued function. For $X \subseteq E(G)$, we write f(X) for $\sum_{e \in X} f(e)$. We use $G \cup H$ to denote the union of two disjoint graphs G and H, and we use $G \cong H$ to denote that G and H are isomorphic. We denote by \mathbb{N} the set of positive integers.

In this paper, we consider a variant of the standard matching problem. A signed matching of a graph G is a function $x : E(G) \to \{-1, 1\}$ satisfying $x (E_G(v)) \leq 1$ for every $v \in V(G)$. The maximum of the values of x (E(G)), taken over all signed matchings x, is called the signed matching number and is denoted by $\beta'_1(G)$. A maximum signed matching is a signed matching x satisfying $x (E(G)) = \beta'_1(G)$.

Let b be a fixed positive integer. A signed b-edge cover is a function $x : E(G) \to \{-1, 1\}$ satisfying $x(E_G(v)) \ge b$ for every $v \in V(G)$. The minimum of the values of x(E(G)), taken over all signed b-edge covers x, is called the signed b-edge cover number and is denoted by $\rho'_b(G)$.

The signed *b*-edge cover problem has been studied in [2]. In the special case when b = 1, ρ'_b is the signed star domination number investigated in [10, 12]. Other signed edge dominations have been investigated in [4, 7, 11, 13].

Note that $\beta'_1(G \cup H) = \beta'_1(G) + \beta'_1(H)$ and $\rho'_b(G \cup H) = \rho'_b(G) + \rho'_b(H)$ for two disjoint graphs G and H, and $\beta'_1(G) = |E(G)|$ for the graphs Gwith $\Delta(G) \leq 1$. Hence, we may assume that all graphs in this paper are connected with maximum degree greater than 1.

In Section 2, we investigate the complexity of the maximum signed matching problem. We show that a maximum signed matching can be found in polynomial time. In Section 3, we investigate the sum of maximum size of signed matchings and minimum size of signed 1-edge covers. We disprove the existence of an analogue of Gallai's theorem. In Section 4, we present sharp bounds on $\beta'_1(G)$ for general graphs. In Section 5, we study $\beta'_1(G)$ for some classes of graphs. Exact values of $\beta'_1(G)$ for paths, cycles, complete graphs and complete bipartite graphs are found.

All graphs considered in this paper are finite undirected graphs without loops or multiple edges. For all graph-theoretic terminology not defined here, the reader is directed to [3].

2. The Complexity of the Maximum Signed Matching Problem

An algorithm is said to run in *strongly polynomial time* if the number of elementary arithmetic and other operations is bounded by a fixed polynomial in the size of the input, where any number in the input is counted only for 1. Strongly polynomial time is of relevance only for algorithms that have numbers among their input; otherwise, strongly polynomial time coincides with the more well-known polynomial time. For more background on strongly polynomial time, the reader is referred to [9]. Our main result for this section is the following theorem.

Theorem 1. A maximum signed matching can be found in strongly polynomial time.

For the proof of Theorem 1, we use the following result from [9].

Theorem 2. Let $b: V(G) \to \mathbb{N}$ be an integer-valued function. A maximum b-matching can be found in strongly polynomial time.

Proof of Theorem 1. We may formulate the maximum signed matching problem to the following.

(2.1)
Maximize
$$\sum_{uv \in E(G)} x_{uv}$$

 $\sum_{uv \in E_G(u)} x_{uv} \le 1$, for every $u \in V(G)$,
 $x_{uv} \in \{-1, 1\}$, for every $uv \in E(G)$.

Notice that $x_{uv} \in \{-1, 1\}$ for each $uv \in E(G)$. So, if d(u) is even for some $u \in V(G)$, then $\sum_{uv \in E_G(u)} x_{uv} \leq 1$ implies that $\sum_{uv \in E_G(u)} x_{uv} \leq 0$.

Thus, (2.1) is equivalent to the following:

(2.2)

$$\begin{aligned}
\text{Maximize } \sum_{uv \in E(G)} x_{uv} \\
\sum_{uv \in E_G(u)} x_{uv} \leq 1, \text{ for every odd } u \in V(G), \\
\sum_{uv \in E_G(u)} x_{uv} \leq 0, \text{ for every even } u \in V(G), \\
x_{uv} \in \{-1, 1\}, \text{ for every } uv \in E(G).
\end{aligned}$$

Now we define $y_{uv} = \frac{1}{2}(1 + x_{uv})$ for each $uv \in E(G)$. It is clear that $y_{uv} \in \{0,1\}$ for each $uv \in E(G)$. Moreover, (2.2) is equivalent to the following:

(2.3)

$$Maximize -|E(G)| + 2 \sum_{uv \in E(G)} y_{uv}$$

$$\sum_{uv \in E_G(u)} y_{uv} \leq \frac{1}{2} (1 + d(u)), \text{ for every odd } u \in V(G),$$

$$\sum_{uv \in E_G(u)} y_{uv} \leq \frac{1}{2} d(u), \text{ for every even } u \in V(G),$$

$$y_{uv} \in \{0, 1\}, \text{ for every } uv \in E(G).$$

Define

$$b(u) = \begin{cases} \frac{1+d(u)}{2}, & d(u) \equiv 1 \pmod{2}, \\ \frac{d(u)}{2}, & d(u) \equiv 0 \pmod{2}. \end{cases}$$

By Theorem 2, we know that (2.3) is polynomial-time solvable.

3. $\rho'_1(G) + \beta'_1(G)$ for General Graphs

Let $k \ge 0$ be any integer. We define a family of trees named T_k as follows. Let $v_0, v_1, \ldots, v_{k+1}$ be a path with length k + 1. For each of the vertices v_0 and v_{k+1} , we add three adjacent leaves. For each of the remaining vertices (if exist) on the path, we add four adjacent leaves. Clearly, T_k is a tree of order 5k + 8. See Figure 1 for drawings of such trees T_0 and T_2 .

480



Figure 1. The trees T_0 and T_2 .

The following result (see [9, Theorem 34.1, p. 165, Vol. A]) is a direct analogue of Gallai's theorem, relating maximum-size *b*-matchings and minimum-size *b*-edge covers.

Theorem 3. Fix b a positive integer. If G is a graph of order n having no isolated vertices, then

$$\rho_b(G) + \beta_b(G) = bn.$$

Surprisingly, our next theorem exhibits that there is no such analogue of Gallai's theorem relating maximum-size signed matchings and minimum-size signed 1-edge covers.

Theorem 4. Let $k \ge 0$ be any integer. For the graph T_k of order n = 5k+8,

$$\rho_1'(T_k) + \beta_1'(T_k) = 4k + 6.$$

Proof. To show that $\rho'_1(T_k) + \beta'_1(T_k) = 4k+6$, we will show $\rho'_1(T_k) = 3k+5$ and $\beta'_1(T_k) = k+1$ in the following.

Let $v_0, v_1, \ldots, v_{k+1}$ be a path with length k+1 in T_k . Notice that T_k has 4k+6 leaves. Let x be a signed 1-edge cover of T_k such that $x(E(T_k)) = \rho'_1(T_k)$. By the definition of $\rho'_1(G)$, we must assign 1 to every leaf of T_k . To minimize $x(E(T_k))$, we should assign -1 to each edge $v_i v_{i+1}$ where $0 \le i \le k$. Hence, $\rho'_1(T_k) = 3k+5$.

Now we prove $\beta'_1(T_k) = k + 1$. Let y be a signed matching of T_k . Notice that the vertex v_i has an even degree for every $0 \le i \le k + 1$. Hence, $y(E_{T_k}(v_i)) \le 0$ for every $0 \le i \le k + 1$. By simple calculations, we know that

$$y(E(T_k)) = \sum_{i=0}^{k+1} y(E_{T_k}(v_i)) - \sum_{i=0}^k y(v_i v_{i+1})$$
$$\leq -\sum_{i=0}^k y(v_i v_{i+1}) \leq k+1.$$

Hence, $\beta'_1(T_k) \leq k+1$. To show that $\beta'_1(T_k) \geq k+1$, it suffices to produce a signed matching y' of T_k such that $y'(E(T_k)) = k+1$. We assign $y'(v_iv_{i+1}) = -1$ for $0 \leq i \leq k$. For each v_i , $0 \leq i \leq k+1$, note that v_i is even, it is possible to assign -1 and 1 appropriately so that $y'(E_{T_k}(v_i)) = 0$. It is not hard to verify that y' is a signed matching of T_k satisfying $y'(E(T_k)) = k+1$, which completes the proof.

It would be interesting to determine sharp lower and upper bounds on $\rho'_1(G) + \beta'_1(G)$ for general graphs G. We leave this as an open problem.

4. Lower and Upper Bounds

In this section, we present sharp lower and upper bounds on the signed matching number of general graphs.

Theorem 5. Let G be a graph of order n with k odd vertices. Then

$$\beta_1'(G) \le k/2.$$

The bound is sharp by Theorem 9.

Proof. Let $S \subseteq V(G)$ be the set of odd vertices. By the fact that every graph has even number of odd vertices, so |S| = k is even.

Taking any signed matching x of G, by the definition, we have that for every $v \in S$, $x(E_G(v)) \leq 1$, and otherwise $x(E_G(v)) \leq 0$. Thus, $\sum_{v \in V(G)} x(E_G(v)) \leq k$, i.e., $2x(E(G)) \leq k$. The proof is now complete.

Theorem 6. For any graph G of order n,

$$\beta_1'(G) \ge -1$$

The bound is sharp.

482

Proof. To start our proof, we construct a graph H from G as follows. If G is eulerian, then H = G. If not, as every graph has even number of odd vertices (say G has k odd vertices), then H is obtained by adding $\frac{k}{2}$ new vertices $w_1, \ldots, w_{\frac{k}{2}}$ to G, and joining each w_i to two distinct odd vertices of G, so that for all $i \neq j$ the vertices w_i and w_j have distinct neighbours. It is clear that H has no odd vertices and each w_i ($1 \leq i \leq \frac{k}{2}$) has degree 2. By Theorem 4.1 on page 94 of [3], H is eulerian. Let C be an eulerian circuit of H. We start with an edge $e \in E(G)$ and assign values -1 and +1 alternately along C. This defines a function $x : E(H) \to \{-1, 1\}$.

It is not hard to see that for every $v \in V(H)$,

$$(4.1) x (E_H(v)) \le 0,$$

and

(4.2)
$$x(E(G)) = x(E(H)) \ge -1.$$

Let x' be the restriction of x on E(G), i.e., x'(e) = x(e) for every $e \in E(G)$. By (4.1) and (4.2), we have that for every $v \in V(G)$, $x'(E_G(v)) \leq 1$, and $x'(E(G)) \geq -1$. Hence, x' is a signed matching of G satisfying that $x'(E(G)) \geq -1$. So, $\beta'_1(G) \geq -1$. As $\beta'_1(C_{2k+1}) = -1$ for any positive integer k, the bound is sharp.

The following result follows immediately from Theorems 5 and 6.

Corollary 7. Let G be eulerian of order n and size m. Then

$$\beta'_1(G) = \frac{1}{2} \left((-1)^m - 1 \right).$$

5. EXACT VALUES FOR CLASSES OF GRAPHS

The following theorem is an easy exercise and is left to the reader.

Theorem 8. Let $n \geq 3$ be an integer. We have

(1)
$$\beta_1'(C_n) = \begin{cases} -1, & n \equiv 1 \pmod{2}, \\ 0, & n \equiv 0 \pmod{2}. \end{cases}$$

(2)
$$\beta_1'(P_n) = \begin{cases} 0, & n \equiv 1 \pmod{2}, \\ 1, & n \equiv 0 \pmod{2}. \end{cases}$$

Our next theorems provide the exact values for complete graphs and complete bipartite graphs.

Theorem 9. Let $n \ge 2$ be an integer. We have the following

$$\beta_1'(K_n) = \begin{cases} n/2, & n \equiv 0, 2 \pmod{4}, \\ 0, & n \equiv 1 \pmod{4}, \\ -1, & n \equiv 3 \pmod{4}. \end{cases}$$

Proof. We consider three cases.

Case 1. n = 2k for some positive integer k. By Theorem 9.19 on page 273 of [3], K_n is 1-factorable. Note that K_n can be decomposed into (2k - 1) 1-factors. So, we can assign x(e) = 1 for each edge e of k 1-factors and assign x(e') = -1 for each edge e' of the remaining k - 1 1-factors. It is straightforward to verify that x is a signed matching of K_n and $x(E(K_n)) = k = \frac{n}{2}$. Hence, $\beta'_1(K_n) \ge \frac{n}{2}$. Note that $\beta'_1(K_n) \le \frac{n}{2}$ by Theorem 5. Thus, $\beta'_1(K_n) = \frac{n}{2}$.

Case 2. n = 4k + 1 for some positive integer k. It is not hard to see that K_n is Eulerian with size m = 2k(4k+1). It follows from Corollary 7 that $\beta'_1(K_n) = 0$.

Case 3. n = 4k + 3 for some positive integer k. Similar to Case 2, K_n is Eulerian with size m = (2k+1)(4k+3). It follows from Corollary 7 that $\beta'_1(K_n) = -1$.

For the graph $K_{1,t}$, one can show that

$$\beta_1'(K_{1,t}) = \begin{cases} 0, & t \equiv 0 \pmod{2}, \\ 1, & t \equiv 1 \pmod{2}. \end{cases}$$

In general, we prove the following theorem.

THE SIGNED MATCHINGS IN GRAPHS

Theorem 10. For positive integers $s \ge 2$ and $t \ge 2$,

$$\beta_1'(K_{s,t}) = \begin{cases} 0, & st \equiv 0 \pmod{2}, \\ \min\{s,t\}, & st \equiv 1 \pmod{2}. \end{cases}$$

Proof. Let $S = \{u_1, \ldots, u_s\}$ and $T = \{v_1, \ldots, v_t\}$ be the partite sets of $K_{s,t}$. We discuss two cases.

Case 1. st is even.

Without loss of generality we assume that s is even. Let x be any signed matching of $K_{s,t}$. By the definition, we have

$$x\left(E_{K_{s,t}}(v)\right) \le 0$$

for every $v \in T$. Hence, $x(E(K_{s,t})) \leq 0$ implying that $\beta'_1(K_{s,t}) \leq 0$.

To show that $\beta'_1(K_{s,t}) = 0$, it suffices to show that there exists a signed matching x such that $x(E(K_{s,t})) = 0$. In fact, x is such a signed matching if $x(u_i v_j) = (-1)^{i+j}$ for all $1 \le i \le s$ and $1 \le j \le t$.

Case 2. Both s and t are odd.

Observe that $\beta'_1(K_{s,t}) \leq \min\{s,t\}$. To show the other direction, it suffices to show that there exists a signed matching x such that $x(E(K_{s,t})) = \min\{s,t\}$.

Case 2.1. s = t.

By Theorem 9.18 on page 272 of [3], the complete bipartite graph $K_{s,s}$ is 1-factorable. In this case $K_{s,s}$ can be decomposed into s 1-factors. Assigning 1 to each edge of $\frac{s+1}{2}$ 1-factors and -1 to each edge of the remaining $\frac{s-1}{2}$ 1-factors, we produce a signed matching x such that $x(E(K_{s,t})) = s$.

Case 2.2. s < t.

Let G and H be the subgraphs induced by the edge sets $\bigcup_{i=1}^{s} \bigcup_{j=1}^{s} \{u_i v_j\}$ and $\bigcup_{i=1}^{s} \bigcup_{j=s+1}^{t} \{u_i v_j\}$, respectively.

Clearly, $E(K_{s,t}) = E(G) \cup E(H)$, $G \cong K_{s,s}$ and $H \cong K_{s,t-s}$. As we have proven in Case 2.1, there exists a signed matching x_G of G such that $x_G(E(G)) = s$. Now we can obtain a signed matching x of $K_{s,t}$ by setting $x(e) = x_G(e)$ for every $e \in E(G)$ and $x(u_i v_j) = (-1)^{i+j}$ for all $1 \le i \le s$ and $s + 1 \le j \le t$. By simple calculations, we have $x(E(K_{s,t})) = s$.

Case 2.3. s > t.

The proof is similar to that of Case 2.2, so we omit it.

References

- R.P. Anstee, A polynomial algorithm for b-matchings: an alternative approach, Inform. Process. Lett. 24 (1987) 153–157.
- [2] A. Bonato, K. Cameron and C. Wang, Signed b-edge covers of graphs, submitted.
- [3] G. Chartrand and L. Lesniak, Graphs & Digraphs, third edition (Chapman and Hall, Boca Raton, 2000).
- [4] W. Chena and E. Song, Lower bounds on several versions of signed domination number, Discrete Math. 308 (2008) 1837–1846.
- [5] S. Goodman, S. Hedetniemi and R.E. Tarjan, *b-matchings in trees*, SIAM J. Comput. 5 (1976) 104–108.
- [6] D. Hausmann, Adjacent vertices on the b-matching polyhedron, Discrete Math. 33 (1981) 37–51.
- [7] H. Karami, S.M. Sheikholeslami and A. Khodkar, Some notes on signed edge domination in graphs, Graphs and Combin. 24 (2008) 29–35.
- [8] W.R. Pulleyblank, Total dual integrality and b-matchings, Oper. Res. Lett. 1 (1981/82) 28-33.
- [9] A. Schrijver, Combinatorial Optimization: polyhedra and efficiency (Berlin, Springer, 2004).
- [10] C. Wang, The signed star domination numbers of the Cartesian product graphs, Discrete Appl. Math. 155 (2007) 1497–1505.
- B. Xu, On signed edge domination numbers of graphs, Discrete Math. 239 (2001) 179–189.
- B. Xu, Note On edge domination numbers of graphs, Discrete Math. 294 (2005) 311–316.
- [13] B. Xu, Two classes of edge domination in graphs, Discrete Appl. Math. 154 (2006) 1541–1546.

Received 18 December 2007 Revised 8 May 2008 Accepted 28 July 2008