

**FORBIDDEN-MINOR CHARACTERIZATION FOR THE  
CLASS OF GRAPHIC ELEMENT SPLITTING  
MATROIDS**

KIRAN DALVI\*, Y.M. BORSE\*\* AND M.M. SHIKARE\*\*

*\*Department of Mathematics*  
*Government College of Engineering, Pune 411 005, India*

**e-mail:** kiran\_dalvi111@yahoo.com

*\*\*Department of Mathematics*  
*University of Pune, Pune 411 007, India*

**e-mail:** ymborse@math.unipune.ernet.in

**e-mail:** mms@math.unipune.ernet.in

**Abstract**

This paper is based on the element splitting operation for binary matroids that was introduced by Azadi as a natural generalization of the corresponding operation in graphs. In this paper, we consider the problem of determining precisely which graphic matroids  $M$  have the property that the element splitting operation, by every pair of elements on  $M$  yields a graphic matroid. This problem is solved by proving that there is exactly one minor-minimal matroid that does not have this property.

**Keywords:** binary matroid, graphic matroid, minor, splitting operation, element splitting operation.

**2000 Mathematics Subject Classification:** 05B35.

1. INTRODUCTION

Let  $M(G)$  and  $M^*(G)$  denote the circuit matroid and the cocircuit matroid, respectively of a graph  $G$ . A matroid is *Eulerian* if its ground set can be expressed as a union of disjoint circuits of the matroid (see [14]). A matroid is *bipartite* if every circuit of it has an even number of elements. Welsh [14]

proved that a binary matroid is Eulerian if and only if its dual is bipartite. As the matroids  $F_7$  and  $M(K_5)$  are Eulerian, their dual matroids  $F_7^*$  and  $M^*(K_5)$  are bipartite. It is easy to see that a binary matroid  $M$  is Eulerian iff the sum of column vectors of  $A$  is zero where  $A$  is a matrix over  $GF(2)$  that represents  $M$ . For undefined notation and terminology in graphs and matroids, we refer [6] and [8].

Fleischner [3] defined the *splitting operation* for a graph with respect to a pair of adjacent edges as follows: Let  $G$  be a connected graph and  $v$  be a vertex of degree at least three in  $G$ . If  $x = uv$  and  $y = wv$  are two edges incident at  $v$ , then splitting away the pair  $x, y$  from  $v$  results in a new graph  $G_{x,y}$  obtained from  $G$  by deleting the edges  $x$  and  $y$ , and adding a new vertex  $v_{x,y}$  adjacent to  $u$  and  $w$ . The transition from  $G$  to  $G_{x,y}$  is called the splitting operation on  $G$ . For practical purposes, we denote the new edges  $v_{x,y}u$  and  $v_{x,y}w$  in  $G_{x,y}$  by  $x$  and  $y$ , respectively (See Figure 1). Fleischner [3] characterized Eulerian graphs and developed an algorithm to find all distinct Eulerian trails in an Eulerian graph using the splitting operation.




Figure 1

In a similar way, Tutte [13] specified the *point splitting operation* for graphs as follows: Let  $G$  be a graph and  $v$  be a vertex of degree at least 4 in  $G$ . Let  $H$  be the graph obtained from  $G$  by replacing  $v$  by two adjacent vertices  $v_1, v_2$  such that each point formerly joined to  $v$  is joined to exactly one of  $v_1$  and  $v_2$  so that in  $H$ ,  $\deg v_1 \geq 3$  and  $\deg v_2 \geq 3$ . We say that  $H$  arises from  $G$  by point-splitting operation. Tutte [13] characterized 3-connected graphs using this operation. Later on, Slater [12] classified 4-connected graphs using  $n$ -point splitting operation which is a natural generalization of the point splitting operation.

Azadi [1] defined an operation which, in a sense, combines the splitting operation and the point splitting operation as follows: Let  $v$  be a vertex of  $G$  and let  $x, y$  be distinct edges of  $G$  incident at  $v$ . Let  $G'_{x,y}$  be the graph obtained from  $G$  such that  $G'_{x,y} = G_{x,y} + v_{x,y}v$ , where  $G_{x,y}$  is the graph obtained from  $G$  by splitting operation with respect to the edges  $x$  and  $y$ .

Then we say that  $G'_{x,y}$  is obtained from  $G$  by the *element splitting operation with respect to the pair* of edges  $x$  and  $y$  (see Figure 2).




Figure 2

Raghunathan *et al.* [7] extended the definition of Fleischner's splitting operation to binary matroids as follows: Let  $A$  be a matrix over  $GF(2)$  that represents the matroid  $M$ . Consider distinct elements  $x$  and  $y$  of  $M$ . Let  $A_{x,y}$  be the matrix that is obtained by adjoining an extra row to  $A$  with this row being zero everywhere except in the columns corresponding to  $x$  and  $y$  where it takes the value 1. Suppose  $M_{x,y}$  is the matroid represented by the matrix  $A_{x,y}$ . Then  $M_{x,y}$  is said to be obtained from  $M$  by *splitting* away the pair  $x, y$ . Various properties concerning the splitting matroid have been studied in [2, 7, 9, 10, 11].

Azadi [1] further extended the operation of element splitting with respect to the pair of edges in graphs to binary matroids as follows: Let  $A$  be a matrix over  $GF(2)$  that represents the matroid  $M$ . Suppose that  $x$  and  $y$  are distinct elements of  $M$ . Let  $A'_{x,y}$  be the matrix that is obtained by adjoining an extra row to  $A$  with this row being zero everywhere except in the columns corresponding to  $x$  and  $y$  where it takes the value 1 and then adjoining an extra column (corresponding to  $a$ ) with this column being zero everywhere except in the last row where it takes the value 1. Suppose  $M'_{x,y}$  is the matroid represented by the matrix  $A'_{x,y}$ . Then  $M'_{x,y}$  is said to be obtained from  $M$  by *element splitting* the pair of elements  $x$  and  $y$ .

Alternatively, the element splitting operation can be defined in terms of circuits of binary matroids [1] as follows:

Let  $M = (S, \mathcal{C})$  be a binary matroid,  $\{x, y\} \subseteq S$ , and  $a \notin S$ . Let

$$\mathcal{C}_0 = \{C \in \mathcal{C} : x, y \in C \text{ or } x, y \notin C\},$$

$\mathcal{C}_1 =$  set of minimal members of  $\{C_1 \cup C_2 : C_1, C_2 \in \mathcal{C}, C_1 \cap C_2 = \phi \text{ and } x \in C_1, y \in C_2 \text{ such that } C_1 \cup C_2 \text{ does not contain any member of } \mathcal{C}_0\}$ , and

$$\mathcal{C}_2 = \{C \cup \{a\} : C \in \mathcal{C} \text{ and contains exactly one of } x \text{ and } y\}.$$

Let  $\mathcal{C}' = \mathcal{C}_0 \cup \mathcal{C}_1 \cup \mathcal{C}_2$ . Then  $M'_{x,y} = (S \cup \{a\}, \mathcal{C}')$ .

If  $x$  and  $y$  are non-adjacent edges of a graph  $G$ , then  $M(G)_{x,y}$  may not

be graphic. Shikare and Waphare [11] characterized graphic matroids whose splitting matroids are also graphic in the following theorem.

**Theorem 1.1** [11]. *The splitting operation, by any pair of elements, on a graphic matroid yields a graphic matroid if and only if the circuit matroid of the corresponding graph has no minor isomorphic to the circuit matroid of any of the following four graphs.*




Figure 3



The element splitting operation on a graphic matroid may not yield a graphic matroid. In this paper, we obtain the forbidden-minor characterization for graphic matroids whose element splitting matroid is graphic. The main result in this paper is the following theorem.

**Theorem 1.2.** *The element splitting operation, by any pair of elements, on a graphic matroid yields a graphic matroid if and only if it has no minor isomorphic to  $M(K_4)$ , where  $K_4$  is the complete graph on 4 vertices.*

## 2. THE ELEMENT SPLITTING OPERATION AND ITS PROPERTIES

In this section we provide necessary lemmas. We assume that  $M$  is a binary matroid and  $x, y$  are distinct elements of  $M$ .

**Lemma 2.1.** *Let  $x$  and  $y$  be elements of a binary matroid  $M$  and let  $r(M)$  denote the rank of  $M$ . Then, using the notations introduced in Section 1,*

- (i)  $M_{x,y} = M'_{x,y} \setminus \{a\}$ ;
- (ii)  $M = M'_{x,y} / \{a\}$ ;
- (iii)  $r(M'_{x,y}) = r(M) + 1$ ;
- (iv) every cocircuit of  $M$  is a cocircuit of the matroid  $M'_{x,y}$ ;

- (v) if  $\{x, y\}$  is a cocircuit of  $M$  then  $\{a\}$  and  $\{x, y\}$  are cocircuits of  $M'_{x,y}$ ;
- (vi) if  $\{x, y\}$  does not contain a cocircuit, then  $\{x, y, a\}$  is a cocircuit of  $M'_{x,y}$ ;
- (vii)  $M'_{x,y} \setminus x/y \cong M \setminus x$ ;
- (viii) if  $M$  is graphic and  $x, y$  are adjacent edges in a corresponding graph, then  $M'_{x,y}$  is graphic;
- (ix)  $M'_{x,y}$  is not eulerian.

**Proof.** (i), (ii), (iii), (v), (vi), (vii) and (viii) are straightforward. The proof of (iv) follows from Lemma 2.4.1 of [4]. If  $A'_{x,y}$  represents the matroid  $M'_{x,y}$ , then the number of one's in the last row of  $A'_{x,y}$  is odd. Hence  $M'_{x,y}$  is not eulerian. This proves (ix). ■

The following result is well known.

**Lemma 2.2** [6]. *A binary matroid is graphic if and only if it has no minor isomorphic to  $F_7, F_7^*, M^*(K_5)$ , or  $M^*(K_{3,3})$ .* ■

**Notation.** For convenience, let  $\mathcal{F} = \{F_7, F_7^*, M^*(K_5), M^*(K_{3,3})\}$ .

**Lemma 2.3.** *Let  $M$  be a graphic matroid and let  $x, y \in E(M)$  such that  $M'_{x,y}$  is not graphic. Then there is a minor  $N$  of  $M$  such that no two elements of  $N$  are in series and  $N'_{x,y} \setminus \{a\}/\{x\} \cong F$  or  $N'_{x,y} \setminus \{a\}/\{x, y\} \cong F$  or  $N'_{x,y} \cong F$  or  $N'_{x,y}/\{x\} \cong F$  or  $N'_{x,y}/\{y\} \cong F$  or  $N'_{x,y}/\{x, y\} \cong F$  for some  $F \in \mathcal{F}$ .*

**Proof.** Since  $M'_{x,y}$  is not graphic,  $M'_{x,y} \setminus T_1/T_2 \cong F$  for some  $T_1, T_2 \subseteq E(M'_{x,y})$ . Let  $T'_i = T_i - \{a, x, y\}$  for  $i = 1, 2$ . Then  $T'_i \subseteq E(M)$  for each  $i$ . Let  $N = M \setminus T'_1/T'_2$ . Then  $N'_{x,y} = M'_{x,y} \setminus T'_1/T'_2$ . Let  $T''_i = T_i - T'_i$  for  $i = 1, 2$ . Then  $N'_{x,y} \setminus T''_1/T''_2 \cong F$ . If  $a \in T''_2$ , then  $F$  is a minor of  $M'_{x,y}/a$  and hence, by Lemma 2.1(i),  $F$  is a minor of  $M$ , which is a contradiction. Suppose  $a \in T''_1$ . By Lemma 2.1(i),  $M_{x,y} = M'_{x,y} \setminus a$ . Hence  $F$  is a minor of  $M_{x,y}$ . It follows from Theorem 2.3 of [11] that  $N$  does not contain a 2-cocircuit and further,  $N_{x,y}/x \cong F$  or  $N_{x,y}/\{x, y\} \cong F$ . This implies that  $N'_{x,y} \setminus \{a\}/x \cong F$  or  $N'_{x,y} \setminus \{a\}/\{x, y\} \cong F$ . Suppose that  $a \notin T''_1 \cup T''_2$ . Hence  $a \notin T_1 \cup T_2$ . If  $T''_1 \cup T''_2 = \emptyset$ , then  $N'_{x,y} \cong F$ . If  $T''_2 = \emptyset$ , then  $N_{x,y} \setminus x \cong F$  or  $N_{x,y} \setminus y \cong F$  or  $N'_{x,y} \setminus \{x, y\} \cong F$ . In the first case,  $a$  forms a 2-cocircuit with  $x$  or  $y$  whichever is remained, and in later case,  $a$  is a coloop. It is a contradiction.

Hence  $T_2'' \neq \phi$ . If  $T_1'' \neq \phi$  then, by Lemma 2.1(vi),  $F$  is minor of  $M$ , which is a contradiction. Hence  $T_1'' = \phi$ . Hence  $N'_{x,y}/x \cong F$  or  $N'_{x,y}/y \cong F$  or  $N'_{x,y}/\{x,y\} \cong F$ .

Assume that  $N$  contains a 2-cocircuit  $Q$ . By Lemma 2.1(iv),  $Q$  is 2-cocircuit in  $N'_{x,y}$ . Since  $F$  is 3-connected, it does not contain a 2-cocircuit. It follows that  $N'_{x,y}$  is not isomorphic to  $F$ . Hence  $N'_{x,y} \setminus \{a\}/x \cong F$  or  $N'_{x,y} \setminus \{a\}/\{x,y\} \cong F$  or  $N'_{x,y}/\{x\} \cong F$  or  $N'_{x,y}/\{y\} \cong F$  or  $N'_{x,y}/\{x,y\} \cong F$ . If  $Q \cap \{x,y\} = \phi$ , then it is retained in all these cases and thus  $F$  has a 2-cocircuit, which is a contradiction. If  $Q = \{x,y\}$ , a contradiction follows from Lemma 2.1(v). Hence  $Q$  contains exactly one of  $x,y$ . Suppose that  $x \in Q$ . Then  $N'_{x,y}/y \not\cong F$ . Let  $x_1$  be the other element of  $Q$ . Let  $L = N/x_1$ . Then  $L$  is a minor of  $M$  in which no pair of elements is in series. Further,  $L'_{x,y} = N'_{x,y}/x_1 \cong N'_{x,y}/x$ . Thus we have  $L'_{x,y} \setminus \{a\} \cong F$  or  $L'_{x,y} \setminus \{a\}/y \cong F$  or  $L'_{x,y} \cong F$  or  $L'_{x,y}/y \cong F$ . Since  $L_{x,y} \cong L'_{x,y} \setminus \{a\}$ , and  $x,y$  are in series in  $L_{x,y}$ , it follows that  $L'_{x,y} \setminus \{a\} \not\cong F$  and also  $L'_{x,y} \setminus \{a\}/y \cong L'_{x,y} \setminus \{a\}/x$ . If  $y \in Q$ , then  $N'_{x,y}/x \not\cong F$ . Also,  $L'_{x,y} \cong N'_{x,y}/y$ . In this case we get  $L'_{x,y} \setminus \{a\}/x \cong F$  or  $L'_{x,y} \cong F$  or  $L'_{x,y}/x \cong F$ . ■

**Definition 2.4.** Let  $M$  be a graphic matroid in which no two elements are in series and let  $F \in \mathcal{F}$ . We say that  $M$  is minimal with respect to  $F$  if there exist two elements  $x$  and  $y$  of  $M$  such that  $M'_{x,y} \setminus \{a\}/\{x\} \cong F$  or  $M'_{x,y} \setminus \{a\}/\{x,y\} \cong F$  or  $M'_{x,y} \cong F$  or  $M'_{x,y}/\{x\} \cong F$  or  $M'_{x,y}/\{x,y\} \cong F$ .

**Corollary 2.5.** Let  $M$  be a graphic matroid. For any  $x,y \in E(M)$ , the matroid  $M'_{x,y}$  is graphic if and only if  $M$  has no minor isomorphic to a minimal matroid with respect to any  $F \in \mathcal{F}$ .

**Proof.** If  $M'_{x,y}$  is not graphic for some  $x,y$ , then by Lemma 2.3,  $M$  has a minor  $N$  in which no two elements are in series and  $N'_{x,y} \setminus \{a\}/\{x\} \cong F$  or  $N'_{x,y} \setminus \{a\}/\{x,y\} \cong F$  or  $N'_{x,y} \cong F$  or  $N'_{x,y}/\{x\} \cong F$  or  $N'_{x,y}/\{y\} \cong F$  or  $N'_{x,y}/\{x,y\} \cong F$  for some  $F \in \mathcal{F}$ . If  $N'_{x,y}/y \cong F$  but  $N'_{x,y}/x \not\cong F$ , then interchange roles of  $x$  and  $y$ . Conversely, suppose that  $M$  has a minor  $N$  isomorphic to a minimal matroid with respect to some  $F \in \mathcal{F}$ . Then  $N'_{x,y} \setminus \{a\}$  or  $N'_{x,y}/\{x\}$  or  $N'_{x,y}/\{x,y\}$  or  $N'_{x,y} \cong F$ , for some  $x,y \in E(M)$ . Then  $M'_{x,y}$  has a minor isomorphic to  $F$  and hence it is not graphic. ■

**Lemma 2.6.** Let  $M$  be a graphic matroid corresponding to a graph  $G$ . If  $M$  is minimal with respect to some  $F \in \mathcal{F}$ , then

- (i)  $M$  has neither loops nor coloops;
- (ii)  $x$  and  $y$  are non-adjacent edges of  $G$  and the minimum degree of  $G$  is at least 3;
- (iii)  $x$  and  $y$  cannot be parallel in  $G$ ;
- (iv) every pair of parallel edges of  $G$  must contain either  $x$  or  $y$ ;
- (v) if  $M'_{x,y}$  or  $M'_{x,y}/\{x\} \cong F_7^*$  or  $M^*(K_5)$ , then  $G$  is simple;
- (vi) if  $M'_{x,y}/\{x\} \cong F_7$  or  $M^*(K_{3,3})$ , then  $G$  is simple or has exactly one pair of parallel edges and one of these two edges must be  $y$ , and further there is no 3-circuit in  $G$  containing both  $x$  and  $y$ ;
- (vii) if  $M'_{x,y}/\{x,y\} \cong F$  then  $G$  is simple and there is no 3-circuit or 4-circuit in  $G$  containing both  $x$  and  $y$ .

**Proof.** (i) On the contrary, suppose  $M$  has a loop, say  $z$ . If  $z$  is different from  $x$  and  $y$ , then it is a loop in  $M'_{x,y}$  and hence in  $F$ , a contradiction. If  $z$  is one of the two elements  $x$  and  $y$ , say  $x$ , then  $M'_{x,y} \setminus \{a\}/\{x\} \cong M \setminus \{x\}$  and  $M'_{x,y} \setminus \{a\}/\{x,y\} \cong M \setminus \{x\}/\{y\}$ . This implies that  $F$  is a minor of  $M$ , a contradiction. Also,  $M'_{x,y}$  contains a 2-circuit, so it cannot be isomorphic to  $F$ . Further  $M'_{x,y}/\{x\}$  and  $M'_{x,y}/\{x,y\}$  contains a loop, a contradiction. Thus,  $M$  cannot have loops.

Suppose that  $M$  has a coloop, say  $w$ . If  $w$  is different from  $x$  and  $y$  then it is preserved in  $M'_{x,y}$  and hence in  $F$ , a contradiction. If  $w$  is one of the two elements  $x$  and  $y$ , say  $x$ , then  $\{y, a\}$  is a 2-cocircuit or  $\{y\}$  is a coloop of  $M'_{x,y}$ . Now, in  $M'_{x,y} \setminus \{a\}/\{x\}$ ,  $y$  becomes a coloop, a contradiction. Also,  $M'_{x,y} \setminus \{a\}/\{x,y\} \cong M/\{x\} \setminus \{y\}$ . This means that  $F$  is a minor of  $M$ , a contradiction. In  $M'_{x,y}$ ,  $\{x\}$  remains a coloop and hence  $M'_{x,y}$  cannot be isomorphic to  $F$ . Moreover, in  $M'_{x,y}/\{x\}$ ,  $\{y, a\}$  remains a 2-cocircuit or  $\{y\}$  remains a cocircuit and in  $F$ , a contradiction. Also,  $M'_{x,y}/\{x,y\} \cong M'_{x,y} \setminus \{x\}/\{y\} \cong M \setminus \{x\}$ , that is  $F$  is a minor of  $M$ , a contradiction. Hence  $M$  cannot have coloops.

(ii) Follows from Lemma 2.1(viii) and Lemma 2.3.

(iii) If  $x$  and  $y$  are parallel in  $G$ , then  $x$  and  $y$  remain parallel in  $M'_{x,y}$ . So, we get a loop in  $M'_{x,y} \setminus \{a\}/\{x\}$ ,  $M'_{x,y}/\{x\}$  and a 2-circuit in  $M'_{x,y}$ , a contradiction. Also,  $M'_{x,y} \setminus \{a\}/\{x,y\} = M_{x,y}/\{x,y\} = M \setminus \{x,y\}$ , a contradiction. Now,  $M'_{x,y}/\{x,y\} = M'_{x,y}/y \setminus x \cong M \setminus x$ , a contradiction to Lemma 2.1(vii). Hence these matroids are not isomorphic to  $F$ , a contradiction.

(iv) Suppose that the edges  $x_1$  and  $x_2$  are in a parallel class of  $G$  that does not contain  $x$  or  $y$ , then  $x_1$  and  $x_2$  remain in parallel in each of the ma-

troids  $M'_{x,y} \setminus \{a\}/\{x\}$ ,  $M'_{x,y} \setminus \{a\}/\{x,y\}$ ,  $M'_{x,y}$ ,  $M'_{x,y}/\{x\}$  and  $M'_{x,y}/\{x,y\}$ , a contradiction. If  $x_1$  and  $x_2$  are in a parallel class containing  $x$  or  $y$ , then we get a loop in  $M'_{x,y} \setminus \{a\}/\{x\}$ ,  $M'_{x,y} \setminus \{a\}/\{x,y\}$ ,  $M'_{x,y}/\{x\}$ ,  $M'_{x,y}/\{x,y\}$  and a 2-circuit in  $M'_{x,y}$ . Hence these matroids are not isomorphic to  $F$ , a contradiction.

(v) As  $F_7^*$  and  $M^*(K_5)$  are bipartite, if  $G$  contains a pair of parallel edges then by (iv) above, it must contain  $x$  or  $y$ . So, we get a 3-circuit in  $M'_{x,y}$  containing  $a$ . Therefore  $M'_{x,y} \not\cong F_7^*$  or  $M^*(K_5)$ . Also, we get a 2-circuit in  $M'_{x,y}/\{x\}$  and  $M'_{x,y}/\{x,y\}$ , a contradiction.

(vi) Suppose that  $G$  is not simple. Then by (iv) above, each pair of parallel edges must contain  $x$  or  $y$ . If  $\{x, x_1\}$  is a 2-circuit for some edge  $x_1$  of  $G$ , then  $\{x, x_1, a\}$  is a 3-circuit in  $M'_{x,y}$  and hence,  $\{x_1, a\}$  is a 2-circuit in  $M'_{x,y}/\{x\}$ , a contradiction. Hence  $G$  has exactly one pair of parallel edges and one of these two edges must be  $y$ .

(vii) If  $G$  contains a pair of parallel edges, it must contain  $x$  or  $y$ , say  $x$ . Then  $M'_{x,y}$  contains a 3-circuit containing  $x$  and  $a$ . Consequently,  $M'_{x,y}/\{x,y\}$  contains a 2-circuit and hence it is in  $F$ , a contradiction. Now, if  $G$  contains 3 or a 4-circuit containing both  $x$  and  $y$ , then  $M'_{x,y}/\{x,y\}$  contains a loop or 2-circuit respectively and hence it is in  $F$ , a contradiction. ■

### 3. THE ELEMENT SPLITTING OPERATION ON GRAPHIC MATROIDS

In this section, we obtain the minimal matroids corresponding to each of the four matroids  $F_7, F_7^*, M^*(K_{3,3})$  and  $M^*(K_5)$  and use them to give a proof of Theorem 1.2.

In the following lemma, we characterize minimal matroids corresponding to the matroid  $F_7$ .




Figure 4



**Lemma 3.1.** *Let  $M$  be a graphic matroid. Then  $M$  is minimal with respect to the matroid  $F_7$  if and only if  $M$  is isomorphic to one of the three circuit matroids  $M(G_1)$ ,  $M(G_2)$  and  $M(G_3)$ , where  $G_1$ ,  $G_2$  and  $G_3$  are the graphs of Figure 4.*

**Proof.** Firstly, we consider the graph  $G_3$  and prove that  $M'(G_3)_{x,y}/\{x\} \cong F_7$ .

Let matrices  $A$  and  $A'_{x,y}$  represent the matroids  $M(G_3)$  and  $M'(G_3)_{x,y}$  respectively. Then

$$A = \begin{array}{cccccccc} & 1 & 2 & 3 & 4 & 5 & x & y \\ \begin{bmatrix} 1 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} & & & & & & & \end{array}.$$

So, we have

$$A'_{x,y} = \begin{array}{ccccccccccc} & 1 & 2 & 3 & 4 & 5 & x & y & a \\ \begin{bmatrix} 1 & 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \end{bmatrix} & & & & & & & & & \end{array}.$$

Therefore

$$A'_{x,y}/\{x\} = \begin{array}{cccccccc} & 1 & 2 & 3 & 4 & 5 & y & a \\ \begin{bmatrix} 1 & 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 1 \end{bmatrix} & & & & & & & \end{array}.$$

Hence  $M'(G_3)_{x,y}/\{x\} \cong F_7$ .

One can check similarly that  $M'(G_1)_{x,y} \setminus \{a\}/\{x\} \cong F_7$ ;  $M'(G_2)_{x,y} \setminus \{a\}/\{x, y\} \cong F_7$ . Thus the matroids  $M(G_1)$ ,  $M(G_2)$  and  $M(G_3)$  are minimal with respect to  $F_7$ .

Conversely, let  $M$  be a minimal matroid with respect to  $F_7$ . Let  $G$  be a graph corresponding to  $M$ . Let the edges  $x$  and  $y$  of  $G$  are such that  $M'(G)_{x,y} \setminus \{a\}/\{x\} \cong F_7$  or  $M'(G)_{x,y} \setminus \{a\}/\{x, y\} \cong F_7$  or  $M'(G)_{x,y} \cong F_7$  or  $M'(G)_{x,y}/\{x\} \cong F_7$  or  $M'(G)_{x,y}/\{x, y\} \cong F_7$ .

By Lemma 2.1(i),  $M(G)_{x,y}/\{x\} \cong F_7$ . If  $M'(G)_{x,y} \setminus \{a\}/\{x\} \cong F_7$ , then by Lemma 3.1 of [11],  $G$  is isomorphic to the graph  $G_1$  of Figure 4. Similarly, if  $M'(G)_{x,y} \setminus \{a\}/\{x, y\} \cong F_7$ , then by Lemma 3.1 of [11],  $G$  is isomorphic to the graph  $G_2$  of Figure 4. Further,  $M'(G)_{x,y} \not\cong F_7$  because  $M'_{x,y}$  is not eulerian by Lemma 2.1(x).

Suppose that  $M'(G)_{x,y}/\{x\} \cong F_7$ . Since  $r(F_7) = 3$ ,  $r(M'(G)_{x,y}) = 4$ . Further  $|E(M'(G)_{x,y})| = 8$ . Consequently,  $r(M(G)) = 3$  and  $|E(M(G))| = 7$ . Thus,  $G$  is a graph with 4 vertices and 7 edges. This implies that  $G$  is non-simple. Also, by Lemma 2.6(vi),  $G$  has exactly one pair of parallel edges. Hence  $G$  can be obtained from a simple graph with 4 vertices and 6 edges by adding an edge in parallel. Since the complete graph  $K_4$  is the only simple graph with 4 vertices and 6 edges (see [5]),  $G$  must be isomorphic to the graph  $G_3$  of Figure 4.

Suppose that  $M'(G)_{x,y}/\{x,y\} \cong F_7$ . Then  $r(M'(G)_{x,y}) = 5$  and  $|E(M'(G)_{x,y})| = 9$ . This implies that  $r(M(G)) = 4$  and  $|E(M(G))| = 8$ . Thus,  $G$  is a graph with 5 vertices and 8 edges. Hence, by Lemma 2.6(ii),  $G$  has degree sequence  $(4,3,3,3,3)$ . By Lemma 2.6(vii),  $G$  is simple and does not have a 3-circuit or a 4-circuit containing both  $x$  and  $y$ . There is only one simple graph with 5 vertices and 8 edges (see [5]) as shown in Figure 5. In this graph, any two edges are either in a 3-circuit or in a 4-circuit. Hence we discard this graph. ■




Figure 5

We characterize minimal matroids corresponding to the matroid  $F_7^*$  in the following lemma.

**Lemma 3.2.** *Let  $M$  be a graphic matroid. Then  $M$  is minimal with respect to the matroid  $F_7^*$  if and only if  $M$  is isomorphic to one of the two circuit matroids  $M(G_4)$  and  $M(G_5)$ , where  $G_4$  and  $G_5$  are the graphs of Figure 6.*




Figure 6

**Proof.** Observe that  $M'(G_4)_{x,y} \setminus \{a\}/\{x\} \cong F_7^*$  and  $M'(G_5)_{x,y} \cong F_7^*$ . Therefore  $M(G_4)$  and  $M(G_5)$  are minimal with respect to  $F_7^*$ .

Conversely, let  $M(G)$  be a minimal graph with respect to  $F_7^*$  and let  $x$  and  $y$  be edges of  $G$  such that  $M'(G)_{x,y} \setminus \{a\}/\{x\} \cong F_7^*$  or  $M'(G)_{x,y} \setminus \{a\}/\{x,y\} \cong F_7^*$  or  $M'(G)_{x,y} \cong F_7^*$  or  $M'(G)_{x,y}/\{x\} \cong F_7^*$  or  $M'(G)_{x,y}/\{x,y\} \cong F_7^*$ . If  $M'(G)_{x,y} \setminus \{a\}/\{x\} \cong F_7^*$ , then by Lemma 2.1(i),  $M(G)_{x,y}/\{x\} \cong F_7^*$ . Hence, by Lemma 3.2 of [11],  $G$  is isomorphic to the graph  $G_4$  of Figure 6. Similarly, if  $M'(G)_{x,y} \setminus \{a\}/\{x,y\} \cong F_7^*$ , then  $M(G)_{x,y}/\{x,y\} \cong F_7^*$ . By Lemma 3.2 of [11], there is no minimal graphic matroid such that  $M(G)_{x,y}/\{x,y\} \cong F_7^*$ . In each of the remaining three cases,  $G$  is simple by Lemma 2.6(v).

Suppose that  $M'(G)_{x,y} \cong F_7^*$ . Then  $r(M(G)) = r(M'(G)_{x,y}) - 1 = 3$ . Further,  $|E(M)| = 7$ . Since  $r(F_7^*) = 4$ ,  $r(M(G)) = 3$ . Consequently,  $G$  is a simple graph with 4 vertices and 6 edges. Hence  $G$  is isomorphic to  $K_4$ , which is the graph  $G_5$  of Figure 6. Suppose that  $M'(G)_{x,y}/\{x\} \cong F_7^*$ . Then  $r(M(G)) = 4$  and  $|E(M(G))| = 7$ . Hence  $G$  is a graph with 5 vertices and 7 edges and has a vertex of degree less than 3, a contradiction to Lemma 2.6(ii). Finally, if  $M'(G)_{x,y}/\{x,y\} \cong F_7^*$ , then  $G$  has 6 vertices and 8 edges and hence a vertex of degree less than 3, a contradiction. ■

The minimal matroids corresponding to the matroid  $M^*(K_{3,3})$  are characterized as follows.

**Lemma 3.3.** *Let  $M$  be a graphic matroid. Then  $M$  is minimal with respect to the matroid  $M^*(K_{3,3})$  if and only if  $M$  is isomorphic to one of the five circuit matroids  $M(G_6), M(G_7), M(G_8), M(G_9)$  and  $M(G_{10})$ , where  $G_6, G_7, G_8, G_9$  and  $G_{10}$  are the graphs of Figure 7.*




Figure 7

**Proof.** Observe that  $M'(G_6)_{x,y} \setminus \{a\}/\{x\} \cong M^*(K_{3,3})$ ;  $M'(G_7)_{x,y} \setminus \{a\}/\{x\} \cong M^*(K_{3,3})$ ;  $M'(G_8)_{x,y} \setminus \{a\}/\{x,y\} \cong M^*(K_{3,3})$ ;  $M'(G_9)_{x,y} \setminus \{a\}/\{x,y\} \cong$

$M^*(K_{3,3})$  and  $M'(G_{10})_{x,y}/\{x\} \cong M^*(K_{3,3})$ . This implies that  $M(G_6)$ ,  $M(G_7)$ ,  $M(G_8)$ ,  $M(G_9)$  and  $M(G_{10})$  are minimal matroids with respect to the matroid  $M^*(K_{3,3})$ .

Conversely, let  $M(G)$  be a minimal matroid with respect to  $M^*(K_{3,3})$ . Let  $x$  and  $y$  be edges of  $G$  such that  $M'(G)_{x,y} \setminus \{a\}/\{x\} \cong M^*(K_{3,3})$  or  $M'(G)_{x,y} \setminus \{a\}/\{x,y\} \cong M^*(K_{3,3})$  or  $M'(G)_{x,y} \cong M^*(K_{3,3})$  or  $M'(G)_{x,y}/\{x\} \cong M^*(K_{3,3})$  or  $M'(G)_{x,y}/\{x,y\} \cong M^*(K_{3,3})$ . If  $M'(G)_{x,y} \setminus \{a\}/\{x\} \cong M^*(K_{3,3})$ , then by Lemma 2.1(i),  $M(G)_{x,y}/\{x\} \cong M^*(K_{3,3})$ . Hence, by Lemma 3.3 of [11],  $G$  is isomorphic to one of the two graphs  $G_6$  and  $G_7$  of Figure 7. If  $M'(G)_{x,y} \setminus \{a\}/\{x,y\} \cong M^*(K_{3,3})$ , then by Lemma 2.1(i),  $M(G)_{x,y}/\{x,y\} \cong M^*(K_{3,3})$ . Hence by Lemma 3.3 of [11],  $G$  is isomorphic to one of the two graphs  $G_8$  and  $G_9$  of Figure 7.  $M'_{x,y}$  is not eulerian, by Lemma 2.1(x). Therefore  $M'(G)_{x,y} \not\cong M^*(K_{3,3})$ .




Figure 8

Suppose that  $M'(G)_{x,y}/\{x\} \cong M^*(K_{3,3})$ . Then  $r(M(G)) = 4$  and  $|E(M(G))| = 9$ . Consequently,  $G$  is a graph with 5 vertices and 9 edges. Suppose that  $G$  is simple. By [5], any simple graph with 5 vertices and 9 edges is isomorphic to the graph of Figure 8. Suppose  $G$  is isomorphic to this graph. Then  $G$  has two edge-disjoint 3-cocircuits. Out of which, by Lemma 2.1(iv), at least one 3-cocircuit is preserved in  $M'(G)_{x,y}/\{x\}$  and hence it is preserved in  $M^*(K_{3,3})$ , a contradiction. Thus  $G$  is non-simple. By Lemma 2.6(vi),  $G$  has exactly one pair of parallel edges. Since the degree of a vertex in  $G$  is at least 3, the degree sequence of  $G$  is  $(6,3,3,3,3)$ ,  $(5,4,3,3,3)$  or  $(4,4,4,3,3)$ . Therefore  $G$  can be obtained from a simple graph with 5 vertices and 8 edges by adding an edge in parallel. There are in all 2 non-isomorphic simple graphs with 5 vertices and 8 edges (see [5]). So, there are in all 3 possibilities for  $G$  as shown in Figure 9.

If  $G$  is isomorphic to one of the two graphs (i) and (ii) of Figure 9, then it has two edge-disjoint 3-cocircuits, and hence at least one of them is survived in  $M'(G)_{x,y}/\{x\} \cong M^*(K_{3,3})$ , a contradiction. Hence  $G$  is isomorphic to third graph which is nothing but the graph  $G_{10}$  of Figure 7.




Figure 9

Finally, suppose that  $M'(G)_{x,y}/\{x,y\} \cong M^*(K_{3,3})$ . Since  $r(M^*(K_{3,3})) = 4$ ,  $r(M'(G)_{x,y}) = 6$ . This shows that  $r(M(G)) = 5$  and  $|E(M(G))| = 10$ . Consequently,  $G$  is a graph with 6 vertices and 10 edges with minimum degree at least 3. So, the degree sequence of  $G$  is  $(4,4,3,3,3,3)$  or  $(5,3,3,3,3,3)$ . By Lemma 2.6(vii),  $G$  is simple. There are in all 4 non-isomorphic simple graphs with 6 vertices and 10 edges having the said degree sequences as shown in Figure 10 (see [5]). By Lemma 2.6(vii),  $G$  does not have a 3-circuit or a 4-circuit containing both  $x$  and  $y$ . As there are no 3-cocircuits and 5-cocircuits in  $M^*(K_{3,3})$ , every such cocircuit in  $G$  contains  $x$  or  $y$ . Suppose  $G$  is the graph (i) or graph (ii) of Figure 10. Then there is only one choice for  $x, y$ , as shown in the figure. For these choices  $M'(G)_{x,y}/\{x,y\}$  is not Eulerian, a contradiction. If  $G$  is the graph (iii) or graph (iv) of Figure 10, then we get a 3-cocircuit or a 5-cocircuit in  $M'(G)_{x,y}/\{x,y\}$  and hence it is in  $M^*(K_{3,3})$ , a contradiction. ■




Figure 10

Finally, we characterize minimal matroids corresponding to the matroid  $M^*(K_5)$  in the following lemma.

**Lemma 3.4.** *Let  $M$  be a graphic matroid. Then  $M$  is minimal with respect to the matroid  $M^*(K_5)$  if and only if  $M$  is isomorphic to one of the three circuit matroids  $M(G_{11})$ ,  $M(G_{12})$  and  $M(G_{13})$ , where  $G_{11}$ ,  $G_{12}$  and  $G_{13}$  are the graphs of Figure 11.*




Figure 11

**Proof.** Observe that  $M'(G_{11})_{x,y} \setminus \{a\}/\{x\} \cong M^*(K_5)$ ;  $M'(G_{12})_{x,y} \setminus \{a\}/\{x\} \cong M^*(K_5)$  and  $M'(G_{13})_{x,y} \cong M^*(K_5)$ . Therefore  $M(G_{11})$ ,  $M(G_{12})$  and  $M(G_{13})$  are minimal matroids with respect to the matroid  $M^*(K_5)$ .

Conversely, let  $M(G)$  be a minimal matroid with respect to  $M^*(K_5)$  and let  $x$  and  $y$  be edges of  $G$  such that  $M'(G)_{x,y} \setminus \{a\}/\{x\} \cong M^*(K_5)$  or  $M'(G)_{x,y} \setminus \{a\}/\{x,y\} \cong M^*(K_5)$  or  $M'(G)_{x,y} \cong M^*(K_5)$  or  $M'(G)_{x,y}/\{x\} \cong M^*(K_5)$  or  $M'(G)_{x,y}/\{x,y\} \cong M^*(K_5)$ . If  $M'(G)_{x,y} \setminus \{a\}/\{x\} \cong M^*(K_5)$ , then by Lemma 2.1(i),  $M(G)_{x,y}/\{x\} \cong M^*(K_5)$ . Therefore, by Lemma 3.4 of [11],  $G$  is isomorphic to one of the two graphs  $G_{11}$  and  $G_{12}$  of Figure 11. If  $M'(G)_{x,y} \setminus \{a\}/\{x,y\} \cong M^*(K_5)$ , then  $M(G)_{x,y}/\{x,y\} \cong M^*(K_5)$ . By Lemma 3.4 of [11], there is no minimal graphic matroid  $M(G)$  such that  $M(G)_{x,y}/\{x,y\} \cong M^*(K_5)$ . By Lemma 2.6(v),  $G$  is simple in the remaining three cases. Suppose that  $M'(G)_{x,y} \cong M^*(K_5)$ . Then  $r(M(G)) = 5$  and  $|E(M(G))| = 9$ . Hence,  $G$  is a graph with 6 vertices and 9 edges having degree sequence  $(3,3,3,3,3,3)$ . There are only two such non-isomorphic simple graphs, (see [5]) as shown in Figure 12. In graph (i) of Figure 12,




Figure 12

for every choice of non-adjacent edges  $x, y$ , there is a 4-circuit containing exactly one of  $x$  and  $y$ . Such circuit becomes a 5-circuit in  $M'_{x,y}$ , a contradiction. Hence, the circuit matroid of this graph is not minimal with respect

to  $M^*(K_5)$ . Hence  $G$  is isomorphic to graph (ii) of Figure 12 which is in fact the graph  $G_{13}$  of Figure 11.

Suppose that  $M'(G)_{x,y}/\{x\} \cong M^*(K_5)$ . Then  $G$  has 7 vertices and 10 edges. Hence  $G$  has a vertex of degree less than 3, a contradiction. Suppose that  $M'(G)_{x,y}/\{x, y\} \cong M^*(K_5)$ . Then  $G$  is a graph with 8 vertices and 11 edges. Hence  $G$  has a vertex of degree less than 3, a contradiction. ■

Now, we prove Theorem 1.2.

**Proof of Theorem 1.2.** Let  $M$  be a graphic matroid and let  $G$  be a graph such that  $M = M(G)$ . On combining Corollary 2.5 and Lemmas 3.1, 3.2, 3.3 and 3.4, it follows that  $M'(G)_{x,y}$  is graphic for every pair  $\{x, y\}$  of edges of  $G$  if and only if  $M(G)$  has no minor isomorphic to any of the matroids  $M(G_i)$ ,  $i = 1, 2, \dots, 13$ , where the graphs  $G_i$  are as shown in Figures 4, 6, 7 and 11. However, we have  $M(G_5) \cong M(G_1) \setminus \{e_2, e_3\} \cong M(G_2)/\{z\} \setminus \{e_1, e_2\} \cong M(G_3) \setminus \{5\} \cong M(G_4)/\{w\} \setminus \{e_1\} \cong M(G_6)/\{e_6\} \setminus \{w, e_1, e_2\} \cong M(G_7)/\{z\} \setminus \{y, e_4, e_5\} \cong M(G_8)/\{z, e_2\} \setminus \{e_1, e_3, e_5\} \cong M(G_9)/\{x, e_1\} \setminus \{w, e_4, e_5\} \cong M(G_{10})/\{3\} \setminus \{5, 7\} \cong M(G_{11})/\{x, e_4, e_5\} \setminus \{w, e_1\} \cong M(G_{12})/\{v, z, e_3\} \setminus \{x, e_1\} \cong M(G_{13})/\{1, 5\} \setminus \{x\}$ . Thus,  $M'(G)_{x,y}$  is graphic if and only if  $M(G)$  has no minor isomorphic to the matroid  $M(G_5)$ . Observe that the graph  $G_5$  is isomorphic to the complete graph  $K_4$ . This completes the proof of the theorem. ■

#### REFERENCES

- [1] G. Azadi, Generalized splitting operation for binary matroids and related results (Ph.D. Thesis, University of Pune, 2001).
- [2] Y.M. Borse, M.M. Shikare and Kiran Dalvi, *Excluded-Minor characterization for the class of Cographic Splitting Matroids*, Ars Combin., to appear.
- [3] H. Fleischner, Eulerian Graphs and Related Topics, Part 1, Vol. 1 (North Holland, Amsterdam, 1990).
- [4] A. Habib, Some new operations on matroids and related results (Ph.D. Thesis, University of Pune, 2005).
- [5] F. Harary, Graph Theory (Addison-Wesley, Reading, 1969).
- [6] J.G. Oxley, Matroid Theory (Oxford University Press, Oxford, 1992).
- [7] T.T. Raghunathan, M.M. Shikare and B.N. Waphare, *Splitting in a binary matroid*, Discrete Math. **184** (1998) 267–271.
- [8] A. Recski, Matroid Theory and Its Applications (Springer Verlag, Berlin, 1989).

- [9] M.M. Shikare and G. Azadi, *Determination of the bases of a splitting matroid*, European J. Combin. **24** (2003) 45–52.
- [10] M.M. Shikare, *Splitting lemma for binary matroids*, Southeast Asian Bull. Math. **32** (2007) 151–159.
- [11] M.M. Shikare and B.N. Waphare, *Excluded-Minors for the class of graphic splitting matroids*, Ars Combin., to appear.
- [12] P.J. Slater, *A classification of 4-connected graphs*, J. Combin. Theory **17** (1974) 281–298.
- [13] W.T. Tutte, *A theory of 3-connected graphs*, Indag. Math. **23** (1961) 441–455.
- [14] D.J.A. Welsh, *Matroid Theory* (Academic Press, London, 1976).

Received 15 October 2008  
Revised 17 December 2008  
Accepted 17 December 2008