

ON CHARACTERIZATION OF UNIQUELY 3-LIST
COLORABLE COMPLETE MULTIPARTITE GRAPHS*

YANCAI ZHAO^{1,2} AND ERFANG SHAN¹

¹*Department of Mathematics*
Shanghai University
Shanghai 200444, P.R. China

²*Department of Science*
Bengbu University
Anhui 233030, P.R. China

e-mail: zhaoyc69@126.com

Abstract

For each vertex v of a graph G , if there exists a list of k colors, $L(v)$, such that there is a unique proper coloring for G from this collection of lists, then G is called a *uniquely k -list colorable graph*. Ghebleh and Mahmoodian characterized uniquely 3-list colorable complete multipartite graphs except for nine graphs: $K_{2,2,r}$ $r \in \{4, 5, 6, 7, 8\}$, $K_{2,3,4}$, $K_{1*4,4}$, $K_{1*4,5}$, $K_{1*5,4}$. Also, they conjectured that the nine graphs are not U3LC graphs. After that, except for $K_{2,2,r}$ $r \in \{4, 5, 6, 7, 8\}$, the others have been proved not to be U3LC graphs. In this paper we first prove that $K_{2,2,8}$ is not U3LC graph, and thus as a direct corollary, $K_{2,2,r}$ ($r = 4, 5, 6, 7, 8$) are not U3LC graphs, and then the uniquely 3-list colorable complete multipartite graphs are characterized completely.

Keywords: list coloring, complete multipartite graph, uniquely 3-list colorable graph.

2010 Mathematics Subject Classification: 05C15.

*Research was partially supported by the National Nature Science Foundation of China (No. 60773078) and Key Disciplines of Shanghai Municipality (S30104).

1. INTRODUCTION

We consider simple graphs which are finite, undirected, with no loops or multiple edges. For the necessary definitions and notations we refer the reader to standard texts, such as [2].

By a k -list assignment L to a graph G we mean a map which assigns to each vertex v of G a set $L(v)$ of size k . A *list coloring* for G from L , or an L -coloring for short, is a proper coloring c , in which for each vertex v , $c(v)$ is chosen from $L(v)$. The idea of list coloring is due independently to Vizing [11] and to Erdős, Rubin, and Taylor [4]. For a recent survey on list coloring we refer the interested reader to Alon [1].

For each vertex v in G , if there exists a list of k colors $L(v)$, such that there exists a uniquely L -coloring for G , then G is called a *uniquely k -list colorable graph* or a $UkLC$ graph for short. The idea of uniquely colorable graph was introduced independently by Dinitz and Martin [3] and by Mahmoodian and Mahdian [8].

If a graph G is not uniquely k -list colorable, we also say that G has property $M(k)$. So G has the property $M(k)$ if and only if for any collection of lists assigned to its vertices, each of size k , either there is no list coloring for G or there exist at least two list colorings. The least integer k such that G has the property $M(k)$ is called the *m -number* of G , denoted by $m(G)$. This conception was originally introduced by Mahmoodian and Mahdian in [9].

Mahdian and Mahmoodian characterized uniquely 2-list colorable graphs as follows:

Theorem 1.1 ([8]). *A connected graph G has the property $M(2)$ if and only if every block of G is either a cycle, a complete graph, or a complete bipartite graph.*

Ghebleh and Mahmoodian studied uniquely 3-list colorability about complete multipartite graphs, and they gave the following results:

Theorem 1.2 ([6]). *Graphs $K_{3,3,3}$, $K_{2,4,4}$, $K_{2,3,5}$, $K_{2,2,9}$, $K_{1,2,2,2}$, $K_{1,1,2,3}$, $K_{1,1,1,2,2}$, $K_{1*4,6}$, $K_{1*5,5}$, $K_{1*6,4}$ are uniquely 3-list colorable graphs.*

Here, $K_{s*r,t}$ denote a complete $(r+1)$ -partite graph in which each part of the r parties is of size s , one party is of size t .

Theorem 1.3 ([6]). *Let G be a complete multipartite graph that is not $K_{2,2,r}$, $r \in \{4, 5, 6, 7, 8\}$, $K_{2,3,4}$, $K_{1*4,4}$, $K_{1*4,5}$, $K_{1*5,4}$, then G is U3LC if and only if it has one of the ten graphs in Theorem 1.2 as an induced subgraph.*

It can be seen from Theorem 1.2 and Theorem 1.3 that, if we can determine whether the nine graphs exempted in Theorem 1.3 are U3LC or not, we can simplify Theorem 1.3. About the nine graphs, Ref. [6] gave an open problem.

Problem 1 ([6]). Verify the property $M(3)$ for the graphs $K_{2,2,r}$, $r \in \{4, 5, 6, 7, 8\}$, $K_{2,3,4}$, $K_{1*4,4}$, $K_{1*4,5}$, and $K_{1*5,4}$.

Recently, except for $K_{2,2,r}$, $r \in \{4, 5, 6, 7, 8\}$, the other graphs in the above problem have been proved to have the property $M(3)$. They are shown below:

Theorem 1.4 ([10]). *The graph $K_{1*5,4}$ has property $M(3)$.*

Theorem 1.5 ([7]). *Graphs $K_{1*4,5}$ and $K_{1*4,4}$ have property $M(3)$.*

Theorem 1.6 ([12]). *The graph $K_{2,3,4}$ has property $M(3)$.*

In the next section of this paper, we will prove that $K_{2,2,8}$ has the property $M(3)$. As its a direct corollary, we show that $K_{2,2,r}$, $r \in \{4, 5, 6, 7\}$ have the property $M(3)$, and thus based on the results we prove together with previous results, we completely characterize the U3LC complete multipartite graphs.

2. $K_{2,2,r}$, $r \in \{4, 5, 6, 7, 8\}$ HAVE PROPERTY $M(3)$

First we give three results as lemmas.

Lemma 2.1 ([5]). *If G is a U3LC complete tripartite graph, then in its uniquely 3-list coloring, at least two colors are used to color all vertices in each part of G .*

Lemma 2.2 ([9]). *If L is a k -list assignment to the vertices in a graph G , and G has a unique L -coloring, then $|\bigcup_v L(v)| \geq k + 1$ and all these colors are used in the unique L -coloring of G .*

Lemma 2.3 ([6]). *If G is a complete multipartite graph which has an induced UkLC subgraph, then G is UkLC .*

In the graph $K_{2,2,8}$, we denote its three parts by $X_1 = \{v_1, v_2\}$, $X_2 = \{v_3, v_4\}$, $X_3 = \{v_5, v_6, \dots, v_{12}\}$, and denote 3-list assigned to the vertex v_i of $K_{2,2,8}$ by $L(v_i) = \{c_{i1}, c_{i2}, c_{i3}\}$, $1 \leq i \leq 12$. Now, we suppose that $K_{2,2,8}$ has a unique L -coloring c such that $c(v_i) = c_{i1}$, $i \in \{1, 2, \dots, 12\}$. For convenience, let $S = \{c_{51}, c_{61}, \dots, c_{12,1}\}$. To obtain our main result, we give more lemmas as follows.

Lemma 2.4. *Colors $c_{11}, c_{21}, c_{31}, c_{41}$ are different from each other.*

Proof. The result follows from Lemma 2.1. ■

Lemma 2.5. *If $v_i \neq v_j$ are in the same part of $K_{2,2,8}$, then $c_{i1} \notin \{c_{j2}, c_{j3}\}$.*

Proof. Otherwise, if $c_{i1} = c_{jk}$, $k \in \{2, 3\}$, then by letting $c'(v_j) = c_{jk} = c_{i1}$ and letting $c'(v_i) = c(v_i)$ for $i \neq j$, we obtain another L -coloring c' of $K_{2,2,8}$, which contradicts the fact that $K_{2,2,8}$ is U3LC . ■

Lemma 2.6. *If $v_i, v_j \in \{v_1, v_2, v_3, v_4\}$ are in two different parts of $K_{2,2,8}$ such that $c_{i1} \in \{c_{j2}, c_{j3}\}$, then $c_{j1} \notin \{c_{i2}, c_{i3}\}$.*

Proof. Otherwise, by letting $c'(v_i) = c_{j1}$, $c'(v_j) = c_{i1}$ and $c'(v_k) = c(v_k)$ for $k \neq i, j$, we obtain another L -coloring c' of $K_{2,2,8}$, a contradiction. ■

Lemma 2.7. *There exists an $i \in \{1, 2, 3, 4\}$ such that $\{c_{i2}, c_{i3}\} \subseteq S$.*

Proof. Otherwise, for each $i \in \{1, 2, 3, 4\}$, $|\{c_{i1}, c_{i2}, c_{i3}\} \setminus S| \geq 2$. Let $L'(v_i) = \{c_{i1}, c_{i2}, c_{i3}\} \setminus S$ for each $i \in \{1, 2, 3, 4\}$. Note that the subgraph of $K_{2,2,8}$ induced by v_1, v_2, v_3 and v_4 is $K_{2,2}$. Then by Theorem 1.1, $K_{2,2}$ has the property $M(2)$. So, there is another L -coloring c' of $K_{2,2}$ with $c'(v_i) \in L'(v_i)$, $i \in \{1, 2, 3, 4\}$, which can be easily extended to another proper coloring of $K_{2,2,8}$ by letting $c'(v_i) = c(v_i)$ for $i \in \{5, 6, \dots, 12\}$, a contradiction. ■

Lemma 2.8. *$\{c_{i2}, c_{i3}\} \subseteq \{c_{11}, c_{21}, c_{31}, c_{41}\}$ for every $i \in \{5, 6, \dots, 12\}$.*

Proof. By Lemma 2.5, if $v_i \neq v_j$ are in the same part of $K_{2,2,8}$, then $c_{i1} \notin \{c_{j2}, c_{j3}\}$. Also obviously $c_{i1} \notin \{c_{i2}, c_{i3}\}$. So $c_{i2}, c_{i3} \notin S$ for $i \in \{5, 6, \dots, 12\}$. Again by Lemma 2.2, $\{c_{i2}, c_{i3}\} \subseteq \{c_{11}, c_{21}, c_{31}, c_{41}\}$. ■

Lemma 2.9. *If X_1 and X_2 are unconsidered, then any three colors in $\{c_{11}, c_{21}, c_{31}, c_{41}\}$ can be used to L -color X_3 .*

Proof. For convenience, we can suppose $c_{i1} = i$, $i \in \{1, 2, 3, 4\}$ by Lemma 2.4. Now without loss of generality, suppose to the contrary that three colors $\{1, 2, 3\}$ cannot be used to L -color X_3 without considering X_1 and X_2 , then $\{c_{i2}, c_{i3}\} \cap \{1, 2, 3\} = \emptyset$ for some $i \in \{5, 6, \dots, 12\}$ and thus $|\{c_{i2}, c_{i3}\} \cap \{1, 2, 3, 4\}| \leq 1$, contradicting the conclusion of Lemma 2.8. ■

Lemma 2.10. *Take any three colors from $\{1, 2, 3, 4\}$, for example $\{2, 3, 4\}$. If any two colors in $\{2, 3, 4\}$ can not be used to L -color seven vertices of X_3 with X_1 and X_2 unconsidered, then there must exist two colors $\{1, 2\}$, or $\{1, 3\}$, or $\{1, 4\}$ which can be used to L -color X_3 with X_1 and X_2 unconsidered.*

Proof. If any two colors in $\{2, 3, 4\}$ can not be used to L -color seven vertices of X_3 with X_1 and X_2 unconsidered, then, noting Lemma 2.8, since $\{3, 4\}$ can not be used to L -color seven vertices of X_3 with X_1 and X_2 unconsidered, there must be at least two $i \in \{5, 6, \dots, 12\}$ such that $\{c_{i2}, c_{i3}\} = \{1, 2\}$; since $\{2, 4\}$ can not be used to L -color seven vertices of X_3 with X_1 and X_2 unconsidered, there must be at least two $i \in \{5, 6, \dots, 12\}$ such that $\{c_{i2}, c_{i3}\} = \{1, 3\}$; since $\{2, 3\}$ can not be used to L -color seven vertices of X_3 with X_1 and X_2 unconsidered, there must be at least two $i \in \{5, 6, \dots, 12\}$ such that $\{c_{i2}, c_{i3}\} = \{1, 4\}$. Without loss of generality, suppose the six sets above are $\{c_{i2}, c_{i3}\}$, $i \in \{7, 8, \dots, 12\}$. Then we check the remaining two sets $\{c_{52}, c_{53}\}, \{c_{62}, c_{63}\}$. If $\{c_{52}, c_{53}\} \cap \{c_{62}, c_{63}\} \neq \emptyset$, suppose $s \in \{c_{52}, c_{53}\} \cap \{c_{62}, c_{63}\}$. Then $1, s$ can color X_3 . If $\{c_{52}, c_{53}\} \cap \{c_{62}, c_{63}\} = \emptyset$, then $\{c_{52}, c_{53}\} \cup \{c_{62}, c_{63}\} = \{1, 2, 3, 4\}$ by Lemma 2.8. Without loss of generality, assume $\{c_{52}, c_{53}\} = \{1, 2\}$ and $\{c_{62}, c_{63}\} = \{3, 4\}$. Then $1, 3$, or $1, 4$ can be used to color X_3 . ■

Theorem 2.1. $K_{2,2,r}$, $r \in \{4, 5, 6, 7, 8\}$ have property $M(3)$.

Proof. First we prove $K_{2,2,8}$ has property $M(3)$ by contradiction. Assume to the contrary that c is a unique 3-list coloring of $K_{2,2,8}$ and $L(v_i) = \{c_{i1}, c_{i2}, c_{i3}\}$, $c(v_i) = c_{i1}$, $i \in \{1, 2, \dots, 12\}$. By Lemma 2.3, suppose $c_{i1} = i$, $i \in \{1, 2, 3, 4\}$. By Lemma 2.7, there exists an $i \in \{1, 2, 3, 4\}$ such that $\{c_{i2}, c_{i3}\} \subseteq S = \{c_{51}, c_{61}, \dots, c_{12,1}\}$. Without loss of generality, suppose such an $i = 1$ and $c_{12} = 5, c_{13} = 6$, that is, $L(v_1) = \{1, 5, 6\}$. To proceed the

proof, we distinguish the following three cases. In each case, we will obtain another L -coloring c' of $K_{2,2,8}$.

Case 1. $|\{c_{22}, c_{23}\} \cap S| = 2$, for example, $c_{22} = a, c_{23} = b, a \neq b, a, b \in S$. If $\{c_{32}, c_{33}, c_{42}, c_{43}\} \cap S \neq \emptyset$, for example, $c_{32} = d \in S$, we can let $c'(v_3) = d, c'(v_1) = m, m \in \{5, 6\} \setminus d, c'(v_2) = n, n \in \{a, b\} \setminus d, c'(v_4) = 4$. By Lemma 2.9, we can let $c'(v_i) = l_i$, where, $l_i \in \{1, 2, 3\} \cap L(v_i), i \in \{5, 6, \dots, 12\}$. Then we obtain another L -coloring c' of $K_{2,2,8}$ (which is exhibited below), a contradiction. Thus assume $\{c_{32}, c_{33}, c_{42}, c_{43}\} \cap S = \emptyset$. Then, by Lemmas 2.4–2.5, it must be that $L(v_3) = \{3, 1, 2\}, L(v_4) = \{4, 1, 2\}$. Let $c'(v_1) = 5, c'(v_2) = a, c'(v_3) = 1, c'(v_4) = 1$, and by Lemma 2.9, we can let $c'(v_i) = l_i$, where, $l_i \in \{2, 3, 4\} \cap L(v_i), i \in \{5, 6, \dots, 12\}$. Then another L -coloring c' of $K_{2,2,8}$ occurs (which is exhibited below), a contradiction.

$$\begin{array}{cc} \underline{156} & \underline{2ab} \\ 3\underline{d} \times & \underline{4} \times \times \end{array} \qquad \begin{array}{cc} \underline{156} & \underline{2ab} \\ 3\underline{12} & \underline{412} \end{array}$$

$$\{c_{32}, c_{33}, c_{42}, c_{43}\} \cap S \neq \emptyset \quad \{c_{32}, c_{33}, c_{42}, c_{43}\} \cap S = \emptyset$$

Case 2. $|\{c_{22}, c_{23}\} \cap S| = 1$, for example, $c_{22} = a \in S$. By Lemma 2.5, $1 \notin \{c_{22}, c_{23}\}$. So $3 \in \{c_{22}, c_{23}\}$ or $4 \in \{c_{22}, c_{23}\}$, for example, $3 \in \{c_{22}, c_{23}\}$. That is, $L(v_2) = \{2, a, 3\}$. We consider two subcases.

Subcase 2.1. $\{c_{32}, c_{33}\} \subseteq S$ or $\{c_{42}, c_{43}\} \subseteq S$, for example, $\{c_{32}, c_{33}\} \subseteq S$. For convenience, write $c_{32} = b, c_{33} = d, \{b, d\} \subseteq S$. Then, by letting $c'(v_2) = a, c'(v_3) = n$, where, $n \in \{b, d\} \setminus a, c'(v_1) = m, m \in \{5, 6\} \setminus n, c'(v_4) = 4$ and letting $c'(v_i) = l_i$, where, $l_i \in \{1, 2, 3\} \cap L(v_i), i \in \{5, 6, \dots, 12\}$, we obtain a new L -coloring c' of $K_{2,2,8}$ (which is exhibited below), a contradiction.

$$\begin{array}{cc} \underline{156} & \underline{2a3} \\ 3\underline{bd} & \underline{4} \times \times \end{array}$$

Subcase 2.2. $\{c_{32}, c_{33}\} \not\subseteq S$ and $\{c_{42}, c_{43}\} \not\subseteq S$. By Lemma 2.5, $4 \notin \{c_{32}, c_{33}\}$. By Lemma 2.6, $2 \notin \{c_{32}, c_{33}\}$. Noting Lemma 2.2, we have $L(v_3) = \{3, 1, b\}, b \in S$.

If $1 \in \{c_{42}, c_{43}\}$, then by letting $c'(v_3) = c'(v_4) = 1, c'(v_2) = a, c'(v_1) = 5, c'(v_i) = l_i$, where, $l_i \in \{2, 3, 4\} \cap L(v_i), i \in \{5, 6, \dots, 12\}$ we obtain another

L -coloring c' of $K_{2,2,8}$ (which is exhibited below), a contradiction. So assume $1 \notin \{c_{42}, c_{43}\}$. Then noting that $3 \notin L(v_4)$ by Lemma 2.5, there must be $\{c_{42}, c_{43}\} = \{2, d\}, d \in S$.

If $b = d$, then $5 \neq b$ or $6 \neq b$, for example $5 \neq b$. By letting $c'(v_3) = c'(v_4) = b, c'(v_1) = m, m \in \{5, 6\} \setminus b, c'(v_2) = 2, c'(v_i) = l_i$, where, $l_i \in \{1, 3, 4\} \cap L(v_i), i = 5, 6, \dots, 12$ we obtain another L -coloring c' of $K_{2,2,8}$ (which is exhibited below), a contradiction. So suppose $b \neq d$. We discuss color a in $L(v_2)$.

(1) If $a \in \{5, 6\}$, for example $a = 5$, then $5 \neq b$ or $5 \neq d$, for example $5 \neq b$. Let $c'(v_1) = c'(v_2) = 5, c'(v_3) = b, c'(v_4) = 4, c'(v_i) = l_i$, where, $l_i \in \{1, 2, 3\} \cap L(v_i), i \in \{5, 6, \dots, 12\}$, another L -coloring c' of $K_{2,2,8}$ occurring (which is exhibited below), a contradiction.

(2) If $a \notin \{5, 6\}$, then there exists at least one color in $\{5, 6, a\}$ such that it does not belong to $\{b, d\}$, for example $5 \notin \{b, d\}$. Then, by letting $c'(v_1) = 5, c'(v_2) = 2, c'(v_3) = b, c'(v_4) = d, c'(v_i) = l_i$, here, $l_i \in \{1, 3, 4\} \cap L(v_i), i \in \{5, 6, \dots, 12\}$, we obtain another L -coloring c' of $K_{2,2,8}$ (which is exhibited below), a contradiction.

$$\begin{array}{cc}
 \begin{array}{cc}
 1\underline{5}6 & 2\underline{a}3 \\
 3\underline{1}b & 4\underline{1}\times \\
 1 \in \{c_{42}, c_{43}\} \\
 1\underline{5}6 & 2\underline{5}3 \\
 3\underline{1}b & 4\underline{2}d \\
 1 \notin \{c_{42}, c_{43}\}, b \neq d \text{ (1)}
 \end{array}
 &
 \begin{array}{cc}
 1\underline{5}6 & 2a3 \\
 3\underline{1}b & 42\underline{d} \\
 1 \notin \{c_{42}, c_{43}\}, b = d \\
 1\underline{5}6 & 2a3 \\
 3\underline{1}b & 42\underline{d} \\
 1 \notin \{c_{42}, c_{43}\}, b \neq d \text{ (2)}
 \end{array}
 \end{array}$$

Case 3. $\{c_{22}, c_{23}\} \cap S = \emptyset$.

By Lemma 2.5, $1 \notin L(v_2)$. So $L(v_2) = \{2, 3, 4\}$. By Lemma 2.6, $2 \notin L(v_3), 2 \notin L(v_4)$. By Lemma 2.5, $3 \notin L(v_4), 4 \notin L(v_3)$. So there exist $a \in S$ and $b \in S$ such that $a \in L(v_3), b \in L(v_4)$, for example, $c_{32} = a, c_{42} = b$.

Subcase 3.1. If $a, b \in S \setminus \{5, 6\}$, then by letting $c'(v_1) = 5, c'(v_2) = 2, c'(v_3) = a, c'(v_4) = b, c'(v_i) = l_i$, where, $l_i \in \{1, 3, 4\} \cap L(v_i), i \in \{5, 6, \dots, 12\}$ we obtain another L -coloring c' of $K_{2,2,8}$ (which is exhibited below), a contradiction.

Subcase 3.2. If $a \in \{5, 6\}$ and $b \notin \{5, 6\}$, or $a \notin \{5, 6\}$ and $b \in \{5, 6\}$, for example $a = 5, b \in S \setminus \{5, 6\}$, then by letting $c'(v_3) = 5, c'(v_1) = 6, c'(v_4) = b, c'(v_2) = 2, c'(v_i) = l_i$, where, $l_i \in \{1, 3, 4\} \cap L(v_i), i \in \{5, 6, \dots, 12\}$ we obtain another L -coloring c' of $K_{2,2,8}$ (which is exhibited below), a contradiction.

Subcase 3.3. If $a, b \in \{5, 6\}$ and $a = b$, for example, $a = b = 5$, then by letting $c'(v_3) = c'(v_4) = 5, c'(v_1) = 6, c'(v_2) = 2, c'(v_i) = l_i$, where, $l_i \in \{1, 3, 4\} \cap L(v_i), i \in \{5, 6, \dots, 12\}$ we obtain another L -coloring c' of $K_{2,2,8}$ (which is exhibited below), a contradiction.

$\underline{156} \quad \underline{234}$	$\underline{156} \quad \underline{234}$	$\underline{156} \quad \underline{234}$
$\underline{3a} \times \quad \underline{4b} \times$	$\underline{35} \times \quad \underline{4b} \times$	$\underline{35} \times \quad \underline{45} \times$
Subcase 3.1	Subcase 3.2	Subcase 3.3

Subcase 3.4. If $a, b \in \{5, 6\}$ and $a \neq b$, for example, $a = 5, b = 6$, then we assert that $\{c_{33}, c_{43}\} \cap S = \emptyset$ holds. Otherwise, without loss of generality, suppose $c_{33} = d \in S$. Noting $d \neq 5$, we can let $c'(v_4) = 6, c'(v_3) = d, c'(v_1) = 5, c'(v_2) = 2, c'(v_i) = l_i$, where, $l_i \in \{1, 3, 4\} \cap L(v_i), i = 5, 6, \dots, 12$, another L -coloring c' of $K_{2,2,8}$ appearing, a contradiction. So $L(v_3) = \{3, 5, 1\}, L(v_4) = \{4, 6, 1\}$. Then, we consider the following two cases.

(1) There exist two colors in $\{2, 3, 4\}$ which can be used to color seven vertices of X_3 with X_1 and X_2 unconsidered. Without loss of generality, suppose $\{3, 4\}$ can be used to color $v_i, i = 5, 6, \dots, 11$, then by letting $c'(v_{12,1}) = c_{12,1}, c'(v_3) = c'(v_4) = 1, c'(v_2) = 2, c'(v_1) = m, m \in \{5, 6\} \setminus c_{12,1}, c'(v_i) = l_i$, where, $l_i \in \{3, 4\} \cap L(v_i), i \in \{5, 6, \dots, 11\}$ we obtain another L -coloring c' of $K_{2,2,8}$ (which is exhibited below), a contradiction.

(2) If any two colors in $\{2, 3, 4\}$ can not be used to color seven vertices of X_3 with the other two parts unconsidered, then by Lemma 2.10, one set of $\{1, 2\}, \{1, 3\}, \{1, 4\}$ can be used to L -color X_3 with the other two parts unconsidered. Without loss of generality, suppose $\{1, 2\}$ can be used to color X_3 . Then by letting $c'(v_2) = 3, c'(v_3) = 5, c'(v_1) = 6, c'(v_4) = 4, c'(v_i) = l_i$, here $l_i \in \{1, 2\} \cap L(v_i), i \in \{5, 6, \dots, 12\}$, we obtain another L -coloring c' of $K_{2,2,8}$ (which is exhibited below), a contradiction.

$$\begin{array}{cc} \underline{156} & \underline{234} \\ 35\underline{1} & 46\underline{1} \end{array}$$

Subcase 3.4 (1)

$$\begin{array}{cc} \underline{156} & \underline{234} \\ 35\underline{1} & 46\underline{1} \end{array}$$

Subcase 3.4 (2)

Consequently, $K_{2,2,8}$ has property $M(3)$. Then, we can easily prove $K_{2,2,r}$, $r \in \{4, 5, 6, 7\}$ have property $M(3)$. For otherwise, assume $K_{2,2,r}$ is U3LC for some $r \in \{4, 5, 6, 7\}$. Since $K_{2,2,r}$, $r \in \{4, 5, 6, 7\}$ is an induced subgraph of $K_{2,2,8}$, by Lemma 2.3, $K_{2,2,8}$ is U3LC, a contradiction. So, $K_{2,2,r}$, $r \in \{4, 5, 6, 7\}$ have property $M(3)$. This completes the proof of Theorem 2.1. ■

Corollary 2.1. $m(K_{2,2,r}) = 3$, $r \in \{4, 5, 6, 7, 8\}$.

Proof. It follows from Theorem 1.1 and Theorem 2.1. ■

Now all the graphs in Problem 1 are proved to have the property $M(3)$. Therefore, we can completely characterize the U3LC complete multipartite graphs as follows.

Theorem 2.2. *A complete multipartite graph is U3LC if and only if it has one of the ten graphs $K_{3,3,3}$, $K_{2,4,4}$, $K_{2,3,5}$, $K_{2,2,9}$, $K_{1,2,2,2}$, $K_{1,1,2,3}$, $K_{1,1,1,2,2}$, $K_{1*4,6}$, $K_{1*5,5}$, and $K_{1*6,4}$ as an induced subgraph.*

Proof. It follows from Theorem 1.3, Theorem 1.4, Theorem 1.5, Theorem 1.6 and Theorem 2.1. ■

Acknowledgements

Comments of the anonymous referee have been very helpful to us when revising the paper.

REFERENCES

- [1] N. Alon, *Restricted colorings of graphs*, in: K. Walker, editor, *Surveys in Combinatorics*, Number 187 in London Math. Soc. LNS, pp. 1–33, 1993.
- [2] J.A. Bondy and U.S.R. Murty, *Graph Theory with Applications* (American Elsevier Publishing Co., INC., New York, 1976).
- [3] J.H. Dinitz and W.J. Martin, *The stipulation polynomial of a uniquely list colorable graph*, *Austral. J. Combin.* **11** (1995) 105–115.

- [4] P. Erdős, A.L. Rubin and H. Taylor, Choosability in graphs, in: Proceedings of West Coast Conference on Combinatorics, Graph Theory and Computing, number 26 in Congr. Number., pp. 125–157, Arcata, CA, September 1979.
- [5] Y.G. Ganjali, M. Ghebleh, H. Hajiabohassan, M. Mirzadeh and B.S. Sadjad, *Uniquely 2-list colorable graphs*, Discrete Appl. Math. **119** (2002) 217–225.
- [6] M. Ghebleh and E.S. Mahmoodian, *On uniquely list colorable graphs*, Ars Combin. **59** (2001) 307–318.
- [7] W.J. He, Y.N. Wang, Y.F. Shen and X. Ma, *On property $M(3)$ of some complete multipartite graphs*, Australasian Journal of Combinatorics, to appear.
- [8] M. Mahdian and E.S. Mahmoodian, *A characterization of uniquely 2-list colorable graphs*, Ars Combin. **51** (1999) 295–305.
- [9] E.S. Mahmoodian and M. Mahdian, *On the uniquely list colorable graphs*, in: Proceedings of the 28th Annual Iranian Mathematics Conference, Part 1, number 377 in Tabriz Univ. Ser., Tabriz, 1997.
- [10] Y.F. Shen and Y.N. Wang, *On uniquely list colorable complete multipartite graphs*, Ars Combin. **88** (2008) 367–377.
- [11] V.G. Vizing, *Coloring the vertices of a graph in prescribed colors*, (in Russian) Discret. Anal. **29** (1976) 3–10.
- [12] Y.Q. Zhao, W.J. He, Y.F. Shen and Y.N. Wang, *Note on characterization of uniquely 3-list colorable complete multipartite graphs*, in: Discrete Geometry, Combinatorics and Graph Theory, LNCS **4381** (Springer, Berlin, 2007) 278–287.

Received 26 February 2009

Revised 2 April 2009

Accepted 14 April 2009