

## THE PROJECTIVE PLANE CROSSING NUMBER OF THE CIRCULANT GRAPH $C(3k; \{1, k\})$

PAK TUNG HO

*Department of Mathematics, Sogang University,  
Seoul 121-742, Korea*

**e-mail:** ptho@sogang.ac.kr

### Abstract

In this paper we prove that the projective plane crossing number of the circulant graph  $C(3k; \{1, k\})$  is  $k - 1$  for  $k \geq 4$ , and is 1 for  $k = 3$ .

**Keywords:** crossing number, circulant graph, projective plane.

**2010 Mathematics Subject Classification:** 05C10.

### 1. INTRODUCTION

The crossing number is an important measure of the non-planarity of a graph. Bhatt and Leighton [1] showed that the crossing number of a network (graph) is closely related to the minimum layout area required for the implementation of a VLSI circuit for that network. In general, determining the crossing number of a graph is hard. Garey and Johnson [3] showed that it is NP-complete. In fact, Hliněný [6] has proved that the problem remains NP-complete even when restricted to cubic graphs. Moreover, the exact crossing number is not known even for specific graph families, such as complete graphs [16], complete bipartite graphs [11, 22], and circulant graph [8, 12, 13, 14, 20, 23]. For more about crossing number, see [2, 21] and references therein.

Attention has been paid to the crossing number of graphs on surfaces [4, 5, 7, 9, 10, 17, 18, 19]. However, exact values are known only for very restricted classes of graphs. In this paper, we compute the projective plane crossing number of the circulant graph  $C(3k; \{1, k\})$ .

**Theorem 1.** *The projective plane crossing number of the circulant graph  $C(3k; \{1, k\})$  is given by*

$$cr_1(C(3k; \{1, k\})) = \begin{cases} k - 1 & \text{for } k \geq 4, \\ 1 & \text{for } k = 3. \end{cases}$$

Note that there are only few infinite classes of graphs whose projective plane crossing number are known exactly. See [9, 19].

Here are some definitions. Let  $G$  be a simple graph with the vertex set  $V = V(G)$  and the edge set  $E = E(G)$ . The *circulant graph*  $C(n; S)$  is the graph with the vertex set  $V(C(n; S)) = \{v_i \mid 1 \leq i \leq n\}$  and the edge set  $E(C(n; S)) = \{v_i v_j \mid 1 \leq i, j \leq n, (i-j) \bmod n \in S\}$  where  $S \subseteq \{1, 2, \dots, \lfloor n/2 \rfloor\}$ .

The *projective plane crossing number*  $cr_1(G)$  of  $G$  is the minimum number of crossings of all the drawings of  $G$  in the projective plane having the following properties: (i) no edge has a self-intersection; (ii) no two adjacent edges intersect; (iii) no two edges intersect each other more than once; (iv) each intersection of edges is a crossing rather than tangential; and (v) no three edges intersect in a common point. Similarly one can define the plane crossing number  $cr(G)$  of the graph  $G$ . In a drawing  $D$ , if an edge (or a set of edges) does not cross other edges, we call it *clean*; otherwise, we call it *cross*. For a drawing  $D$ , the total number of crossings is denoted by  $v(D)$ .

Let  $A$  and  $B$  be two (not necessary disjoint) subsets of the edge set  $E$ . In a drawing  $D$ , the number of crossings crossed by an edge in  $A$  and another edge in  $B$  is denoted by  $v_D(A, B)$ . In particular,  $v_D(A, A)$  is denoted by  $v_D(A)$ , and hence  $v(D) = v_D(E)$ . By counting the number of crossings in  $D$ , we have the following:

**Lemma 2.** *Let  $A, B, C$  be mutually disjoint subsets of  $E$ . Then,*

$$(1) \quad \begin{aligned} v_D(A, B \cup C) &= v_D(A, B) + v_D(A, C), \\ v_D(A \cup B) &= v_D(A) + v_D(B) + v_D(A, B). \end{aligned}$$

The plan of this paper is as follows. In Section 2 we prove the upper bound of the projective crossing number of  $C(3k; \{1, k\})$ . In Section 3, we prove the lower bound of the projective crossing number of  $C(3k; \{1, k\})$  by assuming Lemma 7. In Section 4, we prove Lemma 7, which says that for any drawing of  $C(3k; \{1, k\})$  with all of its cycles being clean, its number of crossing is at least  $k - 1$ .

## 2. UPPER BOUNDS



From now on, we will denote the circulant graph  $C(3k; \{1, k\})$  by  $C(k)$  for simplicity. First we have the following:

**Lemma 3.**  $cr_1(C(3)) \leq 1$ .

**Proof.** One can refer to the drawing of  $C(3)$  in the projective plane in Figure 1. ■

**Lemma 4.**  $cr_1(C(k)) \leq k - 1$  for  $k \geq 4$ .

**Proof.** For a non-planar graph  $G$ , the plane crossing number is strictly greater than the projective plane crossing number, i.e.,  $cr_1(G) \leq cr(G) - 1$ . Lemma 4 follows from  $cr(C(k)) = k$  for  $k \geq 4$ , which is proved in [12]. ■

Figure 1. Drawing of  $C(3)$ .Figure 2.  $F_1(9, 15)$ .

### 3. LOWER BOUNDS

Next, we have the following:

**Lemma 5.**  $cr_1(C(3)) \geq 1$ .

**Proof.** It suffices to show that  $C(3)$  cannot be embedded in the projective plane. Note that  $C(3) - \{v_1v_7, v_2v_8, v_3v_6\}$  is isomorphic to  $F_1(9, 15)$  (see Figure 2) in the list of the minimal forbidden subgraphs for the projective plane (see Appendix A in [15]). This shows that  $C(3)$  cannot be embedded in the projective plane. ■

In fact, we have shown the following:

**Corollary 6.** *If  $e$  is an edge in the cycle  $C_i$  (see the definition below) in  $C(3)$ , then  $cr_1(C(3) - e) \geq 1$ .*

In  $C(k)$ , we define

$$C_i = \{v_i v_{k+i}, v_i v_{2k+i}, v_{k+i} v_{2k+i}\},$$

where  $1 \leq i \leq k$ . We have the following:

**Lemma 7.** *For  $k \geq 4$ , let  $D$  be a drawing of  $C(k)$  such that  $C_i$  is clean for all  $1 \leq i \leq k$ . Then  $v(D) \geq k - 1$ .*

We postpone its proof to Section 4. By assuming Lemma 7, we are in a position to prove the lower bound of  $cr_1(C(k))$ .

**Lemma 8.**

$$(2) \quad cr_1(C(k)) \geq k - 1 \text{ for } k \geq 4.$$

**Proof.** We will prove (2) by induction on  $k$ . First consider  $k = 4$ . Suppose  $D$  is a drawing of  $C(4)$ . We will prove  $v(D) \geq 3$  by contradiction. Suppose that  $v(D) \leq 2$ . Then there exists  $C_i$  which crosses; otherwise, if all  $C_i$  are clean,  $v(D) \geq 3$  by Lemma 7.




Figure 3(a)




Figure 3(b)




Figure 3(c)

Without loss of generality, we may assume that the edge  $v_1v_5$  in  $C_1$  crosses. Then there exists an edge  $e$  in  $D - v_1v_5$  such that  $D - v_1v_5 - e$  is an embedding in the projective plane. Note that  $e$  cannot be the edge in any cycle  $C_1$ : If  $e$  is an edge in  $C_1$  other than  $v_1v_5$ , then  $D - C_1$ , which is a subdivision of  $C(3)$ , is an embedding in the projective plane, which is impossible by Lemma 5. If  $e$  is an edge in  $C_i$  with  $i \neq 1$ , then  $D - C_1 - e$ , which is a subdivision of  $C(3)$  minus an edge in the cycle  $C^i$  is an embedding in the projective plane, which contradicts Corollary 6.

Therefore, by symmetry, we have the following possibilities:  $e = v_2v_3$ ,  $e = v_4v_5$ ,  $e = v_5v_6$ ,  $e = v_6v_7$ ,  $e = v_7v_8$ ,  $e = v_8v_9$ . We will show that it is impossible for  $C(4) - v_1v_5 - e$  to be embedded in the projective plane for each of these cases, which will give the required contradiction.

First, by contracting the edges  $v_5v_6$  and  $v_7v_8$  in  $C(4) - \{v_1v_5, v_4v_5, v_8v_9\}$ , we get a graph which contains a subgraph isomorphic to  $F_4(10,16)$  (see Figure 3(a)) in the list of the minimal forbidden subgraphs for the projective plane (see Appendix A in [15]). Moreover, by contracting the edges  $v_3v_4$  and  $v_5v_6$  in  $C(4) - \{v_1v_5, v_2v_3, v_6v_7\}$ , we get a graph which contains a subgraph isomorphic to  $F_4(10,16)$  (see Figure 3(b)).




Figure 4(a)




Figure 4(b)




Figure 5

Next we are going to show that  $C(4) - \{v_1v_5, v_5v_6\}$  cannot be embedded in the projective plane. Suppose it is not true and let  $D$  be an embedding of  $C(4) - \{v_1v_5, v_5v_6\}$  in the projective plane. Delete the edge  $v_2v_6$  in the drawing. Since  $v_1v_5$  and  $v_5v_6$  are absent, we can always draw an edge connecting  $v_4$  and  $v_9$  which is close to the edges  $v_4v_5$  and  $v_5v_9$  without producing new crossings (see Figure 4(a)). Similarly, since  $v_2v_6$  and  $v_5v_6$  are absent, we can draw an edge connecting  $v_7$  and  $v_{10}$  which is close to the edges  $v_6v_7$  and  $v_6v_{10}$  without producing new crossings (see Figure 4(b)). Therefore, we obtain an embedding of  $C(12, \{1, 4\}) - \{v_1v_5, v_5v_6, v_2v_6\} + \{v_4v_9, v_7v_{10}\}$  in the projective plane, which is impossible since it contains a minor isomorphic to  $F_4(10, 16)$  (see Figure 3(c)).

Finally, one can see that  $C(12, \{1, 4\}) - \{v_1v_5, v_7v_8\}$  contains a minor isomorphic to  $F_5(10, 16)$  (see Figure 5) in the list of the minimal forbidden subgraphs for the projective plane (see Appendix A in [15]).

Therefore, (2) is true for  $k = 4$ . Now suppose that (2) is true for all values less than  $k \geq 5$ . Let  $D$  be a drawing of  $C(k)$  in the projective plane and we are going to show that  $v(D) \geq k - 1$ .

If there exists  $1 \leq i \leq 3k$  such that  $v_iv_{k+i}$  crosses, then by deleting  $v_iv_{k+i}$ ,  $v_{k+i}v_{2k+i}$ ,  $v_{2k+i}v_i$ , we obtain a drawing of a subdivision of  $C(k-1)$ , denote it by  $D'$ . By our construction,  $v(D') \leq v(D) - 1$ . On the other hand,  $v(D') \geq k - 2$  by induction assumption. This implies  $v(D) \geq k - 1$ . Therefore, we may assume that  $v_iv_{k+i}$  is clean in  $D$  for all  $1 \leq i \leq 3k$ , i.e.,  $C_i$  is clean for all  $1 \leq i \leq k$ . Then by Lemma 7, we have  $v(D) \geq k - 1$ . ■

**Proof of Theorem 1.** It follows from Lemma 3, 4, 5 and 8. ■

#### 4. PROOF OF LEMMA 7

This section is devoted to proving Lemma 7. Throughout this section, we assume that  $C_i$  is clean for  $1 \leq i \leq k$ , as we have assumed in Lemma 7.

For  $1 \leq i \leq k$ , let

$$F_i = \{v_i v_{k+i}, v_i v_{2k+i}, v_{k+i} v_{2k+i}, v_i v_{i+1}, v_{k+i} v_{k+i+1}, v_{2k+i} v_{2k+i+1}\}.$$

Note that the set of all  $F_i$  is a partition of the edge set  $E$  of  $C(k)$ , i.e.,

$$(3) \quad E = \bigcup_{i=1}^k F_i \text{ and } F_i \cap F_j = \emptyset \text{ for } i \neq j.$$

For  $1 \leq i \leq k$ , define

$$(4) \quad f_D(F_i) = v_D(F_i) + \frac{1}{2} \sum_{j \neq i} v_D(F_i, F_j).$$

Since we have assumed that each  $C_i$  is clean, there are only two possible ways of drawing  $C_i$ , depending on whether it is contractible or not, which are shown in Figure 6(a) and 6(b).

If  $C_i$  and  $C_{i+1}$  are both contractible, there are three possible ways of drawing  $C_i \cup C_{i+1}$  for each  $i$ , which are shown in Figure 7(a), 7(b) and 7(c).




Figure 6(a).  $C_i$  is contractible.




Figure 6(b).  $C_i$  is non-contractible.

We have the following:

**Proposition 9.** *If  $C_i$  and  $C_{i+1}$  are drawn as in Figure 7(a) or 7(b), then*

$$f_D(F_i) \geq 1.$$

**Proof.** Suppose  $f_D(F_i) < 1$ . By (4),  $v_i v_{i+1}, v_{k+i} v_{k+i+1}, v_{2k+i} v_{2k+i+1}$  do not cross each other. If  $C_i \cup C_{i+1}$  is drawn as in Figure 7(a),  $F_i \cup C_{i+1}$  must be drawn as in Figure 8 since  $C_i, C_{i+1}$  are clean and  $v_i v_{i+1}, v_{k+i} v_{k+i+1}, v_{2k+i} v_{2k+i+1}$  do not cross each other. Since  $C_{i-1}$  is clean,  $C_{i-1}$  must lie entirely in one of the regions  $f_1, f_2$  or  $f_3$ . We may assume that  $C_{i-1}$  lies in the region  $f_1$ , for other cases the proof is the same. If  $C_{i-1}$  lies in  $f_1$ , then  $v_{k+i-1} v_{k+i}$  must cross  $v_i v_{i+1}$  or  $v_{2k+i} v_{2k+i+1}$ . On the other hand, the path  $v_{k+i+1} v_{k+i+2} \cdots v_{2k-i-1}$  must cross  $v_i v_{i+1}$  or  $v_{2k+i} v_{2k+i+1}$ . Hence, by (4),  $f_D(F_i) \geq 1$ . Similarly, one can show that  $f_D(F_i) \geq 1$  if  $C_i \cup C_{i+1}$  is drawn as in Figure 7(b). ■




Figure 7(a)




Figure 7(b)




Figure 7(c)

**Proposition 10.** *If  $C_i \cup C_{i+1}$  is drawn as in Figure 7(c) and  $f_D(F_i) < 1$ , then  $F_i \cup C_{i+1}$  must be drawn as in Figure 9(b).*

**Proof.** Since  $f_D(F_i) < 1$ , by (4),  $v_{k+i}v_{k+i+1}, v_{2k+i}v_{2k+i+1}$  do not cross each other. Then  $F_i \cup C_{i+1}$  must be drawn as in Figure 9(a) or 9(b). If  $F_i \cup C_{i+1}$  is drawn as in Figure 9(a), then  $C_{i-1}$  must lie entirely in one of the regions  $f_1, f_2$  or  $f_3$  since  $C_{i-1}$  is clean. We may assume that  $C_{i-1}$  lies in the region  $f_1$ , for other cases the proof is the same. If  $C_{i-1}$  lies in  $f_1$ , then  $v_{i-1}v_i$  must cross  $v_{k+i}v_{k+i+1}$  or  $v_{2k+i}v_{2k+i+1}$  since  $C_i$  and  $C_{i+1}$  are clean. On the other hand, the path  $v_{i+1}v_{i+2} \cdots v_{k-i-1}$  must cross  $F_i$ . Hence, by (4), we have  $f_D(F_i) \geq 1$ , which contradicts that  $f_D(F_i) < 1$ . ■




Figure 8




Figure 9(a)




Figure 9(b)

Combining Proposition 9 and 10, we have the following:

**Corollary 11.** *If  $F_i \cup C_{i+1}$  is not drawn as in Figure 9(b), then  $f_D(F_i) \geq 1$ .*

**Proof.** By Proposition 10, either  $f_D(F_i) \geq 1$  or  $C_i \cup C_{i+1}$  is not drawn as in Figure 7(c). In the latter case,  $C_i \cup C_{i+1}$  must be drawn as in Figure 7(a) or 7(b). By Proposition 9, again we have  $f_D(F_i) \geq 1$ . ■

**Remark 12.** Hereafter, we say that  $F_j \cup C_{j+1}$  is drawn as in Figure 9(b) if it is drawn as in Figure 9(c), i.e., replacing all the indices  $i$  by  $j$ .




Figure 9(c)






Figure 9(d)

Figure 10.  $F_i \cup C_{i+1} \cup F_j \cup C_{j+1}$ .Figure 11.  $F_1 \cup F_2 \cup C_3$ .

**Proposition 13.** Suppose that  $F_i \cup C_{i+1}$  is drawn as in Figure 9(b). If  $j \neq i-1, i, i+1$  such that  $F_j \cup C_{j+1}$  is drawn as in Figure 9(b), then  $F_i$  and  $F_j$  must cross each other. In particular, we have  $f_D(F_i) \geq 1/2$  and  $f_D(F_j) \geq 1/2$ .

**Proof.** Note that two non-contractible curves in the projective plane must cross each other. Since  $F_i \cup C_{i+1}$  and  $F_j \cup C_{j+1}$  are drawn as in Figure 9(b) where  $j \neq i-1, i, i+1$ ,  $F_i$  and  $F_j$  must cross each other since  $C_i, C_{i+1}, C_j, C_{j+1}$  are clean. See Figure 10 for a possible drawing of  $F_i \cup C_{i+1} \cup F_j \cup C_{j+1}$ . Since  $F_i$  and  $F_j$  cross each other, we have  $v_D(F_i, F_j) \geq 1$ , which implies that  $f_D(F_i) \geq 1/2$  and  $f_D(F_j) \geq 1/2$  by (4). ■



Here is the outline of the proof of Lemma 7. We will consider two cases:

*Case 1.*  $C_i$  is contractible for all  $1 \leq i \leq k$ .

*Case 2.*  $C_i$  is non-contractible for some  $1 \leq i \leq k$ .

For Case 1, by simple arguments, we can show that  $F_1 \cup C_2$  is drawn as in Figure 9(b). Moreover, we can show that  $f_D(F_{i_0}) < 1$  for some  $i_0 \neq 1$ . Then we will consider two cases:

*Case 1.1.*  $i_0 \neq 2, k$ .

*Case 1.2.*  $i_0 = 2$  or  $k$ .

Case 1.1 can be solved easily. For Case 1.2, we will assume that  $i_0 = 2$  since the proof for  $i_0 = k$  is the same. Then we will consider two cases:

*Case 1.2.1.*  $f_D(F_j) \geq 1$  for all  $j \neq 1, 2$ .

*Case 1.2.2.*  $f_D(F_j) < 1$  for some  $j \neq 1, 2$ .

For Case 1.2.1, by assumption,  $f_D(F_j) \geq 1$  for all  $j \neq 1, 2$ . We just need to show that  $f_D(F_1) + f_D(F_2) > 0$ , which implies that  $v(D) = \sum_{j=1}^k f_D(F_j) = f_D(F_1) + f_D(F_2) + \sum_{j \neq 1, 2} f_D(F_j) > k - 2$ , and hence  $v(D) \geq k - 1$  since  $v(D)$  is an integer. For Case 1.2.2, by assumption,  $f_D(F_j) < 1$  for some  $j \neq 1, 2$ . Then we will consider two cases:

*Case 1.2.2.1.*  $j \neq 3, k$ .

*Case 1.2.2.2.*  $j = 3$  or  $k$ .

Case 1.2.2.1 can be solved easily.

For Case 1.2.2.2, we can assume that

$$(5) \quad f_D(F_l) \geq 1 \text{ for } l \neq 1, 2, 3, k.$$

Otherwise, if  $f_D(F_l) < 1$  for some  $l \neq 1, 2, 3, k$ , then it can be reduced to Case 1.2.2.1 by taking  $j = l$ . By simple arguments, we can reduce it to the case when both  $F_3 \cup C_4$  and  $F_k \cup C_1$  are drawn as in Figure 9(b). That is to say,  $F_i \cup C_{i+1}$  is drawn as in Figure 9(b) for  $i = 1, 2, 3, k$ . Then by Proposition 13,  $F_1$  crosses  $F_3$  and  $F_2$  crosses  $F_k$ . Moreover, if  $k \geq 5$ , then  $F_1$  also crosses  $F_k$ . All these implies

$$(6) \quad f_D(F_1) \geq 1, f_D(F_k) \geq 1, f_D(F_2) \geq 1/2, \text{ and } f_D(F_3) \geq 1/2.$$

Combining (5) and (6), we get  $v(D) \geq k - 1$ . For  $k = 4$ , we will use different arguments by making use the fact that  $F_i \cup C_{i+1}$  is drawn as in Figure 9(b) for  $i = 1, 2, 3, 4$ .

Now we are ready to prove Lemma 7.

**Proof of Lemma 7.** By (1), (3) and (4), the total number of crossing of the drawing  $D$  is  $v(D) = v_D(E) = \sum_{i=1}^k f_D(F_i)$ . Therefore, it suffices to prove that  $\sum_{i=1}^k f_D(F_i) \geq k - 1$ . To prove by contradiction, we assume that

$$(7) \quad \sum_{i=1}^k f_D(F_i) < k - 1.$$

We will consider two cases: Case 1.  $C_i$  is contractible for all  $1 \leq i \leq k$  and Case 2.  $C_i$  is non-contractible for some  $1 \leq i \leq k$ .

*Case 1.* Since we have assumed that  $C_i$  is clean for  $1 \leq i \leq k$ , as we have said at the beginning of this section, there are three possible ways of drawing  $C_i \cup C_{i+1}$  for each  $i$ , which are shown in Figure 7(a), 7(b) or 7(c).

Note that (7) implies that  $f_D(F_i) < 1$  for some  $i$ . Without loss of generality, we may assume  $i = 1$ , i.e.,

$$(8) \quad f_D(F_1) < 1.$$

By Proposition 9,  $C_1 \cup C_2$  must be drawn as in Figure 7(c). Hence, by (8) and Proposition 10,  $F_1 \cup C_2$  is drawn as in Figure 9(b) (see Figure 9(d)).

There exists  $i_0 \neq 1$  such that  $F_{i_0} \cup C_{i_0+1}$  is drawn as in Figure 9(b). (Otherwise, if  $F_j \cup C_{j+1}$  is not drawn as in Figure 9(b) for all  $j \neq 1$ ,  $f_D(F_j) \geq 1$  for all  $j \neq 1$  by Corollary 11, which implies  $\sum_{j=1}^k f_D(F_j) \geq \sum_{j \neq 1} f_D(F_j) \geq k - 1$ .) We will consider two cases: Case 1.1.  $i_0 \neq 2, k$  and Case 1.2.  $i_0 = 2$  or  $k$ .

*Case 1.1.* If  $i_0 \neq 2, k$ , i.e.,  $C_{i_0} \cup C_{i_0+1}$  is drawn as in Figure 9(b) for some  $i_0 \neq 1, 2, k$ , then by Proposition 13,  $F_1$  and  $F_{i_0}$  cross each others,

$$(9) \quad f_D(F_1) \geq 1/2 \text{ and } f_D(F_{i_0}) \geq 1/2.$$

Moreover, if there exists  $j \neq 1, 2, i_0, k$  such that  $f_D(F_j) < 1$ , then  $F_j \cup C_{j+1}$  must be drawn as in Figure 9(b) by Proposition 10. By Proposition 13,  $F_j$  and  $F_1$  must also cross each other. Hence,  $f_D(F_1) \geq 1$  since  $F_1$  crosses both  $F_{i_0}$  and  $F_j$ , which contradicts (8). Therefore,

$$(10) \quad f_D(F_j) \geq 1 \text{ for all } j \neq 1, 2, i_0, k.$$

Moreover, we can assume that

$$(11) \quad f_D(F_2) \geq 1 \text{ and } f_D(F_k) \geq 1.$$

(Otherwise,  $f_D(F_2) < 1$  or  $f_D(F_k) < 1$  implies that  $F_2 \cup C_3$  or  $F_k \cup C_1$  is drawn as in Figure 9(b) by Proposition 10. Replacing  $i_0$  by 2 or  $k$ , one can reduce this to Case 1.2.) Combining (9), (10) and (11), we have  $\sum_{j=1}^k f_D(F_j) \geq f_D(F_1) + f_D(F_{i_0}) + \sum_{j \neq 1, i_0} f_D(F_j) \geq k - 1$ .

*Case 1.2.* If  $i_0 = 2$  or  $k$ , then we may assume that  $i_0 = 2$  since the proof for  $i_0 = k$  is the same. Then  $F_2 \cup C_3$  is drawn as in Figure 9(b). We will consider two cases: *Case 1.2.1.*  $f_D(F_j) \geq 1$  for all  $j \neq 1, 2$  and *Case 1.2.2.*  $f_D(F_j) < 1$  for some  $j \neq 1, 2$ .

*Case 1.2.1.* By assumption,

$$(12) \quad f_D(F_j) \geq 1 \text{ for all } j \neq 1, 2.$$

If we can show that

$$(13) \quad f_D(F_1) + f_D(F_2) > 0,$$

then by (12) and (13),

$v(D) = \sum_{j=1}^k f_D(F_j) = f_D(F_1) + f_D(F_2) + \sum_{j \neq 1, 2} f_D(F_j) > k - 2$ , which implies that  $v(D) \geq k - 1$  since the total number of crossing  $v(D)$  is an integer.




Figure 12




Figure 13




Figure 14

Suppose (13) is not true, i.e.,

$$(14) \quad f_D(F_1) = f_D(F_2) = 0.$$

Recall that  $F_1 \cup C_2$  is drawn as in Figure 9(d). Since  $C_3$  is clean,  $C_3$  must lie entirely in regions  $f_1$  or  $f_2$  in Figure 9(d). If  $C_3$  lies in  $f_1$ , then  $v_2v_3$  must cross  $v_{k+1}v_{k+2}$  or  $v_{2k+1}v_{2k+2}$ . By (4),  $f_D(F_2) \geq 1/2$ , which contradicts (14). Therefore,  $C_3$  lies in  $f_2$ . By (4) and (14),  $v_2v_3$ ,  $v_{k+2}v_{k+3}$ ,  $v_{2k+2}v_{2k+3}$  are clean. Then the only possible drawing of  $F_1 \cup F_2 \cup C_3$  is shown as in Figure 11. (It is true up to renaming the vertices. For example, it is possible for  $F_1 \cup F_2 \cup C_3$  to be drawn as in Figure 12. But one can reduce it to Figure 11 by the transformation  $v_j \mapsto v_{j-k}$ .)

Since  $C_4$  is clean, it must lie entirely in one of the regions in Figure 11. Note that  $v_3$ ,  $v_{k+3}$  and  $v_{2k+3}$  do not lie in the the same region in Figure 11. No matter which region  $C_4$  lies in Figure 11, one of the edges  $v_3v_4$ ,  $v_{k+3}v_{k+4}$  and  $v_{2k+3}v_{2k+4}$  must cross the  $F_1$  or  $F_2$  (Note that  $k \geq 4$  is crucial here for  $C_4$  being not equal to  $C_1$ ). Hence,  $f_D(F_1) + f_D(F_2) > 0$  which gives (13).

*Case 1.2.2.* If  $f_D(F_j) < 1$  for some  $j \neq 1, 2$ , then  $F_j \cup C_{j+1}$  must be drawn as in Figure 9(b) by Proposition 10. We will consider two cases: Case 1.2.2.1.  $j \neq 3, k$  and Case 1.2.2.2.  $j = 3$  or  $k$ .

*Case 1.2.2.1.* Since  $F_j \cup C_{j+1}$  is drawn as in Figure 9(b) where  $j \neq 1, 2, 3, k$ ,  $F_j$  must cross  $F_1$  and  $F_2$  by Proposition 13, since  $F_1 \cup C_2$  and  $F_2 \cup C_3$  are drawn as in Figure 9(b). This implies that, by (4),

$$(15) \quad f_D(F_1) \geq 1/2, f_D(F_2) \geq 1/2, \text{ and } f_D(F_j) \geq 1.$$

Note that

$$(16) \quad f_D(F_r) \geq 1 \text{ for all } r \neq 1, 2, 3, j, k.$$

Otherwise, if  $f_D(F_r) < 1$  for some  $r \neq 1, 2, 3, j, k$ , then by Proposition 10,  $F_r \cup C_{r+1}$  is drawn as in Figure 9(b). By Proposition 13,  $F_r$  also crosses  $F_1$ . This implies  $f_D(F_1) \geq 1$  since  $F_1$  cross  $F_j$  and  $F_r$ , which contradicts (8).

We claim that

$$(17) \quad f_D(F_3) \geq 1 \text{ and } f_D(F_k) \geq 1.$$

To see this, suppose that  $f_D(F_3) < 1$ . Then  $F_3 \cup C_4$  is drawn as in Figure 9(b) by Proposition 10. Hence  $F_1$  must cross  $F_3$  and  $F_j$  by Proposition 13, which implies that  $f_D(F_1) \geq 1$  and contradicts (8). On the other hand, if  $f_D(F_k) < 1$ , then  $F_k \cup C_1$  must be drawn as in Figure 9(b) by Proposition 10. Hence  $F_2$  must cross  $F_k$  and  $F_j$  by Proposition 13, which implies that  $f_D(F_2) \geq 1$  and contradicts (8). This proves (17).

Combining (15), (16) and (17), we get  $\sum_{r=1}^k f_D(F_r) = f_D(F_1) + f_D(F_2) + \sum_{r \neq 1, 2} f_D(F_r) \geq k - 1$ .

*Case 1.2.2.2.* If  $j = 3$  or  $k$ , then  $F_k \cup C_1$  or  $F_3 \cup C_4$  is drawn as in Figure 9(b). We may assume that

$$(18) \quad f_D(F_l) \geq 1 \text{ for } l \neq 1, 2, 3, k.$$

(Otherwise, if  $f_D(F_l) < 1$  for some  $l \neq 1, 2, 3, k$ , then it can be reduces to Case 1.2.2.1 by taking  $j = l$ .) It can be reduced to the case when both  $F_3 \cup C_4$  and  $F_k \cup C_1$  are drawn as in Figure 9(b).

To see this, suppose that  $F_3 \cup C_4$  is drawn as in Figure 9(b) and  $F_k \cup C_1$  is not. Then by Corollary 11

$$(19) \quad f_D(F_k) \geq 1,$$

and  $F_3$  must cross  $F_1$  by Proposition 13 since  $F_1 \cup C_2$  is drawn as in Figure 9(b). We claim that  $F_1$  must cross  $F_k$ . Assuming the claim, we have

$$(20) \quad f_D(F_1) \geq 1 \text{ and } f_D(F_3) \geq 1/2.$$




Figure 15(a)




Figure 15(b)




Figure 15(c)




Figure 15(d)

Combining (18), (19) and (20), we get  $\sum_{r=1}^k f_D(F_r) > k - 2$ , which implies that  $v(D) = \sum_{i=1}^k f_D(F_i) \geq k - 1$  since  $v(D)$  is an integer.

To show the claim, i.e.,  $F_1$  crosses  $F_k$ , we note that  $F_1 \cup C_2$  is drawn as in Figure 9(b). See Figure 13. Since  $C_k$  is clean, it must lie entirely in one of the regions in Figure 13. It is impossible for  $C_k$  to lie in  $f_3$ , otherwise, the path  $v_2v_3 \cdots v_k$  crosses  $C_1$ . It is also impossible for  $C_{i-1}$  to lie in  $f_4$ , otherwise,  $v_kv_{k+1}$  crosses  $C_2$ . If  $C_k$  lies in  $f_1$ ,  $v_{3k}v_1$  must cross with  $v_{k+1}v_{k+2}$  or  $v_{2k+1}v_{2k+2}$ , which implies that  $F_k$  crosses  $F_1$ . If  $C_k$  lies in  $f_2$ , then  $F_k$  must cross  $F_1$  since  $F_k \cup C_1$  is not drawn as in Figure 9(b) by our assumption (See Figure 14 for example). Therefore,  $F_1$  must cross  $F_k$ , as we claimed.

Similarly, if  $F_k \cup C_1$  is drawn as in Figure 9(b) and  $F_3 \cup C_4$  is not, then  $\sum_{r=1}^k f_D(F_r) \geq k - 1$ .

Therefore, we can assume that both  $F_3 \cup C_4$  and  $F_k \cup C_1$  are drawn as in Figure 9(b). Then  $F_k$  must cross  $F_2$ , and  $F_1$  must cross with  $F_3$  by Proposition 13. Moreover, if  $k \geq 5$ , then  $F_3$  and  $F_k$  must also cross each other by Proposition 13. All these imply that

$$(21) \quad f_D(F_1) \geq 1/2, f_D(F_2) \geq 1/2, f_D(F_3) \geq 1, \text{ and } f_D(F_k) \geq 1.$$

Combining (18) and (21), we infer  $\sum_{r=1}^k f_D(F_r) \geq k - 1$  if  $k \geq 5$ .

On the other hand, if  $k = 4$ , then  $F_k \cup C_1 = F_4 \cup C_1$ ,  $F_1 \cup C_2$ ,  $F_2 \cup C_3$  and  $F_3 \cup C_4$  are drawn as in Figure 9(b) by assumptions. By Proposition 13,  $F_1$  must cross  $F_3$ , and  $F_2$  must cross  $F_4$ . This implies that

$$(22) \quad f_D(F_i) \geq 1/2 \text{ for } 1 \leq i \leq 4.$$

We will show that  $v(D) \geq 3$ . By contradiction, suppose that  $v(D) \leq 2$ . By (1) and (22), we have

$$(23) \quad f_D(F_1) = f_D(F_2) = f_D(F_3) = f_D(F_4) = 1/2.$$

Since  $F_1$  crosses  $F_3$ , by (4) and (23) we get

$$(24) \quad v_D(F_1, F_3) = 1, v_D(F_1, F_j) = 0 \text{ for } j \neq 3, v_D(F_3, F_j) = 0 \text{ for } j \neq 1.$$

Similarly, since  $F_2$  crosses  $F_4$ , by (4) and (23) we get

$$(25) \quad v_D(F_2, F_4) = 1, v_D(F_2, F_j) = 0 \text{ for } j \neq 4, v_D(F_4, F_j) = 0 \text{ for } j \neq 2.$$

Since  $F_1 \cup C_2$  and  $F_3 \cup C_4$  are drawn as in Figure 9(b), the only possible drawing of  $F_1 \cup C_2 \cup F_3 \cup C_4$  is shown in Figure 15(a) in view of (24) and (25). However, one can show that it is impossible for (24), (25) to hold. For example, if  $F_1 \cup C_2 \cup F_3 \cup C_4$  is drawn in Figure 15(b), then the edge  $v_8v_9$  must cross with  $F_1$  or  $F_3$ , which contradicts (24); and if  $F_1 \cup C_2 \cup F_3 \cup C_4$  is drawn in Figure 15(c), then the edge  $v_2v_3$  must lie entirely in the region  $f$ , as in Figure 15(d), since  $v_D(F_2, F_j) = 0$  for  $j \neq 4$  by (25). However, in Figure 15(d), no matter how  $v_6v_7$

is drawn,  $v_6v_7$  must either (i) cross  $v_2v_3$  which contradicts (25), or (ii) cross  $C_i$  which contradicts that  $C_i$  are all clean, or (iii) cross  $F_1$  or  $F_3$  which contradicts (25). We leave other cases to the reader.

*Case 2.* If there exists  $1 \leq i \leq k$  such that  $C_i$  is non-contractible, then we may assume that  $C_1$  is non-contractible. Then  $C_i$  is contractible for all  $i \neq 1$ . (Otherwise,  $C_i$  crosses  $C_1$  since two non-contractible curves in the projective plane must cross each other. This contradicts the assumption that all  $C_i$  are clean.) Since  $C_i$  and  $C_{i+1}$  are clean and contractible for  $i \neq 1, k$ , there are three possible ways of drawing  $C_i \cup C_{i+1}$ , which are shown in Figure 7(a), 7(b) or 7(c).

We claim that

$$(26) \quad f_D(F_i) \geq 1 \text{ for } i \neq 1, k.$$

To prove this, suppose that  $f_D(F_i) < 1$  for some  $i \neq 1, k$ . By Corollary 11,  $F_i \cup C_{i+1}$  must be drawn as in Figure 9(b), which crosses the non-contractible  $C_1$ . This contradicts that  $C_1$  is clean. This proves (26).

Now we are going to show that

$$(27) \quad f_D(F_1) + f_D(F_k) > 0.$$




Figure 16.  $C_1 \cup C_2$ .

Combining this with (26), we will get  $\sum_{r=1}^k f_D(F_r) > k - 2$ , which gives  $v(D) = \sum_{i=1}^k f_D(F_i) \geq k - 1$  since  $v(D)$  is an integer. Suppose that (27) is not true, i.e.,

$$(28) \quad f_D(F_1) = f_D(F_k) = 0.$$

Since  $C_1$  is non-contractible and  $C_2$  is contractible,  $C_1 \cup C_2$  must be drawn as in Figure 16. On the other hand, by the same reasons,  $C_1 \cup C_k$  must be drawn as in Figure 16 by replacing  $C_2$  by  $C_k$ .

By (4) and (28),  $v_1v_2$ ,  $v_{k+1}v_{k+2}$ ,  $v_{2k+1}v_{2k+2}$  do not cross. From Figure 16, one can see that there are three possible ways of drawing  $F_1 \cup C_2$ , which are shown in Figure 17(a), 17(b) and 17(c).




Figure 17(a)




Figure 17(b)




Figure 17(c)

If  $F_1 \cup C_2$  is drawn as in Figure 17(b) and 17(c), then  $C_3$  must lie entirely in one of the regions since  $C_3$  is clean. Then  $F_2$  must cross with  $F_1$  since there is no region in Figure 17(b) or 17(c) containing all of the vertices  $v_2$ ,  $v_{k+2}$  and  $v_{2k+2}$ . This implies  $f_D(F_1) > 0$ , which contradicts (28).

Therefore,  $F_1 \cup C_2$  must be drawn as in Figure 17(a). By the same argument,  $F_k \cup C_1$  must be drawn as in Figure 17(a) by replacing  $C_2$  by  $C_k$ . Hence,  $F_k \cup F_1 \cup C_2$  must be drawn as in Figure 18(a) or 18(b) since  $F_1$  does not cross  $F_k$  by (28).




Figure 18(a)




Figure 18(b)

Note that  $C_3$  must lie in one of the regions in Figure 18(a) or 18(b). Since there exists no region in Figure 18(a) or 18(b) which contains all of the vertices  $v_2$ ,  $v_{k+2}$  and  $v_{2k+2}$ ,  $F_3$  must cross either  $F_k$  or  $F_1$  ( $k \geq 4$  is needed here for  $F_3$  being not equal to  $F_k$ ). This implies that  $f_D(F_1) > 0$  or  $f_D(F_k) > 0$ , which gives (27).

This finishes the the proof of Lemma 7.  $\blacksquare$



## REFERENCES

- [1] S.N. Bhatt and F.T. Leighton, *A framework for solving VLSI graph layout problems*, J. Comput. System Sci. **28** (1984) 300–343.
- [2] P. Erdős, and R.K. Guy, *Crossing number problems*, Amer. Math. Monthly **80** (1973) 52–58.
- [3] M.R. Garey and D.S. Johnson, *Crossing number is NP-complete*, SIAM J. Algebraic Discrete Methods **1** (1983) 312–316.
- [4] R.K. Guy and T.A. Jenkyns, *The toroidal crossing number of  $K_{m,n}$* , J. Combin. Theory **6** (1969) 235–250.
- [5] R.K. Guy, T. Jenkyns and J. Schaer, *The toroidal crossing number of the complete graph*, J. Combin. Theory **4** (1968) 376–390.
- [6] P. Hliněný, *Crossing number is hard for cubic graphs*, J. Combin. Theory (B) **96** (2006) 455–471.
- [7] P.T. Ho, *A proof of the crossing number of  $K_{3,n}$  in a surface*, Discuss. Math. Graph Theory **27** (2007) 549–551.
- [8] P.T. Ho, *The crossing number of  $C(3k+1; \{1, k\})$* , Discrete Math. **307** (2007) 2771–2774.
- [9] P.T. Ho, *The crossing number of  $K_{4,n}$  on the projective plane*, Discrete Math. **304** (2005) 23–34.
- [10] P.T. Ho, *The toroidal crossing number of  $K_{4,n}$* , Discrete Math. **309** (2009) 3238–3248.
- [11] D.J. Kleitman, *The crossing number of  $K_{5,n}$* , J. Combin. Theory **9** (1970) 315–323.
- [12] X. Lin, Y. Yang, J. Lu and X. Hao, *The crossing number of  $C(mk; \{1, k\})$* , Graphs Combin. **21** (2005) 89–96.
- [13] X. Lin, Y. Yang, J. Lu and X. Hao, *The crossing number of  $C(n; \{1, \lfloor n/2 \rfloor - 1\})$* , Util. Math. **71** (2006) 245–255.
- [14] D. Ma, H. Ren and J. Lu, *The crossing number of the circular graph  $C(2m+2, m)$* , Discrete Math. **304** (2005) 88–93.
- [15] B. Mohar and C. Thomassen, *Graphs on Surfaces* (Johns Hopkins University Press, Baltimore, 2001).
- [16] S. Pan and R.B. Richter, *The crossing number of  $K_{11}$  is 100*, J. Graph Theory **56** (2007) 128–134.
- [17] R.B. Richter and J. Širáň, *The crossing number of  $K_{3,n}$  in a surface*, J. Graph Theory **21** (1996) 51–54.
- [18] A. Riskin, *The genus 2 crossing number of  $K_9$* , Discrete Math. **145** (1995) 211–227.
- [19] A. Riskin, *The projective plane crossing number of  $C_3 \times C_n$* , J. Graph Theory **17** (1993) 683–693.

- [20] G. Salazar, *On the crossing numbers of loop networks and generalized Petersen graphs*, Discrete Math. **302** (2005) 243–253.
- [21] L.A. Székely, *A successful concept for measuring non-planarity of graphs: the crossing number*, Discrete Math. **276** (2004) 331–352.
- [22] D.R. Woodall, *Cyclic-order graphs and Zarankiewicz’s crossing number conjecture*, J. Graph Theory **17** (1993) 657–671.
- [23] Y. Yang, X. Lin, J. Lu and X. Hao, *The crossing number of  $C(n; \{1, 3\})$* , Discrete Math. **289** (2004) 107–118.

Received 2 September 2010

Revised 26 January 2011

Accepted 26 January 2011